

Digitized by the Internet Archive
in 2015

<https://archive.org/details/b21917528>

A KEY

TO

Organic Materia Medica,

WRITTEN FOR THE STUDENTS OF
THE SOUTH-LONDON SCHOOL OF PHARMACY.

BY

DR. JOHN MUTER, M.A., F.C.S.,
Director of the School;

VICE-PRESIDENT OF THE SOCIETY OF PUBLIC ANALYSTS;
AUTHOR OF "PHARMACEUTICAL CHEMISTRY—THEORETICAL AND PRACTICAL," ETC.
PUBLIC ANALYST FOR LAMBETH, NEWINGTON, SOUTHWARK, BERMONDSEY
ROTHERHITHE, AND WANDSWORTH.



SECOND EDITION—REVISED & RE-ARRANGED.

LONDON:

PUBLISHED AT THE SCHOOL OFFICES, KENNINGTON CROSS, S.E.,
By WILLIAM BAXTER, *Secretary*;

SOLD BY MESSRS. SIMPKIN & MARSHALL, STATIONERS' HALL COURT, E.C.; AND
MESSRS. BAILLIERE, TINDAL & COX, KING WILLIAM STREET, STRAND, W.C.

MDCCCLXXVIII.

All Rights Reserved.]

London:

J. DAVIS, Printer, Steam Works, Old Kent Road, S.E.



Preface to the Second Edition.

IN issuing this new and enlarged edition, I have thought it advisable to depart from the former arrangement—suited, as it was, only to our Museum—and to give the book the form usually adopted by writers on *Materia Medica*.

This has afforded me the opportunity of introducing short descriptions of the Classes, Sub-Classes, Orders, and Sub-Orders, both of the Vegetable and Animal Kingdoms; and the whole work is now strictly on the basis of the most modern natural systems of classification. While carrying out this change, however, the book is still rendered a guide to the working drawers of the Museum, by having them catalogued in a tabulated Appendix for use during recognition.

In the Chemical portion, which has been made as complete as space will allow, the researches of others—now become, as it were, stereotyped—have been followed in cases where I have not had the opportunity of repeating the experiments, or where my own observation has fully borne them out; but I have not hesitated to depart from the beaten track in certain points where special experience in the analysis of the drug in question has led me to form different conclusions. A distinct feature has been made of the separation and tests for active principles, so as to aid analysts in their work on drugs; and under the head of “Specially Distinctive Tests” will be found a sort of qualitative course, verified by experiments, which, if followed step by step, will lead to the detection of the article under examination. I purpose, in a future edition, to greatly extend these tables; but their arrangement, and the verification of those proposed by others, is a work of so much time, that I have not been able to carry out all I desired at present. In a word, the whole book has been practically re-written, and, I hope, has been materially improved in every respect.

The Therapeutical portion of the work has been

purposely kept down to the merest recognised generalities, as I consider that, beyond such a point, the inculcation of special views on therapeutics are best left in the hands of the lecturers; and thus the reading of the book by Students will in no way interfere with the doctrines inculcated in their respective Schools of Medicine and Pharmacy.

I have appended some directions on the microscopic mounting and observation of the chief starches of articles of *Materia Medica* and food, in the form of a qualitative course for their detection. This system, originally made public by me in the "*Analyst*" for December, 1876, is one which I elaborated, and have for many years used in the teaching of Microscopy as applied to drugs. It was specially adapted for use with any ordinary microscope, and with no very excellent arrangement of light, so as to enable an ordinary observer to make tolerably shrewd conclusions. The measurements are the result of the carefully-collated results of the examination of over 1000 samples, and may be relied upon as good averages, disregarding extremes. It was my intention to have published illustrations of each of the drugs referred to in the Appendix; but through the

sudden exhaustion of the first edition, I have been compelled to publish without them.

I have to acknowledge the valuable assistance of my friend, Mr. JOSEPH INCE, in revising the proof sheets and aiding in securing the correctness of the typography, as well as in preparing the Index, which will be found very complete. The annotated Catalogue to the working drawers of the Museum is supplied by our *Materia Medica* Demonstrator and Curator, Mr. DODD.

J. M.

*South-London School of Pharmacy,
March, 1878.*



TABLE OF CONTENTS.

The Vegetable Kingdom.

	PAGE.
SUB-KINGDOM I.—PHANEROGAMIA.....	I—417
CLASS I.—DICOTYLEDONES	I—378
DIVISION I.—ANGIOSPERMIA	I—364
SUB-CLASS I.—THALAMIFLORÆ	I—100
<i>Natural Order</i> —Ranunculaceæ	I— 18
Aconiti Folia	2
Aconiti Radix.....	2
<i>Aconitum Napellus.</i>	
Aconiti Radix Indica	9
<i>Aconitum ferox.</i>	
Podophylli Radix	10
<i>Podophyllum peltatum.</i>	
Hellebori Nigri Radix	13
<i>Helleborus niger.</i>	
Staphisagriæ Semina	15
<i>Delphinium Staphisagria.</i>	
Actææ Racemosæ Radix, <i>vel.</i> Cimifugæ Radix	
<i>Actæa racemosa.</i>	

	PAGE.
Hydrastis Radix	18
<i>Hydrastis Canadensis.</i>	
Natural Order—Magnoliaceæ	18— 21
Winteri Cortex, <i>vel.</i> Cortex Winteranus	19
<i>Drimys Winteri.</i>	
Anisi Stellati Fructus	20
<i>Illicium anisatum.</i>	
Natural Order—Menispermaceæ.....	21— 29
Calumbæ Radix	21
<i>Jateorhiza palmata.</i>	
Pareiræ Radix	24
<i>Chondodendron tomentosum.</i>	
Cocculus Indicus	27
<i>Anamirta Cocculus.</i>	
Gulancha.....	29
<i>Tinospora cordifolia.</i>	
Natural Order—Berberidaceæ	29— 30
Natural Order—Papaveraceæ	30— 57
Rhœados Petala.....	30
<i>Papaver Rhœas.</i>	
Papaveris Capsulæ	32
<i>Papaver somniferum.</i>	
Papaveris Semina	34
Opium (Papaver somniferum).....	34
„ Smyrna.....	36
„ Constantinople	36
„ Egyptian	37
„ Persian	37
„ East Indian	37

	PAGE.
Opium Assay (B.P., &c.)	38— 39
Morphia, Morphine	41
Morphiæ Hydrochloras	44
„ Acetas	46
Apomorphia	48
Codeia, Codeine.....	48
Narceia, Narceine.....	49
Meconia, Meconine	49
Narcotia (Anarcotine)	50
Thebaia, Thebaine	50
Papaverine	51
Meconic Acid.....	51
Opium Preparations, Various	51— 56
Sanguinariæ Radix	56
<i>Sanguinaria Canadensis.</i>	
Natural Order—Cruciferæ, vel. Brassicaceæ	57— 62
Armoraciæ Radix	57
<i>Cochlearia Armoracia.</i>	
Sinapis Albæ Semina	59
<i>Sinapis alba.</i>	
Sinapis Nigræ Semina	59
<i>Sinapis nigra.</i>	
Natural Order—Canellaceæ	63
Canellæ Albæ Cortex	63
<i>Canella alba.</i>	
Natural Order—Polygalaceæ	64— 68
Senegæ Radix	64
<i>Polygala Senega.</i>	

	PAGE.
Krameriæ Radix	66
<i>Krameria triandra.</i>	
Natural Order—Guttiferæ	68— 71
Cambogia	68
Special Distinctive Tests	69
<i>Garcinia Morella</i> , var. β . <i>Pedicellata.</i>	
Garciniæ Oleum	71
<i>Garcinia Indica.</i>	
Natural Order—Dipterocarpeæ	71
Balsamum Dipterocarpi	71
<i>Dipterocarpus turbinatus.</i>	
Natural Order—Malvaceæ	72— 74
Althææ Radix	72
<i>Althæa officinalis.</i>	
Gossypium	74
Natural Order—Sterculiaceæ, vel. Byttneriaceæ....	74— 76
Theobromæ Semina	74
<i>Theobroma Cacao.</i>	
Theobromæ Oleum	75
Natural Order—Sapindaceæ	76
Guarana	76
<i>Paullinia sorbilis.</i>	
Natural Order—Linaceæ.....	77— 80
Lini Semina	77
<i>Linum usitatissimum.</i>	
Lini Oleum.....	79
Natural Order—Zygophyllaceæ	80— 84
Guaiaci Lignum	80
<i>Guaiacum officinale</i> , <i>G. sanctum.</i>	

	PAGE.
Guaiaci Resina	81
Special Distinctive Tests	83
<i>Natural Order—Rutaceæ</i>	84—90
Rutæ Oleum	85
<i>Ruta graveolens.</i>	
Cuspariæ, vel. Angosturæ Cortex	86
<i>Galipea Cusparia.</i>	
Buchu Folia	87
<i>Barosma betulina, B. crenulata et B. serratifolia.</i>	
Jaborandi	89
<i>Pilocarpus pinnatifolius.</i>	
<i>Natural Order—Aurantiaceæ</i>	91—98
Aurantii Cortex	91
<i>Citrus vulgaris, var. Bigaradia.</i>	
Aurantii Flores	93
<i>Citrus Bigaradia et C. Aurantium.</i>	
Aurantii Florum Oleum	94
Limonis Cortex	94
<i>Citrus limonum.</i>	
Limonis Oleum	95
Limonum Succus	96
Belæ Fructus	97
<i>Ægle Marmelos.</i>	
<i>Natural Order—Vitaceæ (Ampelidaceæ)</i>	98
Uvæ Passæ	98
<i>Vitis vinifera.</i>	
<i>Natural Order—Simarubaceæ</i>	99—101
Simarubæ Radicis Cortex	99
<i>Simaruba amara.</i>	

	PAGE.
Quassiæ Lignum	99
<i>Picræna excelsa.</i>	
SUB-CLASS II.—CALYCIFLORÆ	101—195
SUB-DIVISION I.—PERIGYNÆ	101—162
Natural Order—Rhamnaceæ	101—103
Rhamni Fructus	101
<i>Rhamnus cathartica.</i>	
Rhamni Succus	102
Frangulæ Cortex	103
<i>Rhamnus frangula.</i>	
Natural Order—Amyridaceæ (Burseraceæ)	104—110
Olibanum	104
<i>Boswellia</i> (species various).	
Elemi	105
Special Distinctive Tests.....	105
<i>Canarium commune.</i>	
Myrrha	106
Special Distinctive Tests	107
<i>Balsamodendron Myrrha, Bissa Böl.</i>	
Bdellium	109
Special Distinctive Tests.....	109
<i>Balsamodendron Mukul, B. Pubescens.</i>	
Natural Order—Anacardiaceæ.....	110—112
Mastiche, Mastich	110
Special Distinctive Tests.....	111
<i>Pistacia Lentiscus.</i>	
Natural Order—Leguminosæ, vel. Fabaceæ	113—147

	PAGE.
<i>Sub-Order I.—Papilionaceæ</i>	113—130
<i>Scoparii Cacumina</i>	113
<i>Sarothamnus Scoparius.</i>	
<i>Fœni-Græci Semina</i>	115
<i>Trigonella Fœnum-græcum.</i>	
<i>Tragacantha</i>	115—117
<i>Astragalus verus.</i>	
<i>Glycyrrhizæ Radix</i>	118—120
<i>Glycyrrhiza glabra.</i>	
<i>Indigo</i>	120
<i>Indigofera tinctoria.</i>	
<i>Setæ Mucunæ</i>	122
<i>Mucuna pruriens.</i>	
<i>Physostigmatis Fabæ</i>	122—124
<i>Physostigma venenosum.</i>	
<i>Kino</i>	124—125
<i>Pterocarpus Marsupium.</i>	
<i>Pterocarpi Lignum</i>	126
<i>Pterocarpus Santalinus.</i>	
<i>Balsamum Tolutanum</i>	126—128
<i>Special Distinctive Tests</i>	128
<i>Myroxylon Toluifera.</i>	
<i>Balsamum Peruvianum</i>	129
<i>Myroxylon Pereiræ.</i>	
<i>Sub-Order II.—Cæsalpinieæ</i>	130—143
<i>Hæmatoxyli Lignum</i>	130—132
<i>Hæmatoxylon Campechianum.</i>	
<i>Senna Alexandrina</i>	133—136
<i>Cassia acutifolia, C. lanceolata, C. obovata.</i>	

	PAGE.
Senna Indica	136
<i>Cassia elongata</i> (B.P.), <i>Cassia angustifolia</i> .	
Bombay or Mecca Senna	136
<i>Cassia angustifolia</i> (wild).	
Cassiæ Fistulæ Fructus	137
<i>Cassia Fistula</i> .	
Cassiæ Grandis Fructus	138
<i>Cassia grandis</i> , <i>Cassia Brasiliana</i> .	
Tamarindus	138—140
<i>Tamarindus Indica</i> .	
Copaiba	140—143
Special Distinctive Tests.....	141
<i>Copaifera multijuga</i> .	
Copaibæ Oleum	143
Sub-Order III.—Mimoseæ	143—147
Gummi Acaciæ	143—146
<i>Acacia</i> (species various).	
Catechu Nigrum	146
<i>Acacia Catechu</i> .	
Natural Order—Rosaceæ	147—162
Sub-Order I.—Chrysobalaneæ	147
Sub-Order II.—Amygdaleæ	148
Amygdalæ Dulces	148
<i>Amygdalus communis</i> , var. <i>dulcis</i> .	
Amygdalæ Amaræ.....	149
<i>Amygdalus communis</i> , var. <i>amara</i> .	
Oleum Amygdalæ Amaræ	150
Amygdalæ Oleum	152
Pruni Fructus	153
<i>Prunus domestica</i> .	

	PAGE.
Pruni Virginianæ Cortex	153
<i>Prunus serotina.</i>	
Lauro-cerasi Folia	154
<i>Prunus Lauro-cerasus.</i>	
Sub-Order III.—Roseæ	155—160
Cusso, Koso, <i>vel.</i> Kouso	155
<i>Hagenia Abyssinica, Brayera anthelmintica.</i>	
Rosæ Gallicæ Petala	156
<i>Rosa Gallica.</i>	
Rosæ Centifoliæ Petala	158
<i>Rosa centifolia.</i>	
Oleum Rosæ, Attar <i>vel.</i> Otto	159
Rosæ Caninæ Fructus	160
<i>Rosa canina.</i>	
Sub-Order IV.—Sanguisorbæ	161
Sub-Order V.—Pomeæ	161
Cydoniæ Semina	161
<i>Pyrus Cydonia.</i>	
SUB-DIVISION II.—EPIGYNÆ	162—195
Natural Order—Cucurbitaceæ.....	162—167
Sub-Orders—1, Nhandirobæ; 2, Siceæ; 3, Cucurbiteæ ..	162
Elaterium	162—165
<i>Ecballium officinarum, E. Elaterium.</i>	
Colocynthis Pulpa.....	165—167
<i>Citrullus Colocynthis.</i>	
Natural Order—Myrtaceæ	168—173
Cajuputi Oleum.....	168
<i>Melaleuca minor.</i>	
Eucalyptus Globulus	168
<i>et Eucalyptus resinifera.</i>	

	PAGE.
Caryophylli	169
<i>Caryophyllus aromaticus.</i>	
Pimenta	171
<i>Eugenia Pimenta.</i>	
Granati Radicis Cortex	172
<i>Punica Granatum.</i>	
Natural Order—Umbelliferæ, vel. Apiaceæ	174—195
Sub-Orders—1, Orthospermeæ; 2, Campylospermeæ;	
3, Cœlospermeæ.....	174
Conii Fructus.....	174—176
<i>Conium maculatum.</i>	
Conii Folia	176
Carui Fructus	178
<i>Carum Carui.</i>	
Carui Oleum	179
Fœniculi Fructus	179
<i>Fœniculum dulce.</i>	
Anisi Fructus.....	181
<i>Pimpinella Anisum.</i>	
Anethi Fructus	182
<i>Anethum graveolens.</i>	
Coriandri Fructus.....	183
<i>Coriandrum sativum.</i>	
Cumini Fructus.....	184
<i>Cuminum Cyminum.</i>	
Sumbul Radix	185
<i>Euryangium Sumbul.</i>	
Asafoetida	186
<i>Narthex Asa-fœtida.</i>	

	PAGE.
Galbanum	189—190
<i>Ferula Galbaniflua.</i> (?)	
Ammoniacum	191
<i>Dorema Ammoniacum.</i>	
Table for the Detection of the Umbelliferous Gum- Resins (Hirschsohn)	193—195
SUB-CLASS III.—COROLLIFLORÆ	195—309
SUB-DIVISION I.—EPIGYNÆ	196—242
Natural Order—Caprifoliaceæ	196
Sambuci Flores	196
<i>Sambucus nigra.</i>	
Natural Order—Cinchonaceæ (Rubiaceæ)	197—222
Catechu Pallidum	197
<i>Uncaria Gambier.</i>	
Cinchonæ Flavæ Cortex	200
<i>Cinchona Calisaya.</i>	
Flat Yellow Bark; Quill Yellow Bark	200—201
Tests—1, Qualitative; 2, Quantitative	202—203
Quinia	204
Quinidia	205
Cinchonia	205
Cinchonidia	205
Quinicine, Cinchonidine	206
Kinic (Quinic) Acid	206
Quinovic Acid	206
Cincho-tannic Acid	206
Cinchona-red	207
Quiniæ Sulphas	209—212

	PAGE.
Cinchonæ Pallidæ Cortex	213
<i>Cinchona officinalis</i> , var. <i>Condaminea</i> .	
Cinchonæ Rubræ Cortex	214
<i>Cinchona succirubra</i> .	
Cinchonæ Lancifoliæ Cortex	215
<i>Cinchona lancifolia</i> .	
Complete Analysis of Cinchona Bark for Alkaloids ..	216—218
Ipecacuanhæ Radix	218—222
<i>Cephaëlis Ipecacuanha</i> .	
Natural Order—Valerianaceæ	222—224
Valerianæ Radix	223
<i>Valeriana officinalis</i> .	
Natural Order—Compositæ	225—240
Sub-Orders—1, Tubulifloræ; 2, Labiatifloræ; 3, Ligulifloræ	
Pyrethri Radix	226
<i>Anacylus Pyrethrum</i> .	
Anthemidis Flores	227—229
<i>Anthemis nobilis</i> .	
Arnicae Radix.....	230
<i>Arnica montana</i> .	
Arnicae Flores	231
Santonica	232—234
<i>Artemisia Cina</i> , <i>Artemisia maritima</i> .	
Inulæ Radix	234—236
<i>Inula Helenium</i> .	
Taraxaci Radix	236—238
<i>Taraxacum officinale</i> , <i>Taraxacum Dens Leonis</i> .	
Lactuca Virosa	238—240

	PAGE.
Natural Order—Lobeliaceæ	240—242
Lobelia	240
<i>Lobelia inflata.</i>	
SUB-DIVISION II.—HYPOSTAMINEÆ	242—245
Natural Order—Ericaceæ	242—245
Uvæ Ursi Folia.....	243—245
<i>Arctostaphylos Uva Ursi.</i>	
SUB-DIVISION III.—EPIPETALÆ	245—309
Natural Order—Sapotaceæ	245—247
Gutta Percha.....	245—247
<i>Isonandra gutta.</i>	
Natural Order—Styraceæ	247—250
Benzoinum.....	247—249
<i>Styrax Benzoin.</i>	
Special Distinctive Tests	249
Acidum Benzoicum	250
Natural Order—Loganiaceæ.....	251—257
Nux Vomica	251
<i>Strychnos Nux vomica.</i>	
Strychnia	252
Brucia	254
Ignatii Semina	256
<i>Strychnos Ignatia.</i>	
Spigeliæ Radix	256
<i>Spigelia Marilandica.</i>	
Natural Order—Gentianaceæ	257—262
Gentianæ Radix	258—260
<i>Gentiana lutea.</i>	

	PAGE.
Chirata	260—262
<i>Ophelia Chirata.</i>	
Natural Order—Asclepiadaceæ	263
Hemidesmi Radix.....	263
<i>Hemidesmus Indicus.</i>	
Natural Order—Convolvulaceæ	264—272
Jalapa	264
<i>Exogonium purga.</i>	
Jalapæ Resina	265
Scammoniæ Radix	268
<i>Convolvulus Scammonia.</i>	
Scammoniæ Resina	268
Scammonium.....	270—272
Natural Order—Solanaceæ.....	272—276
Sub-Orders—1, Rectembryeæ; 2, Curvembryeæ	
Dulcamara	273
<i>Solanum Dulcamara.</i>	
Capsici Fructus	274
<i>Capsicum fastigiatum.</i>	
Natural Order—Atropaceæ	276—288
Belladonnæ Radix.....	276
<i>Atropa Belladonna.</i>	
Atropia	278—280
Atropiæ Sulphas	280
Belladonnæ Folia	281
Stramonii Folia.....	282
<i>Datura Stramonium.</i>	
Stramonii Semina	283

TABLE OF CONTENTS.

xxiii.

	PAGE.
Datura Tatula	284
Hyoscyami Folia	284—287
<i>Hyoscyamus niger.</i>	
Tabaci Folia	287
<i>Nicotiana Tabacum.</i>	
Nicotia	288
Natural Order—Oleaceæ	289—297
Manna	289
<i>Fraxinus Ornus, F. rotundifolia.</i>	
Olivæ Oleum	291—294
<i>Olea Europæa.</i>	
Soaps	294—297
Natural Order—Boraginaceæ	297
Alkanet Root	297
<i>Anchusa tinctoria.</i>	
Symphytum officinale	298
<i>Comfrey.</i>	
Natural Order—Labiatae	298—305
Lavandulæ Oleum	298
<i>Lavandula vera.</i>	
Menthæ Piperitæ Oleum	300
<i>Mentha piperita.</i>	
Menthæ Viridis Oleum	302
<i>Mentha viridis.</i>	
Menthæ Pulegii Oleum	303
<i>Mentha Pulegium.</i>	
Rosmarini Oleum	304
<i>Rosmarinus officinalis.</i>	

	PAGE.
Natural Order—Scrophulariaceæ	306—309
<i>Digitalis Folia</i>	306—309
<i>Digitalis purpurea.</i>	
<i>Digitalinum</i>	307
SUB-CLASS IV.—MONOCHLAMYDEÆ [APETALÆ]	310—378
Natural Order—Polygonaceæ	310—315
<i>Rhei Radix</i>	310—315
<i>Rheum officinale, Rheum palmatum, v. Tanguticum.</i>	
Natural Order—Piperaceæ.....	316—321
<i>Piper nigrum</i>	316
<i>Piper album</i>	317
<i>Piper longum</i>	318
<i>P. officinarum.</i>	
<i>Cubeba</i>	318—319
<i>Cubeba officinalis.</i>	
<i>Cubebæ Oleum</i>	319
<i>Maticæ Folia</i>	320
<i>Piper angustifolium, Artanthe elongata.</i>	
Natural Order—Thymelaceæ	321
<i>Mezerii Cortex</i>	321
<i>Daphne Mezereum, D. Laureola.</i>	
Natural Order—Lauraceæ	323—333
<i>Camphora</i>	323—326
<i>Cinnamomum Camphora, Camphora officinarum.</i>	
<i>Cinnamomi Cortex</i>	326
<i>Cinnamomum Zeylanicum.</i>	
<i>Cinnamomi Oleum</i>	327
<i>Cinnamic Acid</i>	328

TABLE OF CONTENTS.

xxv.

	PAGE.
Cassiæ Cortex	329
<i>Cinnamomum Cassia.</i>	
Lauri Fructus	330
<i>Laurus nobilis.</i>	
Nectrandæ Cortex.....	330
<i>Nectandra Rodiæi.</i>	
Beberiæ Sulphas	331
Sassafras Radix	332
<i>Sassafras officinale.</i>	
Natural Order—Myristicaceæ	334—336
Myristica.....	334
<i>Myristica officinalis.</i>	
Oleum Myristicæ Expressum	335
Oleum Myristicæ	335
Natural Order—Ulmaceæ	336
Ulmi Cortex	336
<i>Ulmus campestris.</i>	
Natural Order—Cannabinaceæ	337—340
Cannabis Indica	337
<i>Cannabis sativa.</i>	
Lupulus	339
<i>Humulus Lupulus.</i>	
Natural Order—Moraceæ	341
Caricæ.....	341
<i>Ficus Carica.</i>	
Mori Succus	341
<i>Morus nigra.</i>	
Natural Order—Euphorbiaceæ	342—350
Cascarillæ Cortex	342
<i>Croton Eluteria.</i>	

	PAGE.
Tiglli Semina.....	344
<i>Croton Tiglium.</i>	
Crotonis Oleum.....	345
Euphorbium	346
<i>Euphorbia resinifera.</i>	
Ricini Semina	346
<i>Ricinus communis.</i>	
Ricini Oleum	347
Kamala	349
<i>Rottlera tinctoria.</i>	
Natural Order—Aristolochiaceæ.....	350—352
Aristolochiæ Radix	350
<i>Aristolochia longa, A. rotunda.</i>	
Serpentariæ Radix	351
<i>Aristolochia Serpentaria.</i>	
Natural Order—Santalaceæ	352
Santali. Lignum.....	352
<i>Santalum album.</i>	
Natural Order—Cupuliferæ, vel. Corylaceæ	353—360
Quercus Cortêx	353
<i>Quercus pedunculata.</i>	
Gallæ	354—356
<i>Diplolepis Gallæ tinctoriæ (Quercus infectoria).</i>	
Tannic Acid	356—358
Gallic Acid.....	359
Natural Order—Altingiaceæ.....	360
Styrax Præparatus	360
<i>Liquidambar orientale.</i>	
Natural Order—Salicaceæ	362

	PAGE.
Salicis Cortex	362
<i>Salix fragilis, S. alba, S. caprea.</i>	
Salicin	363
Salicylic Acid.....	363
DIVISION II.—GYMNOSPERMIA	364—378
Natural Order—Coniferæ	364—378
Laricis Cortex	364
<i>Larix Europæa.</i>	
Terebinthina Veneta	366
<i>Larix Europæa.</i>	
Juniperi Fructus	366
<i>Juniperus communis.</i>	
Juniperi Oleum	367
Juniperi Oleum Empyreumaticum.....	367
<i>Juniperus Oxycedrus.</i>	
Sabinæ Cacumina.....	367
<i>Juniperus Sabina.</i>	
Sabinæ Oleum	368
American Turpentine	369
<i>Pinus palustris, P. tæda, P. pinaster.</i>	
Terebinthinæ Oleum	369
Creasotum	371
Sandarach, <i>vel.</i> Sandarac	373
<i>Callitris quadrivalvis.</i>	
Resina	373
Special Distinctive Tests.....	374
Thus Americanum	375
<i>Pinus tæda, P. palustris.</i>	

	PAGE.
Terebinthina Canadensis.....	375
<i>Abies balsamea.</i>	
Special Distinctive Tests	376
Pix Burgundica.....	376
<i>Abies excelsa.</i>	
Pix Liquida	378
<i>Pinus sylvestris.</i>	
Pix Nigra	378
CLASS II.—MONOCOTYLEDONES	379—417
SUB-CLASS I.—DICTYOGENÆ.....	379—384
Natural Order—Smilacæ	379—384
Sarsæ Radix	379
<i>Smilax officinalis.</i>	
Special Distinctive Tests.....	381
Varieties of Sarsaparilla	382
SUB-CLASS II.—PETALOIDEÆ	384—414
SUB-DIVISION I.—EPIGYNÆ	384—395
Natural Order—Orchidacæ.....	384
Vanilla	385
<i>Vanilla planifolia.</i>	
Salep	385
<i>Orchis mascula.</i>	
Natural Order—Zingiberacæ.....	386—391
Grana Paradisi	386
<i>Amomum Melegueta.</i>	
Curcumæ Rhizoma	386
<i>Curcuma longa.</i>	
Cardamomum	387—389
<i>Elettaria Cardamomum.</i>	

TABLE OF CONTENTS.

xxix.

	PAGE.
Zingiber	389—391
<i>Zingiber officinale.</i>	
Galangal.....	391
<i>Radix Galangæ minoris, Alpinia officinarum.</i>	
Natural Order—Marantaceæ	392
<i>Maranta arundinacea.</i>	
Tous les Mois	392
<i>Canna edulis.</i>	
Natural Order—Iridaceæ	393—395
Crocus.....	393
<i>Crocus sativus.</i>	
SUB-DIVISION II.—HYPOGYNÆ	395—417
Natural Order—Liliaceæ.....	395—404
Aloe Barbadosis	396—399
<i>Aloe vulgaris.</i>	
Aloe Socotrina	399—401
<i>Aloe Socotrina.</i>	
Aloe Capensis	402
<i>Aloe spicata.</i>	
Natal Aloes	402
Scilla	402—404
<i>Urginea Scilla, Scilla maritima.</i>	
Natural Order—Melanthaceæ	405
Sabadilla.....	405
<i>Asagræa officinalis.</i>	
Veratria	405—407
Colchici Cormus	407
Colchici Semina	409

	PAGE.
Veratri Albi Radix	410
<i>Veratrum album.</i>	
Veratri Viridis Radix	411
<i>Veratrum viride.</i>	
Natural Order—Palmaceæ	412—414
Areca	412
<i>Areca Catechu.</i>	
Sanguis Draconis	413
<i>Calamus Draco.</i>	
Special Distinctive Tests.....	413
Sago	414
<i>Sagus</i> (species various).	
Natural Order—Acoraceæ, vel. Aroideæ	414
Calami Aromatici Rhizoma.....	414
<i>Acorus Calamus.</i>	
SUB-CLASS III.—GLUMACEÆ.....	415—417
Natural Order—Graminaceæ	415—417
Hordeum Decorticatum	415
<i>Hordeum distichon.</i>	
Tritici Farina.....	416
<i>Triticum vulgare.</i>	
Amylum	416
Glycerinum Amyli	417
Mucilago Amyli.....	417
Saccharum Purificatum	417
SUB-KINGDOM II.—CRYPTOGAMIA.....	418—430
CLASS III.—ACOTYLEDONES	418—430
SUB-CLASS I.—ACROGENÆ.....	418—420

	PAGE.
Natural Order—Filices.....	418
Filix Mas	419
<i>Aspidium Filix Mas.</i>	
Natural Order—Lycopodiaceæ	420
Lycopodium	420
<i>Lycopodium clavatum.</i>	
SUB-CLASS II.—THALLOGENÆ	421—430
Natural Order—Lichenes	421—424
Cetraria	421
<i>Cetraria islandica.</i>	
<i>Litmus.</i>	422
<i>Roccella</i> (species various), <i>R. tinctoria</i> , <i>R. fuciformis.</i>	
Natural Order—Fungi	425—428
Ergota.....	425—428
<i>Claviceps purpurea.</i>	
Natural Order—Algæ	429
Chondrus Crispus	429
Gracilaria Lichenoides	430

The Animal Kingdom.

SUB-KINGDOM I.—PROTOZOA	433
CLASS—RHIZOPODA	433
Order—Spongida	433
<i>Spongia Officinalis</i>	433
SUB-KINGDOM II.—CÆLEENTERATA	434
SUB-KINGDOM III.—ANNULOIDA.....	434

	PAGE.
SUB-KINGDOM IV.—ANNULOSA	435
DIVISION I.—ANARTHROPODA	435
CLASS—ANNELIDA	435
Order—Hirudinea	436
Hirudo.....	436
<i>Sanguisuga medicinalis</i> , <i>S. officinalis</i> .	
DIVISION II.—ARTHROPODA, OR ARTICULATA ..	437
CLASS—INSECTA	437
Order—Hemiptera	438
Sub-Order—Homoptera	438
Coccus.....	438—440
<i>Coccus cacti</i> .	
Order—Hymenoptera	441—447
Section—Aculeata.....	441
Mel	441
<i>Apis mellifica</i> .	
Sucrose, Glucose, Lævulose	442
Adulterations	444
Cera Flava	445
<i>Apis mellifica</i> .	
Cera Alba	445
Order—Coleoptera	448
Cantharis	448
<i>Cantharis vesicatoria</i> .	
SUB-KINGDOM V.—MOLLUSCA	451
SUB-KINGDOM VI.—VERTEBRATA	452—469
DIVISION I.—ICTHYOPSIDA.....	453—455
CLASS—PISCES	452

	PAGE.
Order—Teleostei	453
Sub-Order—Anacanthini.....	453
Oleum Morrhuae, Oleum Jecoris aselli	453
<i>Gadus Morrhua.</i>	
Ichthyocola	454
<i>Acipenser.</i>	
DIVISION II.—SAUROPSIDA	455—457
CLASS—AVES	455
Order—Carinatae	456
Sub-Order—Rasores.....	456
Albumen Ovi	456
<i>Gallus Bankiva, var. domestica.</i>	
Vitellus Ovi	457
DIVISION III.—MAMMALIA	458—469
Order—Cetacea	458
Cetaceum	458
<i>Physeter macrocephalus.</i>	
Order—Ungulata	459
Sub-Order—Artiodactyla.....	459
Adeps Præparatus	460—461
<i>Sus scrofa.</i>	
Pepsin	462
Lac	463
Analysis of.....	463—464
Saccharum Lactis.....	465
Fel Bovinum Purificatum	465
<i>Bos Taurus.</i>	

	PAGE.
Moschus	466—467
<i>Moschus moschiferus.</i>	
Sevum Præparatum	467
<i>Ovis aries.</i>	
Order—Rodentia	468—469
Castoreum	468
<i>Castor fiber.</i>	

Appendices.

- I.—A Classification and Description of the Chief
Starches existing in the Commoner Articles of Materia
Medica and Food.
- II.—Museum Cabinet. — Drawers for the Recognition of
Materia Medica Specimens arranged for the use of Students.

MATERIA MEDICA
OF THE
VEGETABLE KINGDOM.



SUB-KINGDOM I.—PHANEROGAMIA.

FLOWERING PLANTS PROPAGATED BY SEEDS.

CLASS I.—DICOTYLEDONES.

Plants having 2 cotyledons in the seed ; germination exorhizal ; root true or tap-shaped ; stem exogenous ; leaves usually reticulated in venation ; parts of the flowers arranged in multiples of four or five.

DIVISION I.—ANGIOSPERMIA.

Plants having their ovules enclosed in an ovary.

SUB-CLASS I.—THALAMIFLORÆ.

Having the calyx and corolla usually distinct and the stamens hypogynous.

NATURAL ORDER.—RANUNCULACEÆ.

Usually herbs with a watery acrid juice ; leaves divided and without evident stipules ; calyx 3—6 sepals, deciduous ; corolla 3—15 petals, sometimes suppressed ; æstivation

imbricate; stamens indefinite and hypogynous, with adnate anthers; pistil apocarpous and superior; fruit achenia, follicles, or rarely baccate; seeds with horny albumen.

ACONITI FOLIA.—The fresh leaves and flowering tops of *Aconitum Napellus*, cultivated in Britain.

ACONITI RADIX.—The dried root of the same plant, imported from Germany, or cultivated in Britain.

Botanical Source.—The *Aconitum Napellus* (*Aconite*, *Monkshood*, *Wolfsbane*) is an herbaceous plant, with a conical root and perennial erect stem; leaves numerous, simple, palmatisected, with pinnatifid lobes; dark green above, pale beneath; inflorescence an indefinite raceme, of dark blue flowers, with the calyx petaloid and galeate, constituting the whole of the outwardly visible floral envelope; the petals being distorted and contained within the helmet-shaped calyx; the fruit apocarpous, consisting of three follicles.

Collection.—The leaves and flowering tops are gathered in England when about one-third of the flowers

are expanded; while the root is dug up in winter, or early spring, before the leaves have appeared. The great importance of this matter has been proved by experiment, it having been found that while the average yield of a pound of fresh root collected any time *after flowering* is 8 grains, the quantity obtainable from roots collected after the leaves have expanded, but *before flowering*, is considerably less.

Description.—Aconite leaves have been already described as being simple and palmatisected with pinnatifid lobes, but, more strictly speaking, the segments of the leaf are distinctly trifid and impressed deeply by veins running right to the top of each lobe without many veinlets.

Aconite root, as found in commerce, is from 2 to 4 inches long, conical in shape, and coffee-coloured externally, but whitish internally. It breaks with a short fracture, and is crowned by the remains of the base of the stem, at which point it is knotty, and is usually from $\frac{1}{2}$ to 1 inch in thickness. When a sound root is broken it is seen to have a pure white medullium, which is found to be polygonal when examined with a glass.

Uses.—Aconitia is a very active poison, causing paralysis of the nerves. There is no dose mentioned in B.P., but 1-400th to 1-50th of a grain may be given by the mouth. When subcutaneously injected the dose should not exceed 1-200th of a grain for an adult.

In the early stages of pneumonia, bronchitis, tonsillitis, and croup, the tincture has proved an effectual remedy.

The antidotes are promptly induced vomiting, followed by stimulants and sinapisms to the spine.

Aconite is used as an anodyne and sedative.

The ointment of Aconitia is a useful agent for the relief of sciatica, neuralgia, and lumbago. Care must be taken that aconite be not rubbed in where the skin is broken, or poisonous symptoms may result from its too rapid absorption.

Chemistry.—The active principle of both the root and leaves is an alkaloid called Aconitia (or Aconitina) $C_{30}H_{47}NO_7$, which is combined with an acid known as Aconitic or Equisetic Acid, $H_3C_6H_3O_6$.

The B.P. prescribes the following process for the manufacture of Aconitia in the amorphous form:—

Take of Aconite root in coarse powder, 14 pounds:

rectified spirit, distilled water, diluted sulphuric acid, solution of ammonia, pure ether, of each a sufficiency.

Pour upon the Aconite root three gallons of the spirit, mix them well, and heat until ebullition commences, then cool and macerate for four days. Transfer the whole to a displacement apparatus, and percolate, adding more spirit, when requisite, until the root is exhausted. Distil off the greater part of the spirit from the tincture, and evaporate the remainder over a water-bath until the whole of the alcohol has been dissipated. Mix the residual extract thoroughly with twice its weight of boiling distilled water, and when it has cooled to the temperature of the atmosphere, filter through paper. To the filtered liquid add solution of ammonia in slight excess, and heat them gently over a water-bath. Separate the precipitate on a filter and dry it. Reduce this to coarse powder, and macerate it in successive portions of the pure ether with frequent agitation. Decant the several products, mix, and distil off the ether until the extract is dry. Dissolve the dry extract in warm distilled water acidulated with sulphuric acid; and when the solution is cold, precipitate it by the cautious

addition of solution of ammonia diluted with four times its bulk of distilled water. Wash the precipitate on a filter with a small quantity of cold distilled water, and dry it by slight pressure between folds of filtering paper.

In the spirituous extract of the root we have Aconitate of Aconitia, together with resinous matters, which latter are left behind when the alcoholic extract is treated with water. Ammonia is added which combines with the Aconitic acid, and Aconitia is precipitated. Any ammoniacal salt adherent and other impurities are separated by the ether in which they are insoluble. The Aconitia is subsequently converted into soluble sulphate of Aconitia by means of the diluted sulphuric acid; and, lastly, the sulphate is decomposed by ammonium hydrate, the alkaloid being precipitated, whilst ammonium sulphate is left in solution. By this process we can obtain about 35·5 grains of Aconitia from 1 lb. of dried Aconite root.

Characters and Tests.—A white, usually amorphous solid, soluble in 150 parts of cold, and 50 parts of hot water, and much more soluble in alcohol and in ether; strongly alkaline to reddened litmus, neutral-

ising acids, and precipitated from them by the caustic alkalies, but not by ammonium carbonate, or the sodium or potassium bicarbonates. It melts with heat, and burns with a smoky flame, leaving no residue when burned with free access of air. When rubbed on the skin it causes a tingling sensation, followed by prolonged numbness.

Aconitia communicates a very characteristic violet tint to syrupy phosphoric acid.

Besides the official alkaloid above mentioned, Aconitia has been obtained in a crystalline state. It is then described as having the formula $C_{54}H_{40}NO_2$ and as being nearly insoluble in water, but freely soluble in dilute acids, to form crystalline salts, especially the nitrate. Its best solvent is stated to be chloroform. Alkaloids, called Napelline, Aconella, and others, have been from time to time described, but their existence in the official root is very questionable.

Extractum Aconiti.

The expressed juice of the fresh leaves and flowering tops of Aconite evaporated to an extract.

(100 lbs. of plant yield 50 lbs. of juice or 7 lbs. of extract.)

Dose—1 to 2 grains.

Linimentum Aconiti. (1 in 1)

Aconite Root, in coarse powder	20 ounces
Camphor	1 ounce
Rectified Spirit..... ad.	20 fluid ounces

Moisten the root with spirit for 3 days, then percolate.

Tinctura Aconiti. (1 in 8)

Aconite Root, in coarse powder	2½ ounces
Rectified Spirit..... ad.	20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—5 to 15 minims (1 grain of Aconite in 9 minims.)

Unguentum Aconitiæ.

Aconitia	8 grains
----------------	----------

Dissolve in 30 minims of Rectified Spirit and rub up with
1 ounce of Prepared Lard.

“Fleming’s Tincture,” which is sometimes prescribed, is
nearly four times stronger than the tincture of the B.P.

Adulterations and Substitutions.—Most species of Aconite contain Aconitia. The following is the order of the plants according to their virulency:—*A. ferox*, *A. Napellus*, *A. neomontanum*, *A. Cammarum*, *A. paniculatum*. It has been stated that *A. heterophyllum* (*Atis*), an Indian species, possesses tonic properties, and is not poisonous.

The root being collected in the mountainous parts of Europe by peasants occupied in pasturing sheep and

cattle, it is not surprising that it should often be dug up without due regard to the proper season or even the proper species.

ACONITI RADIX INDICA, Indian Aconite, *Bish* or *Bikh*.—The dried root of *Aconitum ferox*, imported from India (Nepal).

Botanical Source.—The *Aconitum ferox* grows in the lower parts of the Himalayas. The leaf is somewhat broader in the segments than that of *A. Napellus*, and decidedly more symmetrical in shape. The flowers are also larger and of a very dull blue.

Collection.—The roots were formerly collected very much at random, and other species of aconite were often mixed with the *ferox* root, but this is now somewhat improved, and the recent importations are fairly uniform. They are dried by fire-heat.

Description.—Elongated and slightly conical, 3 to 4 inches long by 1 to 1½ inches in diameter at the stem end, which latter has not so well marked remains of the stem, owing to close cutting; brownish externally, but

having patches where the brown colour has been rubbed off and the white interior shows. Generally, the lower ends of the roots have been cut off in digging, and they therefore exhibit a horny whitish fracture at the narrow end.

Uses.—Imported solely for the manufacture of *Aconitia*.

Chemistry.—The active principle is supposed to be *Pseud-aconitine*, an alkaloid existing, both crystalline and amorphous, only slightly soluble in cold ether or rectified spirit, and when boiled in water it fuses and becomes sticky, thus differing in these points from ordinary Aconitine.

PODOPHYLLI RADIX, *Podophyllum* Root.—The dried rhizome and rootlets of *Podophyllum peltatum* imported from North America.

Botanical Source.—*Podophyllum peltatum* (*May apple*, or *Mandrake*) is an herbaceous plant with a creeping horizontal rhizome. The leaves are peltate and sub-divided into 5 or 7 wedged-shaped lobes—in-

florescence solitary, terminal consisting of a white flower having a caducous calyx.*

Description.—The rhizome is reddish brown to greyish brown in colour, somewhat flattened and marked by knotty joints, which each indicate one year's growth. These nodes are surmounted by a depressed scar where the stem has fallen off, and have either rootlets or small whitish marks where they have broken off. As a rule, the portion between the nodes is smooth, but sometimes deeply furrowed in a longitudinal direction.

Uses.—Powerfully cathartic, acting rapidly upon the liver and producing bilious evacuations. It is rarely given *per se*, but nearly always with some other purgative. From the similarity of its action to that of calomel, it has been called the "vegetable mercury."

The powdered rhizome is seldom given in England, but the dose is 10 to 20 grains. The dose of the resin is from $\frac{1}{4}$ to 1 grain. Care should be taken in its administration, as its action is very variable.

NOTE.—*This plant may be looked upon as a connecting link between the Ranunculaceæ and Berberidaceæ, and is by many botanists placed in the latter order.*

Chemistry.—The active principle is a resinous matter, of which the dried root yields from 3 to 5 per cent., extracted by alcohol, and official in the B.P. as Podophylli Resina. It also contains, in considerable quantities, a crystalline alkaloid, called Berberine, an odorous principle, which may be obtained by sublimation in colourless scales, and saponin.

Preparation of Resin of Podophyllum.—Exhaust 1 pound of Podophyllum with 3 pints, or a sufficiency of rectified spirit, by percolation; place the tincture in a still, and draw off the greater part of the spirit. Slowly pour the liquid which remains after the distillation into three times its volume of water, acidulated with 1-24th of its bulk of hydrochloric acid, constantly stirring. Allow the mixture to stand for 24 hours to deposit the resin. Wash the resin on a filter with distilled water and dry it on a stove.

The rectified spirit dissolves the whole of the active resin along with some of the Berberine colouring matter. The acidulated water is to facilitate the separation of the resin, as it settles down very slowly if precipitated by cold water only, and if thrown down by hot water it fuses into a brown cake. The washing with water

removes the greater part of the Berberine. The resin should not be dried at a temperature above 90° F., otherwise it will become very dark in colour.

HELLEBORI RADIX NIGRI, Black Hellebore Root, *Radix Melampodii*.—The dried rhizome and rootlets of *Helleborus niger*, from Southern and Eastern Europe, cultivated in Britain.

Botanical Source.—The *Helleborus niger* (or *Christmas Rose*) is a herb with an almost black rhizome. The leaves are pedate with five ovate lanceolate divisions. Flowers situated on a scape and having a petaloid persistent calyx, with a few large oval bracts. The petals are abortive, forming a circle of short tubular glands, called, by old botanists, nectaries. The plant flowers at Christmas, and bears an apocarpous fruit of several follicles.

Description.—The dried rhizome is very dark in colour, rough, knotty, and irregular in form, and the rootlets are usually more or less broken away. It is in pieces from 1½ to 3 inches long by 3-10ths of an inch thick, and whitish internally. When cut it exhibits a

meditullium, which is distinctly circular and only slightly stellate in very old samples.

It must carefully be distinguished from *Actæa spicata*, which has a much larger rhizome with a red tinge outside, and a cruciform or triangular meditullium; while true Hellebore has a slightly stellate one, and possesses no red tinge. As *Actæa* contains tannin, an aqueous infusion is blackened by ferric salts; which is not the case with an infusion of Black Hellebore.

Uses.—Drastic and emmenagogue (ἐμμήνια the menstrual discharge, and ἀγωγός evoking) used in dropsy, nervous diseases, and chronic skin affections. Dose—10 to 20 grains. Formerly official in the Pharm. Lond.

Chemistry.—Its action depends on a glucoside, Helleborin (0.05 per cent.) $C_{36}H_{42}O_6$. It also contains another glucoside, Helleborein ($C_{26}H_{44}O_{15}$), which is said to be inert. Helleborin is soluble in water and alcohol, and crystallises in shining colourless needles. When boiled with a dilute acid it forms glucose and Helleboresin, $C_{30}H_{38}O_4$; whilst Helleborein yields Helleboretin, $C_{14}H_{20}O_3$, and glucose.

STAPHISAGRIÆ SEMINA, Staves-acre Seeds. The ripe seeds of *Delphinium Staphisagria*, imported from Germany and the South of France.

Botanical Source.—The *Delphinium Staphisagria* is a biennial herb much resembling Aconite in the shape of its leaves, but differing in being slightly pubescent, while Aconite is purely glabrous. The inflorescence is a raceme of blue flowers in which the petaloid and *calcarate calyx* forms a ready means of distinction from the galeate flowers of Aconite. The fruit is a bunch of follicles.

Collection.—The fruits are collected when ripe, just before they dehisce, and the seeds removed and dried.

Description.—About $\frac{1}{4}$ of an inch in length, and in the form of a 4-sided pyramid *with one side much larger than the others and slightly rounded*. The testa is earthy-coloured, wrinkled, and deeply pitted. The seeds are fairly uniform in size, 10 usually weighing about 6 grains.

Uses.—The fatty oil of the seeds, extracted by means

of ether, has been strongly recommended as an ointment, combined with lard (1 to 7) in *prurigo senilis*.

Stavesacre has been sometimes administered as an anthelmintic (ἀντὶ against, and ἔλμυς a worm), but is not a safe remedy for internal use, and is almost entirely used in the form of ointment in cases of itch and to destroy *pediculi*.

Chemistry.—The active principles are two alkaloids, *Delphinia* $C_{24}H_{35}NO_2$ and *Staphisagrine* $C_{16}H_{23}NO_2$. The seeds also contain about 25 per cent of fixed oil.

The alkaloids may be obtained by treating an alcoholic extract with boiling water acidulated with sulphuric acid, and precipitating from this solution by means of ammonium hydrate. They may be separated from each other by treating with ether.

ACTÆÆ RACEMOSÆ RADIX, or **CIMIFUGÆ RADIX**, Cimicifuga Root; *Black Snake Root*, *Black Cohosh*, *Bugbane*.—The dried rhizome and rootlets of *Actæa racemosa*, a native of North America.

Botanical Source.—A perennial herb very much

resembling the *Actæa spicata* of English lanes. The leaves are inclined to be biternate with a deeply serrate margin, and the inflorescence is a long raceme of whitish flowers. The fruit is a dry follicle (*Actæa spicata* bearing a baccate fruit).

Description.—Dark brown short knotty rhizomes about $\frac{1}{2}$ an inch thick and 1 to 3 inches long. They have on one side the remains of stems, and on the other numerous wiry roots, which, in many cases, have their fibrils attached. The roots, when broken, exhibit a very characteristic cruciform or stellate medullium.

Uses.—Stimulant expectorant, but its chief use is in rheumatism, and as an external application in cases of inflammation.

Chemistry.—The active portion is a resinous principle called *Cimicifugin* or *Macrotin*, prepared exactly like *Resina Podophylli*. It is sold in America in brown scales or powder, and the root yields about $3\frac{3}{4}$ per cent.

Preparations.—It is official in the United States' Pharmacopœia, and is given in doses of 20 to 30 grains.

There is also a fluid extract (strength 1 in 1), and a tincture (strength 1 in 8 proof spirit).

HYDRASTIS RADIX, *Yellow Root*.—The dried rhizome and rootlets of *Hydrastis Canadensis* from America.

Description.—This root is official in the United States' Pharmacopœia, and is easily known by its yellow colour, tortuous nature, and fine rootlets.

Uses.—Considered as tonic and anti-periodic, and may be given in a tincture (1 in 4 rectified spirit); dose $\frac{1}{2}$ to 1 drachm.

Chemistry.—Its active principle is obtained in very bitter white crystals, strongly alkaline, and is called *hydrastin*. It is nearly insoluble in water, but freely soluble in alcohol, ether, and chloroform.

NATURAL ORDER.—MAGNOLIACEÆ.

Trees or shrubs; leaves alternate and stipulate; stipules sheathing the leaf bud and deciduous; calyx

3—6 deciduous sepals; corolla with a ternary arrangement and imbricate æstivation; stamens numerous and hypogynous; pistil apocarpous.

WINTERI CORTEX *vel.* CORTEX WINTERANUS; *Winter's Bark*.—The dried bark of *Drimys Winteri*, from the Straits of Magellan and the southern parts of South America generally.

Botanical Source.—The true *Drimys* is an evergreen tree growing from 10 to 40 feet in height.

Description.—A very thick bark in short quills or channelled pieces, often twisted or bent backwards. Ashy grey to brownish externally, and rusty brown internally. The inner side is characterised by sharp striæ or longitudinal ridges with some minute cracks, as if the inner bark had contracted to a great extent in drying. These ridges are seen at the fracture to be the broken ends of white liber tissue. This radiating white liber serves to perfectly distinguish its bark from that of *Canella alba*.

Uses.—Stimulating, tonic and anti-scorbutic. Used in South America for chronic gastric affections and diarrhœa.

Chemistry.—It contains tannic acid or some nearly allied bodies; essential oil and resinous matter. Its decoction, on treatment with potash, becomes dark violet; but is *not* turned intensely bluish brown by tincture of iodine.

Substitutions.—The bark of *Cinnamodendron corticosum*, which has been imported as Winter's Bark and mixed with *Canella alba*. This bark is very like *Canella* but darker in colour, and the inner surface is more fibrous. A decoction is turned intensely purple brown by iodine (distinction from *Drimys*), and is also blackened by ferric chloride (distinction from *Canella*).

ANISI STELLATI FRUCTUS, Star Anise Fruit.—The dried fruit of *Illicium anisatum*. imported from China.

Botanical Source.—The *Illicium* is a small tree about 20 feet high, and a native of China, but cultivated also in Japan.

Description.—The fruits are extremely easy to recognise, consisting, when perfect, of 8 one-seeded carpels arranged round a central attachment like a star. The

carpels are partially dehiscent by the ventral suture showing the brightly polished seed. The odour of aniseed is exceedingly strong.

Uses.—As a substitute for true Aniseed in making liqueurs.

Chemistry.—The carpels contain from 4 to 5 per cent. of a volatile oil corresponding in composition to oil of aniseed, and which solidifies below 36° F. The seeds are destitute of aromatic properties, but contain a fixed oil.

NATURAL ORDER.—MENISPERMACEÆ.

Climbing or trailing shrubs; leaves alternate and exstipulate; flowers diœcious, having a tendency to ternary arrangement in all their parts; fruits with a solitary and peculiarly curved seed.

CALUMBÆ RADIX.—The transversely sliced and dried root of *Fateorhiza palmata*, from the forests of Eastern Africa between the Ibo and Zambesi rivers.

Botanical Source.—The *Fateorhiza palmata* is a perennial climbing plant with palmate leaves and dioecious flowers, which was called *Cocculus palmatus* by old writers. It has been also named *Fateorhiza Calumba* and *F. Miersii*, but the distinctions of these varieties have now been sunk in the one species of *palmata* which includes all.

Collection.—The roots are dug up in March, sliced and then dried in the shade.

Description.—In flat circular or oval slices varying much in diameter, but never more than $\frac{1}{2}$ an inch thick. The cortical portion is thick and hard, while the medullium is thin by contraction during drying. The colour is a dull greenish yellow, and the odour peculiarly musty.

Uses.—Calumba is an aromatic bitter and demulcent tonic, with an entire absence of astringency, and consequently devoid of constipating properties. It is given in cases of indigestion, nausea, and flatulence, usually combined with alkalies or iron preparations.

Chemistry.—Calumbin $C_{21}H_{22}O_7$ is the principal

constituent, but it also contains calumbic acid $C_{22}H_{24}O_7$, berberia ($C_{20}H_{17}NO_4$), resinous extract, a trace of volatile oil, and a large quantity of starch.

Berberia must not be confounded with beberia the alkaloid from the Beeberu.

Calumbin is a neutral body having a yellow colour, slightly soluble in water, alcohol, and ether, but freely soluble in alkalies. Calumba does not contain any tannic or gallic acids.

Extractum Calumbæ.

(1 from 8 made with cold water.)

Dose—2 to 10 grains.

Infusum Calumbæ.

Calumba Root $\frac{1}{2}$ ounce

Cold distilled water..... 10 fluid ounces

Macerate 1 hour, and strain.

Incompatibles—Lead, mercury, or Aq. Calcis.

Dose—1 to 2 fluid ounces.

Cold water is used to avoid dissolving out the starch of the root.

Mistura Ferri Aromatica.—Ph. Dublin.

$\frac{1}{2}$ ounce Calumba to 16 fluid ounces—Peppermint water, with iron and other ingredients.

Tinctura Calumbæ.

Calumba Root, cut small	2½ ounces
Proof spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—¼ to 2 drachms (7 grains of Calumba in 1 fluid drachm).

PAREIRÆ RADIX, Pareira Root.—

The dried root of *Chondodendron tomentosum*, imported from Brazil.

Botanical Source.—The true *Chondodendron* is a tall climbing shrub with leaves 12 inches long, and unisexual flowers. It was formerly supposed that the source of the Pareira Root was the *Cissampelos Pareira*, but Hanbury has conclusively proved that this latter plant, although sometimes used medicinally in the West Indies, is never exported.

Description.—According to Hanbury, the true drug is a long branching woody root, attaining 2 inches or more in diameter, but usually met with much smaller, and dividing into rootlets no thicker than a quill or even than a horse-hair. It is remarkably tortuous and marked with transverse ridges as well as with constrictions.

tions and cracks more or less conspicuous; besides which the surface is strongly wrinkled longitudinally. Roots of about 1 inch in diameter, cut transversely, exhibit a central column of .2 to .4 inch in diameter composed of 10 to 20 converging wedges of large-pored woody tissue with 3 or 4 zones divided from each other by a wavy light-coloured line. Crossing these zones are wedge-shaped rays irregularly and sparsely distributed. It is inodorous, but bitter to the taste, and its decoction is turned inky by tincture of iodine.

Uses.—Possesses a tonic influence on the passages of the genito-urinary system, and is found useful in chronic gonorrhœa and cystitis.

Chemistry.—It contains a bitter principle called *Pelosin*, of which it yields about $4\frac{1}{2}$ per cent. *Pelosin* has the formula of $C_{18}H_{21}NO_3$, is insoluble in water, but soluble in alcohol and ether, and forms bitter salts with acids.

Decoctum Pareiræ.

Sliced Pareira Root.....	1½ ounce
Water, <i>distilled</i>	1 pint

Boil for 15 minutes and strain.

DOSE—1 to 2 ounces.

Brodie's decoction was $\frac{1}{2}$ an ounce to 3 pints of distilled water boiled down to 1 pint.

Dose—8 to 12 ounces daily.

Extractum Pareiræ.

Pareira Root, in coarse powder	16 ounces
Boiling Distilled Water	1 gallon

(100 lbs. of root yield about 6 lbs. of extract.)

Dose—10 to 20 grains.

Extractum Pareiræ Liquidum.

Pareira Root, in coarse powder	16 ounces
Boiling Distilled Water	1 gallon
Rectified Spirit	3 fluid ounces

Product, 16 fluid ounces.

Dose— $\frac{1}{2}$ to 2 drachms. 1 fluid drachm is equal to 1 drachm of root or 4 grains of solid extract.

Substitutions.—*Common false Pareira Brava.* This is what has been for many years the ordinary Pareira of the shops, and until lately regarded as *Cissampelos Pareira*. Its origin is uncertain, but it belongs to the Menispermaceæ. It consists of pieces varying from 6 to 12 inches or more in length, and 1 to 4 inches thick. It is heavy and tortuous, and is derived from both stem and root. In the pieces of the STEM the pith is well defined and about $\frac{1}{4}$ of an inch in diameter, and surrounded by many concentric, or sometimes eccentric

rings. In the root *the pith is absent*, and the rings are very eccentric. It is tough and fibrous, and its decoction is not tinged blue by iodine.

Chondodendron tomentosum stems. Pieces of a large and knotty stem from 1 to 4 inches thick. The thicker pieces exhibit a few *concentric* woody zones, and are frequently hollowed by age. The thinner pieces are dotted over with small dark warts.

True Cissampelos Pareira. This is chiefly found in stems varying in size, but never reaching 1 inch in diameter. The fracture is coarse and fibrous, and there are *no concentric rings*.

Besides these the stems and roots of *Abuta rufescens* have been imported as *white pareira*, and those of the *Abuta amara* as *yellow pareira*.

COCCULUS INDICUS, Levant Nut.—

The dried fruits of *Anamirta Cocculus*, imported from Malabar *via* Bombay and Madras.

Botanical Source.—The *Anamirta* is a climbing shrub with unisexual flowers producing 1 to 3 purple *drupes*, which when dried have the appearance of small berries.

Description.—Blackish and slightly reniform hard fruits about 4-10ths of an inch long. They are wrinkled on the surface, and when shaken close to the ear are found not to have the seed loose in the pericarp, thus distinguishing them from Bay berries.

Uses.—Said to be used chiefly by tanners, but it is currently supposed to be secretly employed by unprincipled persons in brewing, and in poisoning fish. More direct information is wanting. It is not used in medicine and is actively poisonous, producing giddiness and tetanic convulsions.

Chemistry.—It owes its action to about 2-5ths to 1 per cent. of Picrotoxin $C_5H_6O_2$, which resides in the seed. It is a colourless neutral substance, probably a glucoside, and is soluble in 150 parts of cold water, 25 parts of boiling water; in 2 parts of ether, and 3 parts of alcohol.

Picrotoxin may be obtained by separating the oil (50 per cent.) from the kernels by expression, exhausting them with rectified spirit, distilling off the spirit, shaking the residue with boiling water acidulated with hydrochloric acid, evaporating and crystallising.

The aqueous solution of *Picrotoxin* may be distin-

guished from the bitter poisonous alkaloids by not giving a precipitate with any metallic salt or any of the usual re-agents for alkaloids. It, however, somewhat colours potassium dichromate and sulphuric acid, and must, therefore, not be mistaken for strychnia.

Two crystallisable, tasteless, non-poisonous substances, having the same composition, and termed respectively Menispermin and Paramenispermin, are said to have been obtained from the pericarp, but they require re-examination.

GULANCHA.—The dried stems and adventitious roots of *Tinospora cordifolia*, from tropical regions in India.

The drug is in the Pharmacopœia of India, and is considered as tonic, anti-periodic, and diuretic. No proper chemical research has ever been made on it.

NATURAL ORDER.—BERBERIDACEÆ.

Frequently shrubs with alternate compound, and usually spiny leaves and regular symmetrical flowers; calyx 3, 4, or 6 sepals in two whorls, deciduous; petals of corolla

equal to sepals and opposite to them ; stamens equal to petals and opposite to them, hypogynous ; anthers two-celled, dehiscing valvularly ; pistil simple.

There are no plants of this order official in Britain, but the bark of the *Berberis vulgaris*, or common barberry, has been used to adulterate Cortex Granati.

In the Indian Pharmacopœia we find the root bark of *B. Asiatica*, which is tonic and anti-periodic. An extract is also made from it and sold as *Rusot*, for affections of the eye.

In America, the herbalists use the rhizome of *Caulophyllum thalictroides*, under the name of Blue Cohosh, for uterine diseases, and that of *Jeffersonia diphylla* (a root resembling Senega) for rheumatism.

NATURAL ORDER.—PAPAVERACEÆ.

Usually herbs with a milky juice ; leaves alternate and exstipulate ; sepals 2, caducous ; petals 4, crumpled in æstivation ; stamens indefinite, hypogynous ; anthers innate ; pistil syncarpous, with 2 or more parietal placentæ ; seeds numerous and albuminous.

RHŒADOS PETALA.—The fresh petals of the *Papaver Rhæas* from indigenous plants.

Botanical Source.—The *Papaver Rhæas* (red or corn poppy) is a small hairy herb with pinnatisected leaves. The calyx is *caducous*, i.e., falls off when the bud expands, and the corolla is bright scarlet in colour.

Collection.—The petals should be collected in June and July, and used while fresh for making *Syrupus Rhæados*.

Uses.—Only as a colouring matter in the form of *Syrupus Rhæados*. No Morphia appears to be present.

Chemistry.—The petals contain 12 per cent. yellow fatty matter, 20 of gum, 28 of lignin and 40 of red colouring matter. This latter is soluble in water and alcohol, but insoluble in ether. Its watery solution is turned dirty violet by ferric salts, and blackish by caustic potash.

A slightly alkaline substance called Rhæadin, $C_{21}H_{21}NO_6$, has been isolated from this plant. It is crystalline and colourless and only soluble in acids. By exposing its solution in acid to the air it becomes bright scarlet. This colour is bleached by alkalis, but restored by acids.

Syrupus Rhœados.

Red Poppy Petals	13 ounces
Refined Sugar	36 ounces
Distilled Water.....q.s. or	20 fluid ounces
Rectified Spirit.....	2½ fluid ounces

Product, 58 ounces by weight, or 43½ fluid ounces by measure; S.G. 1·330.

PAPAVERIS CAPSULÆ, Poppy Heads.—The dried nearly ripe capsules of *Papaver somniferum*, cultivated in Britain.

Botanical Sources.—(1.) The *Papaver somniferum* (var. *album*) is an annual herb from 2 to 4 feet high. Leaves large, sessile, amplexicaul, deeply sinuate, dentate in the margin. Flowers white with a dark spot near the bottom. Capsules *ovate in shape* with a stellate sessile stigma and *indehiscent*. The seeds are light coloured.

(2.) The *Papaver somniferum* (var. *nigrum*) is altogether smaller and more glaucous. The petals are red or violet. The capsule is *globose* in shape and *dehiscent* by pores under the sessile stigma. The seeds are blackish.

(3.) The *P. somniferum* (var. *setigerum*) is the true wild opium poppy, having somewhat acutely lobed

leaves, the teeth sharp and terminating in a bristle ; indeed the whole plant is more or less covered with scattered sharp stiff hairs. This variety is never seen in England.

Collection.—The capsules should be collected when still slightly unripe, because it is found that any little trace of morphia, which may exist in English grown poppies, entirely disappears on their attaining ripeness.

Uses.—Slightly hypnotic, but very variable in strength and consequently not reliable.

Chemistry.—Similar to Opium, but in a very much lower degree of completeness and richness.

Decoctum Papaveris.

Poppy Capsules, bruised	2 ounces
Distilled Water	1½ pint

Boil 10 minutes ; product 20 fluid ounces.

Extractum Papaveris.

Poppy Capsules, freed from the seeds....	16 ounces
Rectified Spirit.....	2 fluid ounces
Boiling Distilled Water	q.s.

Dose—2 to 5 grains.

Syrupus Papaveris.

Poppy Capsules, bruised and freed from	
seeds	36 ounces
Boiling Distilled Water	2½ gallons
Rectified Spirit	16 fluid ounces
Refined Sugar	64 ounces

Product, 104 ounces by weight, or 78½ fluid ounces by measure; S.G. 1.320.

Dose—1 drachm.

PAPAVERIS SEMINA, Maw Seed.

—The seeds of the Poppy, from which a fixed oil is expressed which has drying properties like linseed oil.

OPIUM.—The inspissated juice (obtained by incisions and subsequent spontaneous evaporation) from the unripe fruit of *Papaver somniferum*, grown in Asia Minor.

Botanical Source.—Has been already described.
(See *Papaveris Capsulæ*.)

Collection.—The collection of opium chiefly takes place in Asia Minor, Persia, India, and Egypt. The incisions are made transversely with a knife into the skin of the capsule a few days after the fall of the petals, when the fruit is about 1½ inch in diameter. Great care

is taken that the incision does not penetrate the skin entirely, as then the juice would flow into the interior of the fruit and be lost. The juice oozes out from the cut and is scraped off. When enough has been thus obtained it is made into a lump or cake, covered with a leaf, and put to dry in the shade.

Uses.—Opium arrests the secretions generally except perspiration, which it assists. In large doses it produces irresistible tendency to sleep and insensibility to pain. In many persons the recovery from a full dose of opium is attended with nausea, headache, giddiness, and constipation. It is employed in inflammatory diseases; in fevers, when it is desired to allay delirium; in diarrhœa, and generally as an anodyne to diminish and relieve the spasms of pain in acute organic disease. Its applications are extremely various, and it is one of the most valuable medicines we possess in the hands of a judicious practitioner.

Antidotes.—Emetics and the stomach pump, coupled with continual rousing of the patient, and affusion of cold water on the head and chest.

The dose of opium in powder is $\frac{1}{2}$ to 2 grains, but 1 grain is always looked upon as the medium adult dose.

Description of Commercial Varieties.

Smyrna Opium.—The highest quality in the market. It is in masses of from $\frac{1}{2}$ to 2 lbs. weight and covered externally with poppy leaves, and the red fruits of a species of *Rumex*. It is pale brownish internally, and when examined under the microscope it is found to be a mass of agglutinated tears of a yellowish colour. It yields on an average about 12 per cent. of *Morphia hydrochloras*, when treated by the B.P. process. Good opium blackens externally, hardens, and loses weight by exposure.

Constantinople Opium.—Practically the same as the Smyrna, only imported *via* Constantinople. It is produced in the north-western districts of Asia Minor, and sent to Constantinople by way of Izmid. It is similar to the Smyrna, although usually more symmetrical in the cakes, and not quite so regular in quality. It is equally good for official use with the Smyrna, and yields at least 8 per cent. of morphia and 4 of narcotine. It would also give about 12 per cent. of the hydrochlorate of morphia by the B.P. process. The cakes are usually wrapped in a poppy leaf, with a very evident midrib running up the middle of the cake.

Egyptian Opium.—An inferior variety, now not often met with in commerce. It is in flattish hard cakes, covered with a leaf having a somewhat *palmate* structure and no *Rumex* fruits. It has a musty smell and does not blacken by keeping, but becomes soft. It rarely yields over 6 per cent. of morphia, but usually the amount is more nearly 3 to 4 per cent.

Persian Opium.—In balls or sticks, usually wrapped in paper. It formerly went almost entirely to China, but lately a good deal has been brought to this country for morphia manufacture. It is very variable, as fine samples have been met with yielding 8 to 10 per cent. of morphia, working on the *undried* article, while other pale samples have only yielded $\frac{1}{2}$ to 3 per cent.

East Indian Opium.—Is produced in large quantities in the Government factories in India, and is entirely exported to China, none ever coming to us. The finest variety is known as the *Patna garden opium*, and yields 7 to 8 per cent. of morphia. As exported to China it is called *cannon ball* opium, from being formed into balls and covered with poppy petals, caused to adhere by a liquid called *lévâ*, which is made from the washing of pots and vessels and a little opium, and then evaporated to a sticky fluid.

B.P. test for the strength of Opium.—Break down 100 grains and steep in one ounce of water for 24 hours, stirring the mixture frequently. Transfer to a displacement apparatus and exhaust by percolation with three ounces more of water. To the fluid thus obtained, placed in a flask, add 100 grains of slaked lime to precipitate the meconic acid and liberate the morphia; boil for ten minutes, transfer it to a filter and wash the undissolved matter with 1 ounce of boiling water. Acidulate the filtered fluid slightly with diluted hydrochloric acid, evaporate to half an ounce, and let it cool. Neutralise cautiously with solution of ammonia, carefully avoiding an excess; remove by filtration the brown matter which separates and wash it with 1 ounce of hot water. Mix the washings with the filtrate, concentrate the whole to half an ounce, and add ammonia in slight excess. After 24 hours, collect the precipitate on a weighed filter, wash it with cold water, and dry at 212° F. It ought to weigh at least from 6 to 8 grains.

In addition to this official process several others have been proposed, but none without a drawback of some kind. The following is good, if carefully performed:—Take 100 grains of opium, previously dried at 212° F., rub it intimately with three times its weight of coarsely

powdered pumice-stone, and percolate with boiling ether to remove narcotine, colouring matter, and wax. Just moisten with a little spirit and then extract with water slightly acidulated with acetic acid. The amount of water employed should not, if possible, exceed 5 ounces. Ammonia is now added in the slightest possible excess, and the whole set aside for 48 hours, when the morphia will be deposited in crystals on the sides of the glass. The mother liquor is to be poured away, the morphia purified by re-crystallisation from boiling alcohol, and the crystals dried and weighed.

In addition to the assay for morphia, the amount of water should always be determined in a sample of opium by taking a weighed portion of thin slices and drying over the water-bath until it ceases to lose weight. Besides these points, the dried opium should be exhausted with cold water, and the residue collected on a weighed filter and again dried at 212° F. The amount of this residue should not exceed 45 per cent. The addition of gum arabic may be detected by precipitating a solution of opium with normal plumbic acetate, concentrating the filtrate to a low bulk and precipitating out the gum arabic with alcohol. Genuine opium should contain no starch or tannic acid. Lastly, a portion of opium dried

at 212° F. should be weighed and freely incinerated, when the amount of ash left should not exceed 8 per cent.

Chemistry.—Opium contains the following principal primary alkaloids:—

Morphia (Morphine)— $C_{17}H_{19}NO_3 \cdot H_2O$ (6 to 12 per cent.)

Codeia (Codeine)— $C_{18}H_{21}NO_3$

Thebaia (Thebaine)— $C_{19}H_{21}NO_3$

Narceia (Narceine)— $C_{23}H_{29}NO_9$

Papaverine— $C_{21}H_{21}NO_4$

Narcotia (Narcotine) (Anarcotine)— $C_{22}H_{23}NO_7$
(6 to 8 per cent.)

} under
1 per cent.

It also contains:—

Meconic Acid— $H_3C_7HO_7 \cdot 3H_2O$ (about 8 per cent.)

Meconine— $C_{10}H_{10}O_4$

Rhœadine— $C_{21}H_{21}NO_6$

} Traces.

Resinous Matters

Caoutchouc

Essential Oil

Mucilage

} Variable.

Besides these there are also numberless secondary constituents, *i.e.*, substances produced from opium by

the action of re-agents often during the processes applied for separating the natural constituents. The following are the principal :—

Apomorphia (Apomorphine)— $C_{17}H_{17}NO_2$

Apocodeia (Apocodeine)— $C_{18}H_{19}NO_2$

Cryptopia— $C_{21}H_{23}NO_5$

MORPHIA (Morphine), $C_{17}H_{19}NO_3H_2O$, may be prepared according to the B.P., by macerating one pound of sliced opium for 24 hours with two pints of water and decanting. Repeat this process with the residue a second and third time and finally subject the residue to strong pressure. Evaporate the united liquors in a water-bath to a pint and strain through calico. Add three-quarters of an ounce of calcium chloride, dissolve in four fluid ounces of water and evaporate until the solution solidifies on cooling. Wrap the mass in a double fold of strong calico, and subject it to a powerful pressure, preserving the dark fluid which exudes, as it frequently still contains some morphia. Triturate the squeezed cake with about half a pint of boiling water, filter and wash the residue with boiling distilled water. Evaporate the filtrate until it again solidifies on cooling and subject it to pressure as before, and if the

mass be still coloured dissolve it in water, again evaporate to solidification, and subject it to pressure as before, always preserving the liquid which exudes. Dissolve the pressed cake in six fluid ounces of boiling water, add a quarter of an ounce of purified animal charcoal and digest for 20 minutes ; filter and wash the filtering charcoal with boiling water and add to the solution ammonium hydrate in slight excess. Collect the morphia which separates as the liquid cools on a filter, and wash with cold water until the washing ceases to give a precipitate with a solution of Argentic nitrate, acidulated with nitric acid.

In this process the water extracts from the opium, the morphia and codeia as meconates together with extractive matter, the narcotine remaining undissolved. On adding calcium chloride and evaporating, a cake is obtained containing calcium meconate and the hydro-chlorates of morphia and codeia, while the extractive matters are removed in the liquid which exudes by pressure. On treating the cake with boiling water the calcium meconate remains insoluble and the hydro-chlorates of morphia and codeia dissolve. This process is repeated until they are obtained nearly pure. The solution containing the hydro-chlorates of the alkaloids,

after further purification by animal charcoal, is treated with ammonium hydrate in slight excess, which precipitates the morphia and re-dissolves the codeia.

Morphia crystallises in hexagonal prisms. It has a bitter taste and fuses at 340° . At 350° the alkaloid sublimes and at a higher temperature it chars and burns away. It is very slightly soluble in cold water (1 in 1000) and more so in boiling water (1 in 400), its solutions having an alkaline re-action. It is soluble in 30 parts of boiling rectified spirit and insoluble in chloroform and ether. It is precipitated by potassium and sodium hydrates, instantly soluble in excess. Its best precipitant is sodium bicarbonate. It is also precipitated from its solution by tannic acid.

Tests.—(1.) A little solid morphia treated with strong nitric acid produces an orange-red coloration which is bleached by stannous chloride.


(2.) A fresh portion of solid morphia, treated with neutral ferric chloride, gives an evanescent blue colour.

(3.) Mixed with six times its weight of sugar, and a drop of strong sulphuric acid added, a purple colour is produced.

(4.) Dissolved in strong sulphuric acid and a drop of water added, followed by a drop of nitric acid, the liquid is, first red, then green, then brown.

(5) Mixed with iodic acid and starch paste a blue colour is produced, due to the liberation of iodine.

MORPHIÆ HYDROCHLORAS, B.P.



Is prepared by dissolving morphia in water by the aid of hydrochloric acid until a perfectly neutral solution is obtained, and crystallising. It is found in silky acicular prisms soluble in cold water (1 in 16) and in boiling water (1 in 1); also freely soluble in spirit. Twenty grains dissolved in half an ounce of warm water and precipitated by a slight excess of ammonium hydrate and the precipitate dried by exposure to the air, should yield 15·18 grains of pure morphia. The dose is a half to one grain.

Liquor Morphiae Hydrochloratis.

Hydrochlorate of Morphia	4 grains
Diluted Hydrochloric Acid	8 minims
Rectified Spirit	2 fluid drachms
Distilled Water	6 fluid drachms

($\frac{1}{2}$ grain of Hydrochlorate of Morphia in 1 fluid drachm.
half the strength of P.L.)

DOSE—10 to 60 minims.

Suppositoria Morphię.

Hydrochlorate of Morphia	6 grains
Benzoated Lard	64 grains
White Wax	20 grains
Oil of Theobroma	90 grains

($\frac{1}{2}$ grain of Hydrochlorate of Morphia in each.) Divide into 15 grains for suppositories, or into 12 equal parts.

Suppositoria Morphię cum Sapone.

Hydrochlorate of Morphia.....	6 grains
Glycerin of Starch	50 grains
Curd Soap (in powder)	100 grains
Starch (in powder)	q.s.

($\frac{1}{2}$ grain of Hydrochlorate of Morphia in each.) Divide the mass into 12 equal parts.

Trochisci Morphię.

Hydrochlorate of Morphia	20 grains
Tincture of Tolu.....	$\frac{1}{2}$ fluid ounce
Refined Sugar, in powder	24 ounces
Gum Acacia, in powder	1 ounce
Mucilage of Gum Acacia	q.s. or 2 fluid ounces
Distilled Water	$\frac{1}{2}$ fluid ounce

Divide into 720 lozenges.

DOSE—I to 6. (Each lozenge contains 1-36th grain of Hydrochlorate of Morphia.

Trochisci Morphię et Ipecacuanhæ.


Hydrochlorate of Morphia	20 grains
Ipecacuanha, in fine powder.....	60 grains
Tincture of Tolu.....	$\frac{1}{2}$ fluid ounce
Refined Sugar	24 ounces

Gum Acacia, in powder	1 ounce
Mucilage of Gum Acaciaq.s. or	2 fluid ounces
Distilled Water	$\frac{1}{2}$ fluid ounce

Divide into 720 lozenges.

DOSE—1 to 6. (Each lozenge contains 1-36th grain of Hydrochlorate of Morphia and 1-12th grain of Ipecacuanha.

MORPHIÆ ACETAS, B.P.



Is prepared by precipitating the morphia from a solution of hydrochlorate of morphia by means of a very slight excess of ammonium hydrate, washing and dissolving in the exact quantity of acetic acid necessary to produce a neutral salt, and evaporating to dryness in a water-bath. Acetate of Morphia is a slightly deliquescent salt, soluble in water and spirit. When first prepared it is a white powder, but by keeping it becomes yellowish, owing to the formation of basic Acetate. A similar loss of acetic acid is also apt to take place through evaporation, and the composition of the commercial salt is therefore by no means constant.

Liquor Morphiæ Acetatis.

Acetate of Morphia	4 grains
Diluted Acetic Acid	8 minims
Rectified Spirit.....	2 fluid drachms
Distilled Water	6 fluid drachms

($\frac{1}{2}$ grain of Acetate of Morphia in 1 fluid drachm, half the strength of P.L.)

DOSE—10 to 60 minims.

Injectio Morphiæ Hypodermica.

Hydrochlorate of Morphia	88 grains	
Solution of Ammonia.....		} Of each a sufficiency
Acetic Acid		
Distilled Water		

(1 grain of Acetate of Morphia in 12 minims.)

Dissolve 88 grains of Hydrochlorate of Morphia in two ounces of water by the aid of a gentle heat, then add solution of ammonia so as to precipitate the Morphia and render the liquid slightly alkaline; allow it to cool; collect the precipitate on a filter, wash it with water, and allow it to drain; then transfer the Morphia to a small porcelain dish containing about an ounce of water: apply a gentle heat, and carefully add acetic acid until the Morphia is dissolved and a very slightly acid solution is formed. Add now sufficient water to make the solution measure exactly two fluid ounces; filter and preserve in a stoppered bottle excluded from the light.

The above process is necessary to ensure a solution of definite strength, owing to the tendency, above mentioned, which Acetate of Morphia has to lose acetic acid. The 88 grains of Hydrochlorate represent 66.79 grains of Morphia, and produce 80 grains of Acetate in the two ounces of solution. Twelve minims of the injection therefore contain one grain of Acetate of Morphia.

The amount to be administered is from one to six minims.

Apomorphia. — $C_{17}H_{17}NO_2$ (that is Morphia $C_{17}H_{19}NO_3 - H_2O$), is prepared by heating Morphia in a sealed tube with hydrochloric acid for three hours, during which process it loses a molecule of water, and is converted into the Hydrochlorate of Apomorphia. Apomorphia is soluble in water and ether but more so in alcohol and chloroform. It gives a red colour with nitric acid, and purple with ferric chloride. It is precipitated from its salts both by potassium and ammonium hydrates, but readily soluble in excess, the potash solution becoming red, and the ammonia greenish, by exposure.

USES.—When subcutaneously injected it produces violent sickness, followed by *hypnosis*, and sometimes by *syncope*, and may, therefore, be useful when sudden vomiting is desired in cases of poison.

Codeia (Codeine), $C_{18}H_{21}NO_3$, may be prepared from the mother liquor after the separation of morphia in the B.P. process, by precipitation with potassium hydrate, and subsequent re-solution, treat-

ment with animal charcoal and crystallisation. It is usually found to the extent of nearly 1 per cent. in opium. It is colourless, bitter, and soluble in water (1 in 50) and in boiling water (1 in 25); it is very soluble in alcohol and chloroform, but requires 40 parts of ether. It gives a dull greenish orange with nitric acid, and is not affected by ferric chloride. It is hypnotic in action, but not so much so as morphia. There is a Syrup of Codeia in the French Pharmacopœia.

Narceia (Narceine), $C_{23}H_{29}NO_9$, may be prepared from the liquid remaining after precipitation with Codeia, by removing the colouring matter with acetate of lead, filtering, passing sulphuretted hydrogen through the filtrate to remove lead, and evaporating to a syrup, when it separates in silky crystals. It is soluble in water (1 in 400), and in boiling water (1 in 100); it is less soluble in alcohol, and insoluble in chloroform and ether. It is soluble in glycerin. With dilute hydrochloric acid it is coloured blue. With strong sulphuric acid it becomes first amber and then deep red.

Meconia (Meconine), $C_{10}H_{10}O_4$, may be separated by shaking up the mother liquor, after the crystallisation

of Narceia, with ether. It is soluble in water, glycerin, and chloroform ; less so in ether and alcohol. Heated gently with sulphuric acid it becomes first emerald green and then purple.

Narcotia (Narcotine, Anarcotine), $C_{22}H_{23}NO_7$, is prepared from the residue left after exhausting the opium with water, by treating with acetic acid, precipitating by ammonium hydrate and purifying by dissolving the precipitate in alcohol, and crystallising. It is in colourless prisms, insoluble in cold water, and slightly soluble in boiling water ; soluble in alcohol and in acids, forming bitter salts. It is precipitated by potassium hydrate, insoluble in excess. By treatment with oxidising agents it yields Opianic Acid and Cotarnia. It is destitute of hypnotic influence, and hence the new synonym proposed.

Thebaia (Thebaine), $C_{19}H_{21}NO_3$, prepared from the alcoholic mother liquor from the crystals of narcotine, by evaporating to dryness, treating with acetic acid, filtering and precipitating with acetate of lead, which precipitates Papaverine, and leaves Thebaia in solution. The excess of lead is removed by sulphuric acid, and the alkaloid precipitated by ammonium hydrate. In

colourless, rectangular prisms, soluble in alcohol, ether, and chloroform; with strong cold sulphuric acid it forms a blood red solution.

Papaverine, $C_{21}H_{21}NO_4$, prepared from the lead precipitate above-mentioned by boiling with alcohol, evaporating the solution to dryness, and treating with dilute hydrochloric acid, filtering, and allowing the Hydrochlorate of Papaverine to crystallise. Very slightly soluble in water, soluble in boiling alcohol. Cold sulphuric acid produces no colour, but on heating the solution becomes first purple and afterwards red.

Meconic Acid, $H_3C_7HO_7.3H_2O$, may be prepared from calcium meconate formed during the manufacture of morphia, by treating it with hydrochloric acid and crystallising. It is in colourless micaceous crystals; sparingly soluble in cold water, freely in hot. It gives precipitates with salts of calcium, barium, and lead, and a blood red with ferric chloride, not discharged by mercuric chloride.

Other Opium Preparations :—

Confectio Opii.

Compound Powder of Opium 192 grains

Syrup..... 1 fluid ounce.

Dose—5 to 20 grains (1 grain of opium in 40 grains, nearly).

Emplastrum Opii.

Opium, in very fine powder	1 ounce.
Resin Plaster	9 ounces.

1 part of opium in 10 parts.

Enema Opii.

Tincture of Opium	30 minims
Mucilage of Starch.....	2 fluid ounces

Mix for one enema.

Extractum Opii.

Opium, in thin slices	16 ounces
Distilled Water	6 pints

(100 lbs. of good Opium yield 50 lbs. of extract.)

Dose— $\frac{1}{2}$ to 2 grains.

Extractum Opii Liquidum.

Extract of Opium	1 ounce
Distilled Water	16 fluid ounces
Rectified Spirit	4 fluid ounces

Product 20 fluid ounces. 1 grain of Extract of Opium in 22 minims.

Incompatibles—Most metallic salts: the alkalies, and their carbonates. Dose—10 to 40 minims.

Linimentum Opii (*Anodyne Liniment*).

Tincture of Opium	2 fluid ounces
Liniment of Soap	2 fluid ounces

(1 of Tinct. Opii in 2.)

Pilula Ipecacuanhæ cum Scilla^A.

Comp. Powder of Ipecacuanha	3 ounces
Ammoniacum, in powder	1 ounce
Squill, in powder	1 ounce
Treacle	q.s.

Dose—5 to 10 grains. (5 grains of Dover's powder in about 10 grains.)

Pilula Plumbi cum Opio.

Acetate of Lead, in fine powder	36 grains
Powdered Opium.....	6 grains
Confection of Roses	6 grains

Dose—3 to 5 grains. ($\frac{1}{2}$ grain of opium and 3 grains of lead in 4 grains.)

Pilula Saponis Composita (Opium Pills).

Opium, in powder	$\frac{1}{2}$ ounce
Hard Soap, in powder	2 ounces
Distilled Water	q.s.

Dose—3 to 5 grains. (1 grain of Opium in 5 grains.)

Pulvis Cretæ Aromaticus c Opio.

Aromatic Powder of Chalk	9 $\frac{1}{2}$ ounces
Opium, in powder	$\frac{1}{4}$ ounce

Dose—10 to 40 grains. (1 grain of Opium in 40 grains.)

Pulvis Ipecacuanhæ Compositus (Dover's Powder).

Ipecacuanha Root, in powder	$\frac{1}{2}$ ounce
Opium, in powder	$\frac{1}{2}$ ounce
Sulphate of Potash.....	4 ounces

Dose—5 to 15 grains. (1 grain of Opium in 10 grains.)

Pulvis Kino Compositus.

Kino, in powder	3 $\frac{3}{4}$ ounces
Opium, in powder	$\frac{1}{4}$ ounce
Cinnamon Bark, in powder	1 ounce
Dose—5 to 20 grains. (1 grain of opium in 20 grains.)	

Pulvis Opii Compositus.

Opium, in fine powder	1 $\frac{1}{2}$ ounce
Black Pepper do.	2 ounces
Ginger do.	5 ounces
Caraway Fruit do.	6 ounces
Tragacanth do.	$\frac{1}{2}$ ounce
Dose—2 to 5 grains. (1 grain of Opium in 10 grains, 24 grains to 1 fluid drachm of syrup, form Conf. Opii P.B.)	

Suppositoria Plumbi Composita.

Acetate of Lead	36 grains
Opium, in powder	12 grains
Benzoated Lard	42 grains
White Wax	10 grains
Oil of Theobroma	80 grains

Divide into 12 equal parts (1 grain of Opium in each).

Tinctura Camphoræ Composita (Paregoric).

Opium, in coarse powder	40 grains
Benzoic Acid	40 grains
Camphor	30 grains
Oil of Anise	30 minims
Proof Spirit	ad 20 fluid ounces

Macerate 7 days, then filter.

Dose—15 minims to 1 drachm. ($\frac{1}{2}$ grain of Opium in 1 drachm.)

Tinctura Opii (Laudanum).

Opium, in coarse powder $1\frac{1}{2}$ ounce or 656 grains
 Proof Spirit ad. 20 fluid ounces

Macerate 7 days, and filter.

Incompatibles—The alkalies, their carbonates, and most metallic salts.

Dose—5 to 40 minims. (1 grain of opium in 14 2-3rd minims.)

Antidotes—The stomach pump, emetics, stimulants, compelled exertion, artificial respiration.

Tinctura Opii Ammoniata (Scotch Paregoric).

Opium, in coarse powder 100 grains
 Saffron, cut small 180 grains
 Benzoic Acid 180 grains
 Oil of Anise 1 fluid drachm
 Strong Solution of Ammonia 4 fluid ounces
 Rectified Spirit 16 fluid ounces

Macerate 7 days, then strain, press, filter, and add spirit to 1 pint.

Dose— $\frac{1}{2}$ to 1 drachm. (1 grain of powdered opium in 96 minims.)

Trochisci Opii.

Extract of Opium 72 grains
 Tincture of Tolu $\frac{1}{2}$ fluid ounce
 Refined Sugar, in powder 16 ounces
 Gum Acacia, in powder 2 ounces
 Extract of Liquorice 6 ounces
 Distilled Water q.s.

Divide into 720 lozenges.

Dose—1 to 6. (Each lozenge contains 1-10th grain of extract opium.)

Unguentum Gallæ c Opio.

Ointment of Galls	1 ounce
Opium, in powder	32 grains

Vinum Opii.

Extract of Opium	1 ounce
Cinnamon Bark, bruised	75 grains
Cloves, bruised.....	75 grains
Sherry	ad 20 fluid ounces

Macerate 7 days, then filter.

It is about $\frac{1}{2}$ stronger than the Edinburgh and Dublin, but 1-5th weaker than the London, and corresponds in strength with Extractum Opii Liquidum. Contains 1 grain of Extract of Opium in 22 minims.

Incompatibles—The alkalies, their carbonates, and most metallic salts.

Dose—10 to 40 minims.

SANGUINARIÆ RADIX, Blood Root. The dried rhizome of *Sanguinaria Canadensis*, a native of Canada.

Botanical Source.—This plant, called frequently the *Red Puccoon*, is a stemless herb, with grey reniform leaves having red veins, and a solitary white flower.

Description.—A large knobby perennial rhizome with a bitter acrid taste, and having internally a bright orange red colour.

Uses.—It is official in the United States' Pharmacopœia, and employed as expectorant, diaphoretic, and emetic in croup and hepatic congestion. In large doses it is cardiac, depressive, and a narcotic poison. The U.S.P. has a tincture (1 in 10) and an alcoholic extract.

Chemistry.—It contains a resin, and an alkaloid called Sanguinaria $C_{19}H_{17}NO_4$.

NATURAL ORDER.—CRUCIFERÆ (*vel*
BRASSICACEÆ.)

Edible and non-poisonous herbs with alternate and exstipulate leaves; inflorescence indefinite; calyx 4 sepals inferior; corolla 4 petals cruciform; stamens 6 tetradynamous and hypogynous; pistil superior; placentæ 2 parietal; fruit, a siliqua or silicula; seeds exalbuminous.

ARMORACIÆ RADIX, Horseradish Root.—The fresh root of *Cochlearia Armoracia*, cultivated in Britain.

Botanical Source.—The horseradish is almost too well known to require description. It has large ovate-oblong radical leaves with a well marked acutely crenate

margin. The flower stalk bears a few small sessile lanceolate and serrate leaves, and the inflorescence is a raceme of small white flowers; the fruits being small ovate siliculæ.

Description.—A long fleshy, *cylindrical* root, up to 1 inch in diameter and giving off certain short shoots at the crown. *It is whitish both externally and internally and it does not turn pink when scraped.* These two last points serve easily to distinguish it from fresh aconite root, which has been sometimes mistaken for it with disastrous consequences.

Uses.—Stimulant, diuretic and diaphoretic. Externally rubefacient, but more feeble than mustard.

Chemistry.—Contains *Sinigrin* and *Myrosin*, by the mutual action of which in the presence of water it develops and yields about 2 per cent. of volatile oil of mustard. (See Black Mustard.)

Spiritus Armoraciæ Compositus.

Horseradish, scraped	20 ounces
Bitter Orange Peel, cut small and bruised	20 ounces
Nutmeg, bruised	½ ounce
Proof Spirit	1 gallon
Water	2 pints

Distil 1 gallon.

DOSE—1 to 2 drachms (8 grains of horseradish in 1 fluid drachm).

If this preparation be desired to be really active the scraped radish should be allowed to be steeped in the water for 24 hours before adding the spirit, which latter prevents the formation of essential oil by coagulating the Myrosin.

SINAPIS ALBÆ SEMINA, White Mustard Seed.—The seeds of *Sinapis alba* from indigenous plants.

SINAPIS NIGRÆ SEMINA, Black Mustard Seed.—The seeds of *Sinapis nigra*, from indigenous plants.

Botanical Source.—Both mustards are annual herbs, with the lower leaves rough and lyrate in the black, and pinnatisected in the white. The upper leaves of the black are small and linear lanceolate. Both have racemes of yellow cruciform flowers, those of the white being larger. The fruits of both are siliquæ, but those of the *black* are small, erect, and *smooth*, while the *white* has spreading fruits shorter and broader and all *markedly hispid*.


Description.—The commercial article is the powder

obtained by grinding the *mixed* seeds. It should contain no flour or starch, turmeric or capsicum; all of which may be detected by the microscope.

Uses.—A local stimulant and rubefacient. Especially useful as a stimulating emetic in cases of narcotic poisoning.

Chemistry.—Both seeds contain 23 per cent. of a fixed oil which is separated by expression, and they also each have about 20 per cent. of mucilage. There is also an albuminous ferment called *Myrosin* which exists to a small extent in the black, and up to 15 per cent. in the white, and is soluble in water being coagulated and rendered inactive by heating the solution above 140° F. or by the addition of alcohol. The real active agent, however, is the *essential oil of mustard*, which produces the acidity, but which does not exist ready formed in either seed, being obtained by a fermentation when *black* mustard seeds are steeped in water after removal of the fixed oil. The essential oil so formed is *Allyl sulphocyanate* C_3H_5CNS , and may be isolated by distillation with the water. The formation of this oil is due to the action of the *Myrosin* upon a substance known as *Sinigrin* or *Potassium Myronate*, which exists only in

the black mustard. This body has the formula of $C_{10}H_{18}KNS_2O_{10}$, and may be regarded as consisting of allyl sulphocyanate, potassium hydrosulphate and glucose, into which three substances it splits up under the influence of the *Myrosin*, thus (taking the most simple view):—



As the amount of *Myrosin* present in the black seeds is sometimes not sufficient to act thoroughly on all the *Sinigrin*, it is advisable to mix in a little white mustard when making the oil. The oil produced is equal to about '5 per cent. of the weight of the mustard employed and is very pale yellow; sp. gr. 1·015. It is a powerful vesicant. *Sinigrin* may be isolated from black mustard (after the removal of the fixed oil) by boiling out with alcohol. It crystallises in needles with a cooling bitterish taste. The acidity of white mustard is due to the action of the *Myrosin* on a substance called *sinalbin* $C_{30}H_{44}N_2S_2O_{16}$, which under the quasi fermentation breaks up into acrinyl sulphocyanate $C_7H_7(CNS)O$ together with sulphate of sinapin $C_{16}H_{25}NO_5SO_4$ and glucose $C_6H_{12}O_6$. Of these the acrid principle is the

acrinyl sulphocyanate. It is a volatile oil, but is not produced in sufficient quantity to be worth isolating, and therefore, the true *Oleum Sinapis* B.P. is the allyl sulphocyanate from the black mustard, as described above.

Cataplasma Sinapis.

Mustard, in powder.....	2½ ounces
Linseed Meal	2½ ounces
Boiling Water	10 fluid ounces

Charta Sinapis.

Solution of Gutta-percha	q.s. or 2 fluid ounces
Black Mustard Seeds, in powder	1 ounce

Mix the mustard with the gutta-percha solution so as to form a semi-fluid mixture; pour it on a dinner-plate, and pass one side of strips of cartridge paper over its surface, then lay the paper on a table with the coated side upwards until dry. Before being applied to the skin, let the mustard paper be immersed for a few seconds in tepid water.

Linimentum Sinapis Compositum.

Oil of Mustard	1 fluid drachm
Ethereal Extract of Mezereon	40 grains
Camphor	120 grains
Castor Oil	5 fluid drachms
Rectified Spirit.....	4 fluid ounces

Dissolve the extract and camphor in the spirit, and add the oils. (12 minims of oil of mustard in 1 fluid ounce.)

NATURAL ORDER. CANELLACEÆ.

CANELLÆ ALBÆ CORTEX, *Canella* Bark. The bark of *Canella alba*, imported from the West Indies.

Botanical Source.—The *Canella alba* (or *Wild Cinnamon*) is a tree growing freely in most of the islands of the West Indies, but specially utilised in the Bahama Islands.

Collection.—The tree is first gently beaten so as to remove the greater part of the epiphloeum, and then the remaining portions of the bark are peeled off, and dried in the shade.

Description.—In crooked and irregular quills or channelled pieces, yellowish-white in colour. The outer surface has here and there patches of the epiphloeum dotted with minute lichens, while the inner surface is marked with longitudinal fissures caused by the beating. It has an aromatic cinnamon-like odour, and a peppery taste.

Uses.—Aromatic bitter tonic and stomachic. Dose,

10 to 30 grains, in powder. Still much sought after by old-fashioned country people as a powder mixed with aloes, known as *Hiera Picra*.

It is an ingredient in *Vinum Rhei*, 60 grains to 1 pint.

Chemistry.—The essential constituent is a volatile oil about 94 per cent. *Canella* also contains 8 per cent. of mannite, gum resin, starch, and a bitterish extract.

The bark is quite free from tannic and gallic acids, and is thus distinguished from *Winter's Bark*, by giving no re-action with ferric salts.

NATURAL ORDER. POLYGALACEÆ.

SENEGÆ RADIX. The dried root of *Polygala Senega*, imported from North America.

Botanical Source.—The *Polygala Senega* is a perennial plant with short herbaceous stems, and spikes of white flowers resembling those of the ordinary *Milk-wort*. It grows in rocky places, chiefly in the Western States.

Description.—A knotty root-stock, with contorted roots, yellowish grey in colour, and characterised by a

keel-shaped ridge running like a shrunken sinew through the principal root. The bark has a sourish odour and an acrid taste, causing a flow of saliva.

Uses.—Diuretic, diaphoretic, and stimulant, but in large doses emetic and cathartic. Useful in bronchial affections and whooping cough.

Chemistry.—Senega root contains a little volatile oil, traces of resin, gum, salts of malic acid, and sugar, but appears to owe its medicinal properties to *Polygalic Acid*, or *Senegin*, of which it yields $5\frac{1}{2}$ per cent. It is closely allied to *Saponin*, and is amorphous, and insoluble in ether and cold water; it forms a frothing solution with boiling water, and dissolves in alkaline liquids with a greenish yellow colour.

Infusum Senegæ.

Senega Root, bruised	$\frac{1}{2}$ ounce
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Dose—1 to 2 ounces.

Tinctura Senegæ.

Senega, in coarse powder	$2\frac{1}{2}$ ounces
Proof Spirit	ad 20 fluid ounces

Macerate 48 hours, then percolate.

Dose— $\frac{1}{2}$ to 2 drachms. (7 grains of Senega in 1 fluid drachm.)

Adulteration.—The root of *Panax quinquefolium* (American Ginseng) is a spindle-shaped root, quite distinct in appearance from true Senega.

RADIX KRAMERIÆ, Rhatany Root.
The dried root of *Krameria triandra*, from Peru.

Botanical Source.—The *Krameria triandra* is a small shrub with silver-grey leaves and red flowers, growing in sandy and elevated spots in Peru and Bolivia.

Description.—It is about an inch in diameter, with numerous branches, reddish brown rough exterior, and yellowish red interior; the bark being easily separable from the wood.

Uses.—Astringent, like catechu. A constituent in most astringent dentifrices.

Chemistry.—The only part of the drug having active properties is the cortical portion, which contains about 20 per cent. of a peculiar form of tannin called Rhatania-tannic Acid, closely related to catechu-tannic acid. Rhatany also contains wax, gum, and uncrystallisable sugar.

The root contains no gallic acid.

Rhatania-tannic acid does not give a precipitate with tartar-emetic, and is converted by boiling with dilute acids into sugar and *rhatany red* $C_{26}H_{22}O_{11}$.

Extractum Krameriz.

Rhatany, in coarse powder 16 ounces
Distilled Water 1½ gallon or q.s.

Dose—5 to 20 grains.

Infusum Krameriz.

Rhatany Root, bruised ½ ounce
Boiling Distilled Water 10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Incompatibles—Salts of iron, lime water, and all substances acted on by vegetable astringents.

Dose—I to 2 ounces.

Pulvis Catechu Compositus. (1 in 5.)

Tinctura Krameriz.

Rhatany, in coarse powder 2½ ounces
Proof Spirit ad 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—½ to 2 drachms. (7 grains of Rhatany in 1 fluid drachm.)

Substitutions.—There are several other species, the chief of which is the Savanilla or New Granada Rhatany, derived from the *Krameria tomentosa*, which

is in straight pieces of a dull purplish brown colour, with a thick smooth cortical portion marked with longitudinal furrows.

Sections of Peruvian Rhatany are turned grey by ferric chloride, whereas Savanilla is turned violet.

NATURAL ORDER.—GUTTIFERÆ.

This order is characterised by a combination of opposite and leathery leaves, indefinite hypogynous stamens, and sessile radiant stigmas.

CAMBOGIA, Gamboge. The gum-resin obtained from *Garcinia Morella* (var β . *Pedicellata*) imported from Siam.

Botanical Source.—The plant yielding Gamboge is a tree with laurel-like leaves, and small yellow diœcious flowers.

Collection.—The Gamboge is chiefly secreted in large vessels contained in the bark. It is obtained by making an oblique incision in the bark, extending half way round the trunk, and fixing a joint of bamboo at the lower end of the cut to receive the sap which slowly exudes for several months; this forms ripe gamboge.

Sometimes the juice obtained from the broken leaves and twigs is allowed to flow into cocoa-nut shells, and afterwards removed to earthen vessels, in which it hardens by exposure, forming lump gamboge. Each tree yields on an average a pipe about 20 inches by $1\frac{1}{2}$ inch, and suffers no injury provided the tapping is not more frequent than every other year.

Uses.—Drastic, purgative, and anthelmintic. Useful in obstinate costiveness and dropsy. Dose 1 to 4 grains.

Chemistry.—The pipe-gamboge of Siam has been found by Sir R. Christison to contain 72·2 per cent. of resin, 23 per cent. of gum, and 4·8 per cent. of moisture. The resin forms coloured solutions with ammonia or the fixed alkalies, and gives a copious yellow precipitate with *basic* acetate of lead and brown precipitates with solutions of iron and copper.

Specially Distinctive Tests. — (1.) Imperfectly soluble both in chloroform and ether.

(2.) The ether solution, after filtration, does not become turbid with alcohol.

(3.) Imperfectly soluble in alcohol, and the solution gives a yellow but clear mixture with ammonium hydrate.

(4.) The filtered alcoholic solution evaporated, and a portion of the residue treated with alcoholic ferric chloride (1 in 10), gives a dark colour.

(5.) Alcoholic plumbic acetate gives no precipitate with the remainder of the residue dissolved in alcohol.

Pilula Cambogiæ Composita.

Gamboge	1 ounce
Barbadoes Aloes	1 ounce
Comp. Powder of Cinnamon	1 ounce
Hard Soap, in powder	2 ounces
Syrup.....	q.s.

Dose—5 to 10 grains. (1 grain each Aloes and Gamboge in 6 grains nearly.)

Varieties.—(1st). *Pipe Gamboge*, in sticks or cylinders, bright yellow in colour, and varying up to 2 inches in thickness. They are striated externally by the action of the bamboo moulds.

(2nd). *Lump or Cake Gamboge*, in round cakes or masses of several pounds' weight, and inferior in quality.

(3rd). *Coarse Gamboge*, simply fragments and inferior pieces, much mixed and very dirty, and of low value.

Adulteration.—Pieces of wood and starch; an emulsion made with boiling water, and cooled, does not become green with the solution of iodine (absence

of the latter). The Cambodians themselves sometimes adulterate the lump variety with rice flour or the powdered bark of the tree. The adulterations all remain undissolved after the gum resin has been successively exhausted by alcohol and water.

GARCINIÆ OLEUM, Mangosteen Oil, *Kokum Butter*.—A concrete oil obtained by bruising the seeds of *Garcinia Indica*, and subsequently boiling them with water, when the fat rises and may be skimmed off. It is used in the Pharmacopœia of India for making suppositories, and it has been imported to serve as an ingredient in artificial butter.

NATURAL ORDER.—DIPTEROCARPEÆ.

BALSAMUM DIPTEROCARPI, Gurjun Balsam, *Wood Oil*.—An oleo-resin obtained from *Dipterocarpus turbinatus* and several other allied species by incision and heat. Imported from Singapore and also from Southern India.

This oleo-resin contains about 45 of volatile oil and 55 of resin. It is used as a substitute for Balsam of Copaiba, from which it may however be distinguished

by heating in a sealed glass tube to 428° F., when it will become a solid mass, while the Copaiba remains fluid.

NATURAL ORDER.—MALVACEÆ.

Herbs, shrubs, or trees, with alternate and usually palmated and stipulate leaves; inflorescence axillary—calyx inferior, persistent and valvate in æstivation; corolla regular and twisted in æstivation; stamens indefinite, monadelphous and hypogynous, with one-celled reniform anthers; pistil usually syncarpous with a branched stigma; fruit a carcerule or more rarely a capsule.

ALTHÆÆ RADIX, Marshmallow Root, *Racine de Guimauve*.—The scraped and dried root of *Althæa officinalis* from the South of France and Spain; also cultivated in Britain.

Botanical Source.—The true marshmallow is a herb with a perennial root and erect pubescent stem 2 to 3 feet high. The leaves are ovate in shape, having a serrate margin and two distinct lobes standing out from the broadest portion, the whole being downy on both sides. The inflorescences are axillary cymes of a

pale pinkish blue. The flowers are regular with an epicalyx, monadelphous stamens and carcerule fruit.

The plant is to be distinguished from the common mallow of our road sides (*Malva sylvestris*) which has palmately-lobed leaves, *not pubescent*, and pink flowers, and is altogether a lower growing plant.

Collection.—The roots of the biennial cultivated plant are collected and the outer coating and rootlets are scraped off before drying.

Description.—White sticks about the thickness of a finger, deeply furrowed longitudinally and marked here and there with brown scars.

Uses.—Demulcent and emollient; used in making jujubes.

Chemistry.—The chief value of the root lies in the large quantity of mucilage it yields, about 25 per cent. This is supposed to be *bassorin*, $C_{12}H_{20}O_{10}$, differing from arabin in being precipitated by normal plumbic acetate. It also contains about 2 per cent. of *asparagin*, a neutral and inert body, together with starch and sugar.

Syrupus Althææ (*Pharm. Lond.*)

Macerate $1\frac{1}{2}$ ounce in one pint of cold water, and then add 3 lbs. sugar and $\frac{1}{2}$ fluid drachm rectified spirit to each fluid ounce. **DOSE**—1 to 5 fluid drachms.

GOSSYPIUM, Cotton Wool, *Carded Cotton*.—The hairs from the *testa* of various species of gossypium ; imported chiefly from America, and carded.

Description.—Under the microscope cotton fibres are seen to be transparent, flattened, and peculiarly twisted, and they are thus easily distinguished from flax and silk.

Uses.—As a dressing for wounds and burns.

Chemistry.—Cotton is nearly pure *cellulose*, $C_6H_{10}O_5$, by treatment with a mixture of strong nitric and sulphuric acids, and subsequent copious washing with water, it forms nitro-cellulose in one of three degrees of substitution. The dinitro-cellulose, $C_6H_8(NO_2)_2O_5$, is *pyroxylin*, B.P., which is soluble in a mixture of alcohol and ether to form *collodion*, while the trinitro-cellulose, $C_6H_7(NO_2)_3O_5$, is the ordinary explosive known as gun-cotton.

NATURAL ORDER.—STERCULIACEÆ.

[*Ascribed also to* BYTTNERIACEÆ.]

THEOBROMÆ SEMINA, Cocoa Nibs, *Cacao*.—The slightly roasted and partially

crushed seeds of *Theobroma Cacao*. Imported in their raw state from Mexico and the West Indies.

Botanical Source.—The *T. Cacao* is a small tree bearing large capsules filled with seeds, which each possess a succulent and membranous arillode.

Description.—Found in commerce in “nibs,” and also in powder as prepared cocoa, usually then deprived of part of its fat and mixed with starch and sugar.

Uses.—Dietetic; nourishing without astringency.

Chemistry.—The seeds contain about 88 per cent. of actual kernel, and this yields about 50 to 55 per cent. of fat, 22 to 24 of sugar, 17 of vegetable albumen, and 1 to 1.5 per cent. of *Theobromine*, $C_7H_8N_4O_2$. This latter is an alkaloid closely resembling *Caffeine*.

THEOBROMÆ OLEUM, Cocoa Butter.—A concrete oil obtained in Britain by expression and heat from the ground seeds of *Theobroma Cacao*.

Description.—A very pale yellowish fat with a pleasant odour of chocolate, having a specific gravity of .961, and fusing between 84° and 86° F.

Chemistry.—*Cocoa Butter* is chiefly *stearin* with a little *palmatin* and *olein*, and is therefore capable of saponification by alkalies. It is entirely soluble in twice its weight of *benzol* at 50° F. It enters into the composition of

Suppositoria—

Acidi Tannici.....	1 in 2
Hydrargyri.....	1 in 2
Morphiæ.....	1 in 2
Plumbi Composita	4 in 9

NATURAL ORDER.—SAPINDACEÆ.

This order is that to which the well-known horse-chesnut tree (*Æsculus Hippocastanum*) belongs.

GUARANA, Guarana Bread.—The pounded and prepared seeds of *Paullinia sorbilis*, imported from Brazil.

Collection.—The seeds are pounded into a soft mass and then made into rolls or cakes and dried.

Description.—In heavy dark red granular-looking cakes or masses, like large sausages.

Uses.—Recommended in nervous headache and some kinds of neuralgia.

Chemistry.—The active principle is *Theine* or *Caffeine*, which exists in the large proportion of 5 per cent.

NATURAL ORDER.—LINACEÆ.

This order is allied to the Malvaceæ, but has exstipulate leaves, calyx with imbricate æstivation, and stamens definite, and only slightly monadelphous at the bottom. The ovaries have their styles, as well as their stigmas, distinct. The order is remarkable for the fine development of the fibres of the endophlœum.

LINI SEMINA, Linseed.—The seeds of *Linum Usitatissimum*, cultivated in Britain.

Botanical Source.—The *L. usitatissimum* is an annual herb about $1\frac{1}{2}$ foot high with linear-lanceolate, sessile, glabrous and alternate leaves. The flowers are in terminal cymes, and the corolla is sky-blue in colour, with petals having a twisted æstivation. It is extensively cultivated in Europe for the manufacture of linen and flax, both of which are spun from the inner bark fibres, the latter being separated by macerating the plant in water and then stripping off the bark and beating with a flat stick till the fibres separate.

Description.—Small flat oval seeds with acute edges and a smooth polished brown testa and yellowish cotyledons.

Uses.—Emollient and demulcent.

Chemistry.—The testa of the seeds contains about 15 per cent. of the special mucilage already described under *Althæa* (page 73), while the embryo or body of the seed yields about 30 per cent. of fixed oil and 25 per cent. of vegetable albumen and some sugar.

Farina Lini.

The pulverised cake of linseed left after expression of the fixed oil. This article, known as *Linseed Meal*, is used in making all the official *Cataplasmata* except *C. Fermenti*.

Cataplasma Lini.

Linseed Meal	4 ounces
Olive Oil	$\frac{1}{2}$ fluid ounce
Boiling Water	10 fluid ounces

The water extracts the mucilaginous matter from the meal, and causes it to swell, while the oleaginous matter which has been removed in the manufacture of the meal is restored by the addition of the more elegant and savoury olive oil.

Infusum Lini.

Linseed	160 grains
Fresh Liquorice Root, sliced.....	60 grains
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel for 4 hours, and strain.

Incompatibles—Most metallic salts.

Adulteration.—Linseed simply crushed is usually fairly good, but the meal is subject to adulteration with mustard, rape, and cotton seed cakes. The two former may be recognised by the development of acrid essential oils when the meal is macerated in water, and the latter by the microscope. A cooled decoction of pure linseed meal should give no blue with iodine.

LINI OLEUM, Linseed Oil.—The oil expressed without heat from the seeds of the plant above described.

Description.—A yellow viscid oil, with a slight but somewhat sickly odour, which has a specific gravity of .902, and dries by exposure to the air. *Boiled linseed oil* is made by heating the ordinary oil with oxide of lead. It dries much more rapidly than the raw oil. An inferior oil is obtained by expressing at 200° F.

Uses.—Made into an emulsion with lime water, it affords an excellent application to burns and scalds. It is the chief oil used in painting.

Chemistry.—The raw oil is chiefly olein, with a little stearin and myristin. When saponified, and the soap decomposed by acids, it yields 95 per cent. of fatty

acids, chiefly linoleic acid, $C_{16}H_{26}O_2$. This acid is peculiar to drying oils, and is not even homologous with the ordinary oleic acid. It is converted by the action of the air into oxylinoic acid, $C_{16}H_{26}O_5$, which is resinous. Boiled oil prepared as above rapidly changes into a solid transparent varnish called *Linoxyn*, $C_{32}H_{54}O_{11}$. Linseed oil is soluble in alcohol and ether, and when treated with a caustic alkali it becomes blood red. It is not affected by nitric acid of 1.18 sp. gr. (distinction from hemp seed oil, which turns green), but is turned green by sulphuric acid, sp. gr. 1.47. Its solubility in alcohol would serve to detect it if added to olive oil.

NATURAL ORDER.—ZYGOPHYLLACEÆ.

GUAIACI LIGNUM, *Guaiacum* Wood, *Lignum Vitæ*.—The wood of *Guaiacum officinale*, imported from St. Domingo and Jamaica, and reduced to chips and coarse powder by the turning lathe.

Botanical Source.—The *G. officinale* is a large evergreen tree, with paripinnate leaves and pale blue flowers. Another source of the wood of commerce is the *G. sanctum*.

Description.—Greenish brown chips, becoming more green by exposure to light, and very heavy, having a sp. gr. of 1.33, and consequently sinking in water.

Uses.—An ingredient in Decoctum Sarzæ Compositum.

GUAIACI RESINA, Guaiacum Resin.

—A resin obtained by exudation, incision or heat, from the stem of *G. officinale* above described.

Collection.—By exudation, either naturally or after incision, the resin is obtained in small masses hardened by spontaneous evaporation. This is the fine *Guaiacum in tears*. The more ordinary (*Lump Guaiacum*) is obtained by boiling the chips in salt and water and skimming off the resin as it rises to the surface, or by cutting the wood into billets and boring a hole through longitudinally, heating one end in the fire, when the melted resin flows out of the other end. In this way the wood will yield about 25 per cent.

Description.—In tears or brittle masses of a greenish brown colour, with a resinous fracture and translucent edges; as usually seen in powder it is

greyish green. When placed in the mouth it softens, and produces after the first moment an acrid sensation. An alcoholic solution of the resin turns the inside of a potato-peeling blue.

Uses.—Acrid stimulant and alterative diaphoretic. Given in chronic forms of rheumatism, and formerly used for secondary syphilis. Dose, 20 to 30 grains.

Chemistry.—The active principles are Guaiaconic Acid, $C_{38}H_{40}O_{10}$ (70 per cent.), and Guaiaretic Acid, $C_{20}H_{26}O_4$ (10 per cent.). They are isolated by extracting the resin by alcoholic potash and filtering, when the solution deposits crystals of potassium Guaiaretate, from which the acid may be liberated by hydrochloric acid. The alcoholic mother liquor is then acidulated with hydrochloric acid, and the mass which separates is washed with water and extracted by ether, which leaves on evaporation the *Guaiaconic Acid*. It is to this latter that the property of turning green in the presence of oxidising agents possessed by Guaiacum is due. Guaiacum may be detected in any mixture by evaporating to dryness and distilling with a gradually-increasing heat, when it yields first a colourless aromatic liquid which gives a green with potassium hydrate, and

afterwards a sublimate of pearly crystals, which are turned blue by warm sulphuric acid and green by ferric chloride.

Specially Distinctive Tests.—*Ordinary Guaiacum* is—1. Completely soluble in chloroform; soluble in ether, and the latter solution is not rendered turbid on the addition of alcohol.

2. It is perfectly soluble in alcohol, and the solution is coloured blue by alcoholic ferric chloride (1 in 10), and is also precipitated by saturated alcoholic plumbic acetate.

3. Sulphuric acid dissolves the solid resin to a cherry-red solution. *Peruvian Guaiacum* differs by giving a greenish brown with the ferric chloride, and the plumbic acetate precipitate re-dissolves on boiling, giving a solution which is coloured red by bromine in chloroform (1 in 20).

Mistura Guaiaci.

Guaiacum Resin, in powder	$\frac{1}{2}$ ounce
Sugar	$\frac{1}{2}$ ounce
Gum Acacia, in powder	$\frac{1}{4}$ ounce
Cinnamon Water.....	20 fluid ounces

Dose— $\frac{1}{2}$ to 2 ounces (11 grains of Guaiacum Resin in 1 fluid ounce).

Tinctura Guaiaci Ammoniata.

Guaiacum Resin, in fine powder 4 ounces
 Aromatic Spirit of Ammoniaad. 20 fluid ounces

Macerate 7 days.

Incompatibles—Acids; acidulous, earthy and metallic salts.

Dose— $\frac{1}{2}$ to 1 drachm (11 grains of Guaiacum Resin in 1 fluid drachm).

Guaiacum Resin is also contained in Pilula Hydrargyri Subchloridi Composita (1 in $2\frac{1}{2}$).

Adulteration.—The chief admixture to which this body is subject is *common resin*, detected by pouring a tincture into water, and then carefully adding solution of potassium hydrate until the solution just clears. If common resin be present a further addition of potash will cause a precipitate. The smell on heating is also a simple guide.

NATURAL ORDER.—RUTACEÆ.

Trees or shrubs (rarely herbs) with exstipulate and dotted leaves and perfect flowers; calyx 4—5 segments, imbricate; corolla 4—5 petals usually valvate; stamens, usually 5 or 10, hypogynous; pistils superior syncarpous 2—5 carpels, more or less inclined to be distinctly lobed; fruit capsular.

RUTÆ OLEUM, Oil of Rue.—The oil distilled in England from *Ruta graveolens*.

Botanical Source.—The common *Rue* is a small evergreen under-shrub with pale green bipinnate leaves, with the leaflets thick and inclined towards an obovate shape. The flowers are greenish yellow, and the inflorescence is a corymbose cyme. The stamens are hypogynous and the ovary 4-5 lobed.

Collection.—The plant is cut after flowering when the fruits are nearly ripe, and the whole cuttings are distilled with water.

Description.—Pale yellow, with a characteristic odour. Sp. gr. '911.

Uses.—Anti-spasmodic and emmenagogue. Used in hysteria, epilepsy, and flatulent colic. Externally it is an acrid stimulant and rubefacient. There is no dose mentioned in the B.P., but from 1 to 3 minims may be administered.

Chemistry.—Its chief constituent is *Euodic aldehyd*, $C_{11}H_{22}O$, together with a hydrocarbon isomeric with turpentine, and *lauric aldehyd*, $C_{12}H_{24}O$. By prolonged exposure to oxidising agents (as HNO_3) it forms *rutic acid*, $HC_{10}H_{19}O_2$, with a little *pelargonic acid*, $HC_9H_{17}O_2$.

CUSPARIÆ *vel.* ANGOSTURÆ
CORTEX, Cusparia or Angostura Bark.

—The dried bark of *Galipea Cusparia*, imported from tropical South America.

Botanical Source.—The *G. Cusparia* is a small tree growing chiefly in Venezuela.

Description.—In slightly quilled or flattish pieces about $\frac{1}{4}$ inch thick, greyish white externally and cinnamon-coloured internally. The edges are *sharp* and *bevelled* in a very characteristic manner, and the internal surface is easily capable of being split into laminæ. A drop of nitric acid colours the internal surface a very *dull* red. The bark of *Strychnos Nux vomica* containing *Brucia* is turned blood red on the addition of nitric acid.

Uses.—Stimulant and tonic. In hot climates it appears to be somewhat anti-periodic, as it has been employed with success in fevers of a malignant type. 10 to 40 grains of the powdered bark may be given.

Chemistry.—It contains about $\frac{3}{4}$ per cent. of volatile oil, about $1\frac{1}{2}$ per cent. of a neutral bitter principle called *Cusparin*, soluble in alcohol and acids, but sparingly so in water, and which may be obtained by spontaneous

evaporation of the tincture; it is rendered greenish by nitric acid. Cusparia also contains a hard and soft resin, the latter coloured dark red by nitric acid. The bark is free from tannin, but gives a copious precipitate with ferric chloride.

Infusum Cuspariæ.

Cusparia Bark, in coarse powder ½ ounce
 Distilled Water, 120° 10 fluid ounces

Infuse in a covered vessel for 2 hours, and strain.

Incompatibles—Most metallic salts and tincture of galls.

Dose—1 to 2 ounces.

BUCHU FOLIA.—The dried leaves of *Barosma betulina*, *crenulata* and *serratifolia*, imported from the Cape of Good Hope.

Botanical Source.—All the species of *Barosma* are erect shrubs, with small coriaceous leaves with conspicuous oil glands on the under-surface and margin.

Description.—*Barosma betulina* is about $\frac{3}{4}$ -inch long, coriaceous obovate, with a recurved truncated apex and sharp spreading teeth.

Barosma crenulata is about an inch long, ovate-lanceolate, obtuse, minutely crenated, five nerved.

Barosma serratifolia is from 1 to 1½ inch long, linear-lanceolate, tapering equally at each end, sharply and finely serrated, three nerved.

Uses.—Stimulant, diuretic and diaphoretic, acting specially on the urinary organs. In catarrh of the bladder, enlarged prostate, gleet and stricture it is very useful. Dose of powdered leaves, 20 to 30 grains in wine.

Chemistry.—Buchu leaves contain mucilaginous matter and about 1½ per cent. of volatile oil, which, on exposure to cold, deposits crystals of Barosma camphor having a peppermint odour. .

Infusum Buchu.

Buchu Leaves, bruised ½ ounce
Boiling Distilled Water 10 fluid ounces

Infuse in a covered vessel 1 hour, and strain.

Incompatibles—Salts of iron and astringent preparations.

DOSE—1 to 4 ounces.

Tinctura Buchu.

Buchu Leaves, bruised 2½ ounces
Proof Spirit ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—1 to 2 fluid drachms (7 grains of Buchu in 1 fluid drachm.

Substitution.—The leaves of *Empleurum serrulatum*

(Rutaceæ) grow in the same localities, and are sometimes imported and sold as Buchu. They very much resemble *B. serratifolia*, but are much narrower and terminate in an acute point. They have a different odour from Buchu.

JABORANDI.—The leaves mixed with the young branches of *Pilocarpus pinnatifolius*, a native of Brazil.

Botanical Source.—The *P. pinnatifolius* is a shrub with a light grey bark and compound imparipinnate leaves with each leaflet as large or even larger than a leaf of cherry laurel. Inflorescence in racemes.

Description.—Yellowish or greenish brown leathery leaflets from 4 to 6 inches long and from $\frac{3}{4}$ to 2 inches broad with the margin entire and slightly re-curved. The apex is rounded and slightly emarginate, and the veins are well marked and terminate in an undulating line just within the margin.

Uses.—The action of Jaborandi is sialagogue (*σίαλον* saliva, and *ἀγωγός* evoking) and sudorific in large doses causing nausea. About 60 grains of the dry or $\frac{1}{2}$ grain of pilocarpia will cause profuse salivation and sweating within 15 minutes, which will continue for about 1 hour.

Chemistry.—The leaves of this plant contain an alkaloid called *pilocarpia*, to which the sialagogue and sudorific properties of the plant appear to be due. It is a colourless crystalline substance, soluble in water, alcohol, and in chloroform, forming crystalline salts with acids of which the hydrochlorate is deliquescent and soluble in alcohol, and the nitrate is permanent and insoluble in alcohol. It has been obtained (by Mr. Gerrard) by the following process:—Prepare a soft extract with proof spirit; digest this with water, filter and wash; evaporate the filtrate to a soft extract, cautiously add ammonia in slight excess, shake with chloroform, separate the latter and evaporate; the residue is impure *pilocarpia*, which may subsequently be purified by solution in acidulated water, and re-crystallisation from chloroform.

The leaves also contain a volatile oil, tannic acid, and acrid resin.

Adulteration.—There are no adulterations yet known, but the leaves might very readily be mistaken for those of cherry laurel. The latter, however, are *alternate leaves*, while Jaborandi has *opposite leaflets*. The leaves of the laurel are also slightly serrate towards the apex, while Jaborandi leaflets are perfectly entire.

NATURAL ORDER.—AURANTIACEÆ.

Trees or shrubs closely allied to Rutaceæ by abounding in oil vesicles, but they are distinguished from every other order by their characteristic fruit—namely, the *hesperidium*.

AURANTII CORTEX, Bitter Orange Peel.—The rind of the ripe fruit *Citrus vulgaris*, variety *Bigaradia*, imported from Seville and the south of Europe generally.

Botanical Source.—The orange tree is from 16 to 20 feet high, with simple ovate-acuminate leaves articulated to a winged petiole, and fine large white fragrant flowers. The fruit is a *hesperidium*, having the epicarp and mesocarp united in the rind, and the endocarp running in and dividing the pulp into lobes. The *C. bigaradia*, or bitter orange, has axillary spines on its branches, and a small deep-coloured fruit, while the *C. Aurantium*, or sweet orange, has no spines and a larger and paler fruit.

Description.—Good orange peel should be very

thinly cut, and carefully dried. It should not be of too dark a colour, and should have as little white rind as possible.

Uses.—Bitter stomachic, tonic, and stimulant.

Chemistry.—The peel contains a bitter glucoside called *hesperidin*, a trace of gallic acid, and a little volatile oil.

Infusum Aurantii.

Bitter Orange Peel, cut small	½ ounce
Boiling Distilled Water	10 fluid ounces

Infuse 15 minutes in a covered vessel, and strain.

Incompatibles—Salts of iron.

DOSE—1 to 2 ounces.

Infusum Aurantii Compositum.

Bitter Orange Peel, cut small	¼ ounce
Fresh Lemon Peel, „	60 grains
Cloves, bruised.....	30 grains
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel 15 minutes, and strain.

Incompatibles—Salts of iron.

DOSE—1 to 2 ounces.

Syrupus Aurantii.

Tincture of Orange Peel	1 fluid ounce
Syrup.....	7 fluid ounces

DOSE—1 drachm.

Tinctura Aurantii.

Bitter Orange Peel, bruised..... 2 ounces
 Proof Spirit.....ad. 20 fluid ounces

Macerate 7 days, then strain, press, and filter.

DOSE—I to 2 drachms.

Tinctura Aurantii Recentis.

Bitter Orange Peel, fresh 6 ounces
 Rectified Spiritad. 20 fluid ounces

Carefully cut from the orange the coloured part of the rind, in thin slices, and macerate 6 ounces of this in a pint of spirit for 7 days. Then pour off the liquid, press the dregs, mix the liquid products and filter. Finally, add sufficient spirit to make 1 pint.

DOSE—I or 2 drachms.

Vinum Aurantii.

British wine, containing 12 per cent. of alcohol, made by the fermentation of a saccharine solution, to which the fresh peel of the bitter orange has been added.

Also contained in Infusum Gentianæ Compositum, 120 grains to 1 pint; Mistura Gentianæ, 60 grains to 1 pint; and Tinctura Gentianæ Composita, $\frac{3}{4}$ ounce to 1 pint.

AURANTII FLORES, Orange Flowers.—The fresh flowers of both *Citrus Bigaradia* and *C. Aurantium*. They are used in the following preparations :—

Aqua Aurantii Floris.

The distilled water of the flowers of *Citrus Bigaradia*, and of *C. Aurantium*, imported from France.

Syrupus Aurantii Floris.

Orange Flower Water 8 fluid ounces

Distilled Water q.s. or 16 fluid ounces

Refined Sugar 48 ounces

Product 4½ lbs. by weight, or 3 lb. 6 oz. by measure,
sp. gr. 1.330.

Dose—1 drachm.

AURANTII FLORUM OLEUM, Oil of Neroli.—The oil prepared by distilling the fresh flowers with water in a copper still. It is commonly adulterated by the oil obtained by distilling the leaves instead of the flowers, called *Huile de petit grain*. Pure oil should have a sp. gr. of .889 at 52° F., and should assume an intense permanent crimson colour when shaken with sodium bisulphite.

LIMONIS CORTEX, Lemon Peel.—The outer part of the rind of the fresh fruit of *Citrus limonum*, imported from Southern Europe.

Botanical Source.—The *C. Limonum* is a much branched and thorny shrub, with oval leaves having serrulate margin and a winged petiole. The fruit is a pale yellow egg-shaped hesperidium.

Uses.—Aromatic tonic, similar to orange peel.

Chemistry.—Similar to that of orange peel, and is employed in the following *preparations*:—Infusum Aurantii Compositum, 120 grains to 1 pint; Infusum Gentianæ Compositum, $\frac{1}{2}$ ounce to 1 pint.

Syrupus Limonis.

Fresh Lemon Peel	2 ounces
Lemon Juice, strained	20 fluid ounces
Refined Sugar	36 ounces

Product, 3½ lbs. by weight, or 2 lb. 9 oz. by measure;
sp. gr. 1.340.

Dose—1 drachm.

Tinctura Limonis.

Fresh Lemon Peel, sliced thin.....	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 7 days, strain, press, and filter.

Dose—½ to 2 drachms (7 grains of lemon peel in 1 fluid drachm).

LIMONIS OLEUM, Oil of Lemon.—

The oil expressed or distilled from fresh lemon peel, chiefly produced in and imported from Sicily.

It is pale yellow in colour, with a sp.gr. of .850. It consists chiefly of a hydrocarbon isomeric with turpentine, $C_{10}H_{16}$, and is soluble in 7 parts of rectified spirit.

By long keeping it deposits a fatty crystalline body called *citroptene*.

It is an ingredient in Linimentum Potassii Iodidi c Sapone and Spiritus Ammoniæ Aromaticus.

LIMONUM SUCCUS, Lemon Juice.

—The freshly expressed juice of the ripe fruit of Citrus Limonum.

Uses.—Refrigerant, ant-alkaline, and anti-scorbutic—also used for making citric acid (for process and description of this body see the author's "Pharmaceutical Chemistry."

Chemistry.—Lemon juice contains as its active principle about 7 per cent. of citric acid, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7 \cdot \text{H}_2\text{O}$, together with a little sugar, gum, and traces of malic acid. Artificial lemon juice may be made by dissolving $1\frac{1}{2}$ ounce of citric acid and $\frac{3}{4}$ ounce of sugar in 1 pint of water, and adding 3 or 4 drops of oil of lemon. The average fresh juice shows a sp. gr. of 1.039, and is supposed to contain $32\frac{1}{2}$ grains citric acid per fluid ounce.

Table of the amount of lemon juice required to saturate 20 grains respectively of the following alkalies.

TWENTY GRAINS.	LEMON JUICE.	CITRIC ACID.
Potassium Bicarb.	3½ fluid drachms	= 14 grains
Potassium Carb.	4 fluid drachms	= 17 grains
Sodium Bicarb.	4 fluid drachms	= 17 grains
Sodium Carb.	2½ fluid drachms	= 10 grains
Ammonium Carb.	6 fluid drachms	= 24 grains

Beside the species already mentioned we may note the *Citrus Bergamia* which yields the oil of bergamot, and the *Citrus Limetta* from which the well-known anti-scorbutic lime-juice is derived.

BELÆ FRUCTUS.—The dried half-ripe fruit of *Egle Marmelos*, from Malabar and Coromandel.

Description.—A roundish fruit about the size of a large orange, with a hard woody rind; usually imported in dried slices, or in fragments consisting of portions of the rind and adherent pulp and seeds. Rind about a line and a half thick, covered with a smooth pale brown or greyish epidermis, and internally, as well as the dried pulp, brownish red or cherry-red. The moistened pulp is mucilaginous.

Mangosteen fruit (*Garcinia Mangostana*) is sometimes substituted for it, but this has a much thicker rind and no adhering pulp. The fruit of *Bael* is a hesperidium.

Use.—Held in repute in India as a remedy for dysentery, but found of little use in this country.

Chemistry.—The drug appears to contain only a very slight trace of tannin. The pulp and seeds furnish a large quantity of mucilage and a little sugar.

Extractum Belæ Liquidum.

Bael Fruit.....	16 ounces
Distilled Water	12 pints
Rectified Spirit	2 fluid ounces
Product, 16 fluid ounces (1 fluid ounce of extract is equal to 1 ounce of the fruit).	

DOSE—1 to 2 drachms.

NATURAL ORDER.—VITACEÆ.

[AMPELIDACEÆ.]

The Vine Order of climbing shrubs with swollen nodes and minute green flowers arranged in a thyrsus, with stamens equal in number to the petals, and a succulent superior fruit called a *nuculanium*.

UVÆ PASSÆ, Raisins.—The ripe fruit of *Vitis vinifera*, dried by the sun or by artificial heat. Imported from Spain. They enter into the composition of Tinctura Cardamomi Composita, and Tinctura Sennæ Composita.

NATURAL ORDER.—SIMARUBACEÆ.

SIMARUBÆ RADICIS CORTEX,
Simaruba Root Bark.—The dried cortical portion of the root of *Simaruba amara*, imported from Jamaica.

Description.—In long strips of thin bark folded upon themselves, of a pale greyish colour, and very fibrous in structure.

Uses.—It may be used in the same cases as quassia. It gained at one time some repute in the treatment of diarrhœa and dysentery, but is now little employed. Dose of the powder, 10 to 40 grains.

Chemistry.—The active principle is quassin, and it also contains a little volatile oil, resin, and ulmin.

QUASSIÆ LIGNUM, Quassia Wood.
—The wood of *Picræna excelsa*, imported from Jamaica.

Botanical Source.—A tall tree growing on the mountains of Jamaica.

Description.—A dense, tough, yellowish white, odourless, intensely-bitter wood, generally in chips or raspings, but sometimes in billets.

Uses.—Quassia is a pure bitter, similar in operation but rather more powerful than gentian. It is employed in stomach complaints of an atonic or functional character. As it is not affected by iron or mercury, it is a useful vehicle for the administration of the salts of those metals. It is fatal to flies and other small insects.

Chemistry.—The active bitter principle is a neutral, odourless, crystallisable substance called Quassin. $C_{10}H_{12}O_3$, which is yielded to both dilute alcohol and water. It is insoluble in ether, and gives a precipitate with tannic acid. The wood yields about 1-10th of this substance. It is entirely free from astringent matter.

Extractum Quassiæ.

Quassia Wood, rasped	16 ounces
Distilled Water	q.s.

100 lbs. of wood yield 2 and 1-12th lbs. of extract.

DOSE—3 to 5 grains (1 grain is equal to 8 fluid ounces Inf. Quassia).

Infusum Quassiæ.

Quassia, in chips	60 grains
Cold Distilled Water	10 fluid ounces

Infuse in a covered vessel for $\frac{1}{2}$ hour, and strain.

Incompatibles—Salts of lead and silver.

DOSE—1 to 2 ounces.

Tinctura Quassiæ.

Quassia $\frac{3}{4}$ ounce
 Proof Spirit.....ad. 20 fluid ounces

Macerate 7 days, and filter.

Dose— $\frac{1}{2}$ to 2 drachms (2 grains of Quassia to 1 fluid drachm.)

SUB-CLASS II.—CALYCIFLORÆ.

Plants having the calyx united, the corolla distinct, and the stamens perigynous or epigynous. This sub-class has two sub-divisions:—

1. Perigynæ, having the stamens perigynous and the ovary superior.
2. Epigynæ, with the stamens epigynous and the ovary inferior.

SUB-DIVISION I.—PERIGYNÆ.

NATURAL ORDER.—RHAMNACEÆ.

RHAMNI FRUCTUS, Buckthorn Berries.—The dried fruits of *Rhamnus cathartica*, from indigenous plants.

Botanical Source.—The *R. cathartica* is a robust shrub with thorns. The leaves are small oval, acutely serrate, with 3 or 4 simple prominent convergent veins and stipulate. Flowers diœcious and fruit baccate.

Collection.—In September, when the fruit ripens.

Description.—Small round fruits resembling a black currant, with a greenish purple juice, and containing 4 triangular seeds.

Uses.—A violent hydragogue (ῥδωε water, ἀγωγός evoking) purgative, with a tendency to cause griping.

Chemistry.—Contains two crystalline glucosides, Rhamnine, $C_{18}H_{22}O_{10}$, and Rhamnegine, $C_{24}H_{32}O_{14}$. The former is only soluble in hot alcohol, and insoluble in ether; while the latter is soluble in water also. Both give deep orange solutions when treated with alkalis. The purgative principle of the berries is, however, supposed not to be due to these glucosides, but to a bitter principle as yet not properly investigated.

Rhamni Succus (Buckthorn juice).—The recently expressed juice of the ripe berries, used in making—

Syrupus Rhamni.

Buckthorn Juice	4 pints
Ginger, sliced	$\frac{3}{4}$ ounce
Pimento, bruised	$\frac{3}{4}$ ounce
Refined Sugar	5 lbs.
Rectified Spirit.....	6 fluid ounces

Dose—1 drachm. Sp. gr. 1.320.

FRANGULÆ CORTEX, Alder Bark.

—The bark from the trunk and larger branches of *Rhamnus frangula*, growing in England and imported from Germany.

Botanical Source.—The *R. frangula* is distinguished from the *R. cathartica* by being more tree-like in habit, having larger leaves, which are entire and have 7 pairs of prominent veins, and by the fruit only having 2 or at most 3 *flat* seeds.

Description.—The bark is greyish externally, and brownish yellow internally, smooth, compact, and fibrous, with a sweetish taste.

Uses.—A purgative, official in Germany, and esteemed for its combined certainty and mildness. The best preparation is a watery fluid extract 1 in 1.

NATURAL ORDER.—AMYRIDACEÆ.

[BURSERACEÆ.]

OLIBANUM, Frankincense.—The gum resin obtained by incision and exudation from the stem of various species of *Boswellia*. Really produced in Africa and Southern Arabia, but imported *via* India.

Description.—Usually in pale yellowish or nearly colourless distinct pear-shaped tears, but sometimes stalactiform and slightly agglutinated. Always covered externally by a fine white dust, and even where that is wiped off the tears appear translucent and milky, and have a characteristic turpentine-like odour. When rubbed with water it forms a white emulsion.

Uses.—Supposed to be the *Frankincense* of the Bible, and still used for burning in the Romish and Greek churches.

Chemistry.—*Olibanum* contains about 5 to 7 per cent. of volatile oil, 33 per cent. of gum identical with *arabin*, and the remainder a resin, $C_{40}H_{30}O_6$, which is not soluble in alkalies.

ELEMI.—A concrete resinous exudation probably from *Canarium commune*, chiefly imported from Manilla.

Description.—A soft adhesive and unctuous mass, becoming resinous by age. Its colour is yellowish white, and it has a fragrant odour resembling that of fennel. It should be almost entirely soluble in rectified spirit.

Uses.—*Elemi* is seldom used in British medicine, except in the form of an ointment sometimes prescribed as a stimulating application to old wounds.

Chemistry.—*Elemi* contains 10 per cent. of a fragrant colourless neutral oil, 60 per cent. of an uncrystallisable resin soluble in cold alcohol, and 25 per cent. of a crystallisable resin, called amyrin, insoluble in cold alcohol.

Specially Distinctive Tests.—1. Soluble in chloroform and ether, the latter solution not being rendered turbid by alcohol.

2. Only partially soluble in alcohol, and the filtered solution is not precipitated by alcoholic plumbic acetate (1 in 10).

3. A portion of the alcoholic solution evaporated leaves a residue insoluble in boiling sodium carbonate.

4. The alcoholic and chloroform solutions are coloured violet respectively by alcohol saturated with dry hydrochloric acid gas, and by bromine in chloroform (1 in 20).

Unguentum Elemi.

Elemi.....	¼ ounce
Simple Ointment.....	1 ounce

MYRRHA, Myrrh.—A gum-resinous exudation from the stem of *Balsamodendron Myrrha*, collected in Arabia Felix and Abyssinia.

Botanical Source.—The *B. Myrrha* is a spiny shrub with ternate leaves and nearly sessile flowers.

Collection.—Myrrh, when first exuded, is said to be of an oily, then of a buttery appearance, yellowish white, gradually assuming a golden tint, and becoming reddish as it hardens.

There are two well-known commercial varieties.

1. Turkey, the finer kind, is of two qualities—(a) picked myrrh, in large pieces; and (b) grain myrrh.
2. East Indian myrrh (*Bissa Böl*); this is an inferior quality and much adulterated. Hanbury states that there are certainly three varieties of myrrh, which may well be derived from as many distinct species of myrrh-tree. He leaves the subject open to further elucidation.

Description.—Good commercial myrrh is in irregularly shaped masses of a reddish-brown colour, and slightly translucent. It has a dull irregular fracture, and an aromatic and characteristic odour.

Uses.—Tonic and stimulant. It acts upon the mucous membrane; applied locally it is a mild astringent. The tincture when diluted forms a useful gargle in relaxed throats. Dose of powder, 10 to 30 grains.

Chemistry.—Myrrh contains 2·5 per cent. of volatile oil, 40 to 50 per cent. of gum, and the remainder resin. Both the resin and gum contain two modifications, soft and hard. Soft resin, supposed to be a mixture of hard resin and volatile oil, has a strong odour, becomes softened by moderate heat, and is insoluble in ether. Hard resin has no odour, is not so readily softened, is insoluble in ether, and soluble in caustic alkalies, forming myrrhates. The first modification of the gum, probably arabin, is soluble in water, forming precipitates with salts of silver, mercury, and lead. The second modification is insoluble in water, and similar to bassorin. The volatile oil is soluble in alcohol, ether, and fixed oils.

Specially Distinctive Tests.—Ordinary myrrh is

1. Imperfectly attacked both by chloroform and ether.
 2. Alcoholic solution gives a turbid mixture with ammonium hydrate, and no precipitate with alcoholic ferric chloride (1 in 10).
 3. Not completely soluble in ether-alcohol.
 4. Contains no sulphur, and yields no umbelliferone by dry distillation.
 5. A little of the solid extract from alcohol solution is not coloured by a saturated solution of chlorinated lime.
 6. The alcoholic solution is precipitated by alcoholic plumbic acetate.
 7. The evaporated residue from a solution in petroleum spirit is coloured violet by impure chloral hydrate.
- East Indian myrrh does not give the reactions 6 and 7.

Tinctura Myrrhæ.

Myrrh, in coarse powder	2½ ounces
Rectified Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—½ to 1 drachm (7 grains of Myrrh in 1 fluid drachm).

Myrrh is also contained in Decoctum Aloes Compositum, 3 grains to 1 fluid ounce; Mistura Ferri Composita, 6 grains to 1 fluid ounce; Pilula Aloes

et Myrrhæ, 1 in 6; Pilula Asafœtidæ Composita, 1 in $3\frac{1}{2}$; Pilula Rhei Composita, 1 in 8 nearly.

Adulterations.—Various gums and resins which may be distinguished by the transparency of their fractured surfaces and want of taste and odour. Also inferior kinds derived from other species of *Balsamodendron*, as *B. pubescens* and *B. Mukul*, which are the source of Bdellium or false myrrh, and are imported from Africa and the East Indies. Good myrrh should form a nearly white emulsion with water.

BDELLIUM.—A gum resin yielded by *Balsamodendron Mukul* and *B. pubescens*. Both species indigenous in Scinde.

Description.—In rounded pieces, of dull dark red colour; taste bitter and slightly acrid. Used as a substitute for Myrrh.

Specially Distinctive Tests.—*Indian Bdellium* behaves like myrrh as far as reactions 1, 2, and 3 already noticed.

4. It contains sulphur, detected by fusion with nitre and treating with barium chloride, but yields no umbelliferone.

5. It forms with saturated solution of sodium carbonate an emulsion which cannot be filtered.

6. The tincture is either not affected at all by alcoholic plumbic acetate, or yields a precipitate which disappears on warming.

NATURAL ORDER.—ANACARDIACEÆ.

MASTICHE, Mastich.—A resinous exudation obtained by incision from the stem of *Pistacia Lentiscus*, produced in the island of Scio.

Botanical Source.—The *P. Lentiscus* is an ever-green diœcious shrub.

Collection.—Numerous vertical incisions are made about the middle of June in the bark of the stems and principal branches, from which the resin speedily flows, and soon hardens and dries; after 15 or 20 days it is carefully collected in baskets lined with paper or clean cotton wool.

Description.—Mastich is in smooth yellowish brittle tears, about the size of grains of mace, and covered with a dusty powder. It is entirely soluble in ether and oil of cloves, and about 90 per cent. is soluble in alcohol. the remainder being only soluble in ether and turpentine.

Uses.—Solutions of Mastich in ether and chloroform are used for stopping teeth and coating pills. Dissolved in turpentine or alcohol it forms a good common varnish.

Chemistry.—Mastich contains 90 per cent. of resin ($C_{20}H_{32}O_3$), which is soluble in alcohol, and 10 per cent. of Masticin, which is insoluble in alcohol. It also contains a very little volatile oil.

Masticin is a translucent colourless tough substance, insoluble in boiling alcohol or in caustic alkalies, but soluble in ether and turpentine.

Among other interesting plants of the Anacardiaceæ, may be noted—

Pistacia Terebinthus, which yields Chian Turpentine.

Pistacia vera, ,, ,, the Pistacia Nut.

Anacardium occidentale, ,, ,, the Cashew Nut.

Mangifera Indica, ,, ,, the Mango Fruit.

Specially Distinctive Tests.—Ordinary Mastich is—1. Entirely soluble in chloroform and ether, and the latter solution is rendered turbid by alcohol.

2. The alcoholic solution gives no turbidity with saturated solution of ferric chloride.

3. It is only partially soluble in petroleum spirit, boiling below 100° F., and the mixture is coloured reddish violet by a saturated solution of iodine in petroleum spirit, boiling at 140° F.

Bombay Mastich is—1. Soluble in chloroform and ether, and an addition of alcohol to the latter solution causes no turbidity.

2. It is perfectly soluble in alcohol, and the solution is coloured greenish brown by an alcoholic solution of ferric chloride (1 in 10).

3. An alcoholic solution gives a precipitate with a saturated alcoholic solution of plumbic acetate, which is not re-dissolved by boiling.

4. It is not soluble in saturated solution of sodium carbonate.

5. The petroleum spirit solution leaves, on evaporation, a colourless extract, in which impure chloral hydrate (containing alcoholate) gives no decided colour.

Alexandrian Mastich behaves like *Bombay*, except that the plumbic acetate precipitate dissolves by boiling, and the solution is not coloured by the addition of solution of bromine in chloroform (1 in 20).

NATURAL ORDER.—LEGUMINOSÆ

[Or FABACEÆ].

Herbs, shrubs, or trees, with usually compound, stipulate and alternate leaves; calyx monosepalous, 5 parted, persistent and inferior, odd sepal anterior; corolla (in British plants), 5 petals irregular, with odd petal posterior; stamens definite monadelphous or diadelphous; pistil, simple and superior; fruit, a legume or lomentum (rarely a drupe); seeds, exalbuminous.

The order is divided in 3 sub-orders—

1. *Papilionaceæ*, with truly papilionaceous corolla.
2. *Cæsalpinieæ*, corolla not truly papilionaceous.
3. *Mimoseæ*, with valvate æstivation and indefinite distinct stamens.

SUB-ORDER I.—PAPILIONACEÆ.

SCOPARII CACUMINA, Broom Tops.

—The dried tops of *Sarothamnus Scoparius*, cut from indigenous plants.

Botanical Source.—The common Broom is a shrub, with straight, glabrous, and angular branches. The leaves are ternate and very small, while the flowers are axillary and bright yellow, with the corolla papilionaceous, the stamens monadelphous, and the legume hairy, becoming blackish when ripe.

Collection.—The young first year's branches, which shoot up almost devoid of leaves, or with only a minute ovate and sessile leaflet, are cut, and either used at once for pressing out the juice, or dried for preservation.

Uses.—Chiefly administered as a diuretic in dropsy.

Chemistry.—The tops contain a neutral gelatinous or sometimes yellow crystalline body, called Scoparin, $C_{21}H_{22}O_{10}$, which remains, when a decoction is evaporated, until it solidifies on cooling, and the resulting gelatinous mass is well washed with cold water acidulated with hydrochloric acid. These acid liquors when distilled with sodium hydrate, and the product rectified by solution in acid and re-distillation with alkali, yield an oily volatile alkaloid, called Sparteia, $C_{15}H_{26}N_2$, sparingly soluble in water and boiling at 550° F. which forms crystalline salts with acids.

Decoctum Scoparii.

Broom Tops, dried	1 ounce
Distilled Water	20 fluid ounces

Boil 10 minutes in a covered vessel, strain and make up product to 20 fluid ounces.

Dose—2 to 4 ounces.

Succus Scoparii.

Broom Juice	3 pints
Rectified Spirit	1 pint

Filter after standing 7 days.

Dose—1 drachm to 2 drachms (1 fluid drachm is equal to 4 drachms of broom tops).

FŒNI-GRÆCISEMINA, Fenugreek.

—The seeds of *Trigonella Fœnum-græcum* from Southern Europe.

Botanical Source.—This plant is an annual herb with whitish flowers, indigenous to the Mediterranean coasts.

No proper investigation has been made of fenugreek seeds, which are only used in veterinary practice ; in curry powder, and cattle foods.

TRAGACANTHA, Gum Tragacanth.

A gummy exudation from the stem of *Astragalus verus* and other species collected in Asia Minor.

Botanical Source.—The various species of *Astragalus* are low shrubs, having pinnate leaves with linear and hispid leaflets and curious strong and spiny petioles. The flowers are papilionaceous and yellow.

Collection.—Formerly it was only the gum which exuded spontaneously that was collected, but now incisions are made near the root, and the gum flows out and dries in the fine white cakes so highly prized in the market.

Description.—Good Tragacanth is in thin flakes, dull white and translucent, and having peculiar wavy marks on their surface. It should be horny, flexible and not easily fractured, and the surface should not be shiny. When dried at 120° F. it loses 14 per cent. of water, and may then be powdered.

Uses.—As a suspender of heavy insoluble powders in water, and as a pill excipient.

Chemistry.—Placed in water tragacanth swells and absorbs 50 times its own weight, forming a clear jelly. If now 50 parts more water be added and the whole stirred and filtered it is found that a portion is soluble while the remainder remains as an insoluble jelly, rapidly drying to a very coherent mass. The soluble

gum is like *arabin*, inasmuch as it is precipitated by alcohol and ammonium oxalate, but it differs in being precipitated by normal plumbic acetate and not by borax or ferric chloride. The insoluble gum is called *Bassorin*, $C_{12}H_{20}O_{10}$, and is associated in the tragacanth with a little starch. Good samples should not yield more than 3 per cent. of ash.

Mucilago Tragacanthæ.

Tragacanth	60 grains
Distilled Water	10 fluid ounces

Pulvis Tragacanthæ Compositus.

Tragacanth, in powder	1 ounce
Gum Acacia do.	1 ounce
Starch do.	1 ounce
Refined Sugar do.	3 ounces

Dose—20 grains to 1 drachm.

Tragacanth also enters into the composition of *Confectio Opii*, 1 in 120, nearly; *Pulvis Opii Compositus*, 1 in 30.

Adulterations.—*Moussul gum*, or “hog tragacanth,” is known by its much darker colour. *Caramania Gum* and *Lead Carbonate* are both found as commercial falsifications.

GLYCYRRHIZÆ RADIX, Liquorice Root.—The dried (and fresh) underground stem and root of *Glycyrrhiza glabra*. Cultivated in England.

Botanical Source.—The *G. glabra* is a herb with a perennial root stock and compound imparipinnate leaves, usually having 1—3 oval leaflets. The inflorescence is an axillary raceme of pale blue flowers, having papilionaceous corolla and diadelphous stamens.

Collection.—The plant which is chiefly cultivated in Mitcham and in Yorkshire, is allowed to grow for four years before being touched, and the roots are then dug up after the aerial stems die off in late autumn or early winter.

Description.—The dried root sometimes occurs in long cylindrical branched pieces, but is usually found in pieces about 4 inches long and as thick as a finger: it is tough and stringy, dirty brown externally, and yellow internally. It gives with water a solution precipitated by sulphuric acid.

Uses.—The powdered root is used to dust pills with to keep them from adhering together. It is also an excellent pill excipient for soft or liquid bodies, such as oleo-resins, acting by giving consistence. It may be

employed with advantage to cover the taste of aloes and other nauseous drugs.

Chemistry.—Liquorice root contains a peculiar sweetish principle, called *glycyrrhizin*, $C_{24}H_{36}O_9$, sugar, asparagin, starch, albumen, malates and phosphates of lime and a little acrid oily matter. *Glycyrrhizin* is precipitated from a strong decoction upon the addition of an acid, or solution of acid tartrate of potassium, or normal or basic acetate of lead. It is a yellowish powder having an acid re-action and a strong sweetish-bitter taste. By boiling with a dilute acid it is resolved into a resinous amorphous bitter substance called *Glycyrretin*, the formula of which has not been settled, and an uncrystallisable sugar having the characters of glucose.

The root should be decorticated before use, as the cortical portion contains a somewhat acrid matter.

Extractum Glycyrrhizæ.

Liquorice Root, in coarse powder	16 ounces
Distilled Water	4 pints

Extractum Glycyrrhizæ Liquidum.

Macerate 16 ounces of the root in 2 pints of water for 12 hours, strain and press. Repeat this process with another 2 pints of water. Mix the liquors, heat to 212° F., strain and evaporate to a density of 1.160. Finally add 1-8th of its bulk of rectified spirit, let it stand for 12 hours and filter.

Pulvis Glycyrrhizæ Compositus.

Senna, in fine powder	2 ounces
Liquorice Root, in fine powder.....	2 ounces
Refined Sugar, in powder	6 ounces

Mix thoroughly, pass through a fine sieve, and finally rub it lightly in a mortar.

• Dose—30 to 60 grains.

Liquorice root is also an ingredient in *Confectio Terebinthinæ*, *Decoctum Sarsæ Compositum*, *Infusum Lini*, *Pilula Hydrargyri*, and *Pilula Ferri Iodidi*.

The extract enters into the composition of *Decoctum Aloes Compositum*, *Confectio Sennæ*, *Mistura Sennæ Composita*, *Tinctura Aloes*, and *Trochisci Opii*.

The *Spanish Juice* of commerce is an extract made by boiling the crushed fresh root with water in copper vessels and evaporating to a proper consistence, when it is rolled into sticks with the hands and stamped with the name of the maker.

The most famous is the "Solazzi" Juice, but there are several other varieties such as "Barracco," "Corigliano" and "Pignatelli."

INDIGO.—A blue pigment, prepared from various species of *Indigofera*.

Botanica Source.—This is chiefly *I. tinctoria*, a small shrubby plant, cultivated in Bengal and elsewhere in India.

Collection.—The plants are cut before flowering and placed in troughs covered with water. Fermentation sets in and is continued for 12 to 14 hours, until a sherry-coloured liquid is obtained, which on agitation in contact with air deposits the indigo as a precipitate. This latter is then filtered out, pressed, cut into cakes and dried.

Uses.—Solution of sulphate of Indigo is in the B.P. as a test for free chlorine, which bleaches the colour.

Chemistry.—The blue colouring matter is *Indigotin*, existing in ordinary commercial samples to the extent of 50 to 60 per cent. Its formula is C_8H_5NO , and it is thus metameric with *benzoyl cyanide*, $C_7H_5O.CN$. It fuses at 550° , and sublimes in prisms, and it is entirely insoluble in water and all ordinary solvents. Digested with strong sulphuric acid it dissolves to form a deep blue liquid called *sulphindigotic acid* (commonly called sulphate of Indigo). When treated with reducing agents in the presence of alkalies (such as ferrous sulphate and lime) Indigo dissolves to a colourless solution, and if any

article be dipped into this liquid and hung up in the air it becomes dyed blue, because the Indigo is re-deposited by oxidation.

SETÆ MUCUNÆ, Cowhage, *Cow-itch*.

—The hairs from the legumes of *Mucuna pruriens* from the West Indies.

These hairs possess the property of penetrating the skin, and cause great itching. When mixed with treacle, Cowhage is used as a mechanical anthelmintic against thread and round worms. Its use is to be followed by castor oil.

PHYSOSTIGMATIS FABÆ, Calabar Beans, *Eseré Nut*, *Ordeal Bean*.—The seeds of *Physostigma venenosum*, from Western Africa.

Botanical Source.—The plant resembles the common “scarlet runner,” but grows much higher and has a woody stem.

Description.—Oblong seeds about $1-1\frac{1}{2}$ inch long, and about $\frac{3}{4}$ inch broad, with a hard and rough but slightly polished testa of a dark chocolate-brown colour. At one edge they are nearly straight, but at the other

they are arched. Running along the arched side is a deep furrow. At one end of this furrow is seen the *microphyle* as a slight V-shaped depression, and the *raphê* is also observed running up the middle of the furrow as a slightly raised line.

Uses.—Diminishes the action of the heart, and also induces paralysis through its action on the spinal cord. Locally applied it powerfully *contracts* the pupil of the eye, and is chiefly used by oculists in presbyopia. It has been recommended in tetanus, delirium tremens, and epilepsy, but it is a dangerously powerful drug. Dose of the powder, 1 to 4 grains. The best antidote is the hypodermic injection of 1-50th of a grain of sulphate of atropia (repeated, if necessary, at the end of two hours) and artificial respiration.

Chemistry.—The active principle is an alkaloid called *Physostigma*, $C_{30}H_{21}N_3O_4$, which exists chiefly in the kernel. It is soluble in ether, chloroform, alcohol, benzol, acids, and very sparingly in water. Its solutions become red or sometimes intensely blue on exposure to the air, a partial decomposition of the alkaloid taking place. It also becomes red on heating, *per se*, to 212° F. The kernels yield to boiling ether .5 per cent. of fatty

oil. They also contain 23 per cent. of legumin, 48 per cent. of starch, some mucilage, and a little sugar.

Extractum Physostigmatis.

Calabar Bean, in coarse powder	16 ounces
Rectified Spirit.....	4 pints

100 lbs. of beans yield about $2\frac{1}{2}$ lbs. of extract.

Dose—1-16th to 1-4th grain.

KINO, Kino, Gum Kino.—The inspissated juice obtained from incisions made in the trunk of *Pterocarpus Marsupium*. Imported from Malabar.

Botanical Source.—The *P. Marsupium* is a tall tree.

Collection.—The juice is caused to exude through a long incision, and is collected in vessels at the foot of the tree. After a few hours' exposure to the sun and air it hardens, and is then packed in boxes.

Description.—In small, hard, brittle, reddish black fragments, glistening and translucent at the edges. It dissolves in the mouth with a very astringent taste and colours the saliva blood-red. It is entirely soluble in boiling water, but partially separates on cooling, leaving a red solution. It is almost entirely soluble in rectified spirit, but quite insoluble in ether.

Uses.—Astringent, but not so strong in that respect as catechu. Used in diarrhœa.

Chemistry.—Dilute acids cause in solutions a precipitate of *Kino-tannic Acid*, to which the astringency of Kino is due. An excellent test for its presence is to shake up a solution with *Ferrum redactum*, when a beautiful violet colour is produced. The colouring matter of the juice is called *Kino red*.

Pulvis Kino Compositus.

Powdered Kino	3 $\frac{3}{4}$ ounces
„ Opium	$\frac{1}{4}$ ounce
„ Cinnamon	1 ounce

Dose—5 to 20 grains (3 $\frac{3}{4}$ in 5).

Tinctura Kino.

Kino, in coarse powder	2 ounces
Rectified Spirit	ad. 20 fluid ounces

Macerate 7 days and filter.

Dose— $\frac{1}{2}$ to 2 drachms (5 $\frac{1}{2}$ grains of Kino in 1 fluid drachm).

Also an ingredient in Pulvis Catechu Compositus, 1 in 5.

Substitutions.—1. Gambia Kino from *Pterocarpus erinaceus* from Western Africa.

2. Butea gum, or Bengal Kino, from *Butea frondosa*. Used in India in preference to ordinary Kino, and containing 74 per cent. of tannic acid.

3. Botany Bay Kino, the produce of several species of *Eucalyptus* from Australia. This is often imported in a semi-fluid state.

PTEROCARPI LIGNUM, Red Santal Wood, Red Sanders Wood.—The wood of *Pterocarpus santalinus*, imported from Ceylon.

Botanical Source.—The *P. santalinus* is a small tree, a native of Southern India.

Description.—In heavy dark red billets, or in chips having the darker colour in streaks.

Uses.—Feebly astringent, but is only employed in pharmacy as a colouring for Compound Tincture of Lavender. It has numerous uses in the arts.

Chemistry.—The red colouring matter appears to be a resinoid substance, and is termed santalic acid or *santalin*, $C_{15}H_{14}O_5$. It is insoluble in water, turpentine and fixed oils, but soluble in alcohol, ether and acetic acid. Santalic acid will neutralise alkalies, and form with them uncrystallisable salts.

BALSAMUM TOLUTANUM,
Balsam of Tolu.—A balsam obtained by exudation after incision into the bark of *Myroxylon Toluifera*. Imported from New Granada.

Botanical Source.—The *M. Toluifera* is a lofty evergreen tree, sometimes rising to a height of 40 feet before branching.

Collection.—The balsam is obtained by cutting several V-shaped notches into the bark, at the base of which the wood is a little hollowed out, so as to allow calabashes of the size and shape of tea-cups to be fixed. Sometimes as many as twenty calabashes may be seen on various parts of the same trunk, and the *bleeding* is allowed to go on for eight months in the year; so that ultimately the trees become much exhausted and thin in foliage.

Uses.—Stimulant and expectorant. Used in asthma and chronic diseases of the mucous membranes.

Dose, 10 to 20 grains.

Chemistry.—The balsam yields an amorphous resin, apparently identical with that of Balsam of Peru. It also contains a large portion of cinnamic acid, $\text{HC}_9\text{H}_7\text{O}_2$, together with 1 per cent. of tolenc $\text{C}_{10}\text{H}_{16}$, a volatile oil, obtained by distilling the balsam with water.

By destructive distillation, it yields the same compounds as Balsam of Peru, under similar treatment, and also a little phenol and styrol.

The presence of cinnamic acid may be proved by heating with potassium dichromate and sulphuric acid, when *benzoic aldehyd* (oil of bitter almonds) will be given off.

Specially Distinctive Tests.—1. Completely soluble in chloroform and alchohol, but only imperfectly in ether.

2. Exhausted with petroleum spirit and the solution evaporated, a residue is obtained which is not distinctly coloured by sulphuric acid.

3. Yields much benzoic aldehyd when heated with potassium dichromate and sulphuric acid.

Syrupus Tolutanus.

Balsam of Tolu	1½ ounce
Refined Sugar	32 ounces
Distilled Water.....q.s. or	20 fluid ounces
Product, 48 ounces by weight, or 36 fluid ounces by measure, sp. gr. 1·330.	
Dose—I drachm.	

Tinctura Tolutana.

Balsam of Tolu	2½ ounces
Rectified Spirit	ad. 20 fluid ounces
Dissolve and filter.	
Dose—20 to 40 minims (1 grain of Balsam of Tolu in 9 minims.)	
This tincture is used to flavour the Trochisci of Tannin, Morphia, Morphia and Ipecacuanha, and of Opium.	

It is also an ingredient in Tinctura Benzoini Composita.
1 in 40.

BALSAMUM PERUVIANUM,
Balsam of Peru.—A balsam exuding from the trunk of *Myroxylon Pereiræ* after the bark has been scorched and removed. Imported from the western part of the state of San Salvador, known as the *Balsam Coast*, in Central America,

Botanical Source.—The *M. Pereiræ* is a lofty tree, of an exceedingly handsome character.

Collection.—The bark is well bruised with the back of an axe, and it is then charred by means of torches. After the lapse of several days the charred bark falls off and linen rags are applied to the stem, which soak up the exuding balsam. When fully saturated, the rags are thrown into a boiler with water, when the balsam sinks to the bottom; the water is then decanted and the balsam poured into *tecomates* or gourds, ready for the market. It formerly came to Europe through Peru, and so obtained its name.

Description.—A reddish black liquid, sp. gr. 1·15, as thick as syrup, and having a balsamic odour. It is soluble in 5 parts of rectified spirit, and should not alter in volume when treated with water.

Uses.—Stimulant and expectorant. Dose 10 to 15 minims in emulsion with acacia or on sugar. Used also in the form of ointment with castor oil, as a dressing for bed sores and indolent ulcers.

Chemistry.—Contains about 60 per cent. of a body called *Cinnamein*, and 38 per cent. of a black amorphous resin. These two bodies may be separated by mixing some balsam with thrice its weight of carbon disulphide, when the former dissolves, while the latter precipitates. The carbon solution leaves, on evaporation, a brown thick liquid, volatile at 570° F., and having a sp. gr. of 1.1. This is the so-called *cinnamein*, which is really *benzyl cinnamate*, $C_7H_7C_9H_7O_2$, as may be proved by the fact of its conversion into benzoic and cinnamic acids by the action of caustic alkalies.

Specially Distinctive Tests.—Behaves exactly like Tolu balsam, except that the residue in test 2 is coloured yellow-brown, passing to violet by sulphuric acid.

SUB-ORDER II.—CÆSALPINIÆ.

HÆMATOXYLI LIGNUM, Logwood.

—The sliced heart-wood of *Hæmatoxylon Campechianum*. Imported from Campeachy, Honduras and Jamaica.

Botanical Source.—The *H. Campechianum* is a spreading tree, indigenous to Central America, and naturalised in the West Indies.

Collection.—Trees about 10 years old are chosen for felling.

Description.—Usually in chips, of a reddish brown colour and greenish lustre, with a feeble odour and sweetish taste, colouring the saliva pink.

Uses.—Mild astringent, and useful in diarrhœa of young persons, in hæmorrhages, and as an injection in leucorrhœa; but its use requires caution. Its employment in the art of dyeing is far more important.

Chemistry.—The useful agent in logwood is the colouring matter called *Hæmatoxylin*, of which it yields 9 to 12 per cent. The formula of this body is $C_{16}H_{14}O_6$. It is colourless when pure, and soluble in hot water and alcohol, but sparingly in cold water and ether. Its solution becomes red on exposure to light, and violet on adding even a trace of alkali.

This change is due to the formation of *Hæmatein*, $CH_{16}H_{12}O_6 \cdot 3H_2O$, and the colour may be destroyed by various reducing agents, like sulphurous acid, which

re-form the original Hæmatoxylin. A further test for the presence of Hæmatoxylin is found in the fact that it forms pyrogallic acid when fused with potassium hydrate. It may also be precipitated by alum, plumbic acetate, ferric salts and gelatin, and it reduces Fehling's solution of copper.

Decoctum Hæmatoxyli.

Cinnamon, in coarse powder.....	60 grains
Logwood, in chips	1 ounce
Distilled Water	1 pint

Boil 10 minutes, add cinnamon towards the end; make up the product to 20 fluid ounces.

Incompatibles.—Acids; lime water; and most metallic salts.

DOSE—1 to 2 ounces.

Extractum Hæmatoxyli.

Logwood, in fine chips	16 ounces
Boiling Distilled Water	1 gallon

(100 lbs. of wood yield about 18 lbs. of Extract.)

DOSE—10 to 30 grains.

Adulteration.—Chips of Brazil wood (from various species of *Cæsalpinia*), a decoction of which turns *red* instead of *violet*, with alkalies. Oxidised by nitric acid. pure logwood solution will only yield oxalic acid. but if *Brasilin* be present, picric acid will also be formed.

SENNA ALEXANDRINA, Alexandrian Senna.—The leaflets of *Cassia acutifolia*, (*C. lanceolata* and *C. obovata* B.P.) carefully freed from the leaf-stalks, flowers and pods of the same. Imported from Alexandria.

Botanical Source.—The *C. acutifolia* is a shrub about 2 feet high, with paripinnate leaves and axillary racemes of yellow flowers.

Collection.—The plants are cut down in September, and sometimes also in April, and laid on rocks in the sun till dry. The leaflets are then carefully stripped off and packed in bags made from palm leaves.

Description.—Ovate or lanceolate, acute and mucronate (with *mucro* a sharp point or edge) leaflets, about $\frac{3}{4}$ to $1\frac{1}{4}$ inch long, and unequal at the base. They are conspicuously veined, and have often an incurved margin and a brown midrib. They are very finely pubescent, especially close to the midrib, and are greyish green in colour.

Uses.—Purgative, producing watery stools by acting on the small intestines. It is by some considered rather heating, and is apt to gripe and nauseate.

Dose, 30 to 120 grains.

Chemistry.—The chief principle is an uncrystallisable glucoside called *Cathartic acid*, $C_{180}H_{192}O_{82}N_4S$, which may be obtained by mixing a syrupy aqueous extract of senna with an equal volume of alcohol, and filtering out the mucilage thus thrown down, when the addition of a further quantity of alcohol throws down a dark substance, from which when treated with hydrochloric acid Cathartic acid is precipitated. It is almost insoluble in water and alcohol, but entirely soluble in ether and chloroform, and it dissolves in warm dilute alcohol. It is precipitated by most acids, but not by tannic. Cathartic acid when boiled with hydrochloric acid is resolved into glucose and *Cathartogenic acid*. Senna contains *Chrysophanic acid* and some tartaric and oxalic acids, with traces of malic acid, also a peculiar crystalline sugar, called *catharto-mannite*, which is insusceptible of fermentation.

Confectio Sennæ (*Lenitive Electuary*).

Senna, in fine powder.....	7 ounces
Coriander Fruit, in fine powder	3 ounces
Figs	12 ounces
Tamarinds	9 ounces
Cassia Pulp	9 ounces
Prunes	6 ounces
Extract of Liquorice	3 ounce
Refined Sugar	30 ounces
Distilled Water	24 fluid ounces

Product 75 ounces.

Dose—1 to 2 drachms ($6\frac{1}{2}$ grains of Senna in 1 drachm).

Infusum Sennæ.

Senna.....	1 ounce
Ginger, sliced	30 grains
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Incompatibles.—Strong acids; lime water; and most metallic salts.

Dose—I to 2 ounces.

Mistura Sennæ Composita.

Sulphate of Magnesia.....	4 ounces
Extract of Liquorice	$\frac{1}{2}$ ounce
Tincture of Senna	$2\frac{1}{2}$ fluid ounces
Compound Tinct. of Cardamoms	10 fluid ounces
Infusion of Senna	20 fluid ounces
Dose—I to $1\frac{1}{2}$ ounce (about 1 drachm of magnes. sulph. in 5 fluid drachms).	

Syrupus Sennæ.

Senna, broken small	16 ounces
Oil of Coriander	3 minims
Refined Sugar	24 ounces
Distilled Water.....q.s. or	5 pints
Rectified Spirit.....	2 fluid ounces

Product, 42 ounces by weight, or 32 fluid ounces by measure; sp. gr. 1.310.

Dose—I to 4 drachms.

Tinctura Sennæ.

Senna, broken small	$2\frac{1}{2}$ ounces
Raisins, freed from seeds	2 ounces
Caraway Fruit, bruised	$\frac{1}{2}$ ounce
Coriander Fruit, bruised.....	$\frac{1}{2}$ ounce
Proof Spirit	20 fluid ounces

Macerate 48 hours, then percolate.

Dose—I to 4 drachms (28 grains of senna, in $\frac{1}{2}$ fluid ounce).

Adulterations.—1. The leaves of *Cynanchum*, or *Solenostemma Argel* (Nat. Ord. Asclepiadaceæ), which are thicker than those of senna, equal at the base, and not distinctly veined. The small corymbs of white star-like flowers, and the slender pear-shaped follicles of the same plant may also be found.

2. The leaves of *Coriaria myrtifolia*, which are ovate, acute, and unequal at the base, but have a *distinct vein running up just inside each margin*.

3. The leaves of *Tephrosia Apollinea*, which are obovate and not at all mucronate.

SENNA INDICA, Tinnevelly Senna.
—The leaves of *Cassia angustifolia* (*C. elongata* B.P.), from plants cultivated in Southern India.

Description.—Yellowish green leaflets, about 1 to 2 inches long, with an unequal base, and almost glabrous.

Chemistry and Uses.—The same as Alexandrian Senna, to which it is to be preferred, owing to its freedom from adulteration. It is said however not to be quite so active.

BOMBAY or MECCA SENNA, East Indian Senna.—The produce of *wild* plants of

C. angustifolia, collected in Arabia, and exported to Bombay and from thence to Europe.

Description.—Of the most inferior description, consisting of brown and broken leaflets with many stalks, flowers, and pods.

CASSIÆ FISTULÆ FRUCTUS,
Purging Cassia.—The indehiscent lomentum of *Cassia Fistula*, imported from the East and West Indies.

Botanical Source.—This plant is a tree growing 30 feet high, with pendulous racemes of yellow flowers, originally indigenous in the East Indies, but cultivated in the West Indies and in Brazil.

Description.—Long narrow cylindrical pods about $\frac{3}{4}$ to 1 inch in diameter, having the ventral and dorsal sutures distinctly marked, and when curved with the ventral suture inside. The interior of the pod is divided into a number of small divisions, by means of *spurious dissepiments*, and in each division is an oval, flat reddish brown seed with a shiny testa and a well-marked *raphé* as a dark line on one side. The seeds are embedded in a black sweetish pulp usually partially dried up in the imported pod.

Chemistry.—The pulp contains sugar, gum, pectin, and a cathartic principle.

Cassia Pulpa (Cassia Pulp).—The pulp of the fruit separated from the pods. It is an ingredient in *Confectio Sennæ* (1 in 8½ nearly).

CASSIÆ GRANDIS FRUCTUS,
Horse Cassia.—The lomentum of *C. grandis* or *C. Brasilina*, from Central America. The pulp of this fruit is much used in veterinary practice. It is a very much larger fruit, and the pulp is bitter and astringent.

TAMARINDUS, Tamarind. — The preserved pulp of the ripe fruit of *Tamarindus Indica*, chiefly imported from the West Indies.

Botanical Source.—A large tree with compound pinnate leaves, and having indehiscent legumes.

Description.—The fruit is an oblong legume from 3 to 6 inches long, and with 4 to 12 seeds. It consists of a hard and brittle epicarp, a pulpy mesocarp, and a tough membranous endocarp surrounding each seed like a cell. On the surface of the pulp run very characteristic strong fibres.

Collection.—The fruit is plucked when fully ripe from June till August, and the outer shell having been broken and stripped off, the pulp and seeds are placed in layers in a cask with a sprinkling of raw sugar between. When the cask is sufficiently packed, boiling syrup is poured over, and when cold the whole is closed up for export. This is the process for preserving the *brown* or *West Indian tamarinds* which are used in this country; but the *black* or *East Indian tamarinds*, more generally employed on the Continent, are packed without sugar.

Uses.—Mildly laxative; used for making a refrigerant drink in fevers. The cotyledons of the seeds, which have a bland mucilaginous taste, serve in India as a food during times of scarcity.

Chemistry.—Unsweetened tamarind pulp contains 34·35 parts of large-celled parenchyma, 27·55 per cent. of water, 9·4 per cent. of citric acid, 1·55 per cent. of tartaric acid, 0·45 per cent. of malic acid, 3·25 per cent. of acid tartrate of potash, 4·7 per cent. of gum, 6·25 per cent. of pectin, and 12·5 per cent. of sugar; but as sugar is added to preserve the West Indian pulp, it will be found to contain twice as much sugar, and proportionately less

of the other ingredients. The testa of the seed contains a large quantity of tannin. A piece of bright iron left in contact with the pulp for about an hour should not exhibit any deposit of copper, with which the tamarinds are sometimes contaminated through being placed in copper vessels.

Preparation.—An ingredient in *Confectio Sennæ*.

COPAIBA, Copaiva.—The oleo-resin obtained from incisions made in the trunk of *Copaifera multijuga* and other species of *Copaifera*, chiefly from the valley of the Amazon. Its true botanical origin is doubtful, and its orthography is not definite.

Collection.—A large wedge-shaped notch is cut right into the heart of the tree near its base, and from this the balsam flows and is collected.

Description.—An oily thick liquid varying in colour from deep brownish to pale amber, and in sp. gr. from '94 to '99. It has a characteristic odour, and should not become gelatinous after heating to 270° F. Copaiva should be soluble in its own weight of benzol.

Uses.—Acts specially on the mucous membranes of the urinary organs and the rectum. Employed as a

specific in gonorrhœa and also in hæmorrhoids. The dose of the oleo-resin is from $\frac{1}{2}$ to 1 drachm. As usually administered the balsam is enclosed in gelatin capsules, but it is also made into an emulsion with mucilage or alkalies, or into pills by means of an equal weight of magnesium carbonate.

The dose of the oil is from 5 to 20 minims.

Chemistry.—40 to 60 per cent. of a volatile oil, $C_{10}H_{16}$, isomeric with turpentine, holding in solution an acid resin called *copaivic acid*, $C_{20}H_{32}O_2$. By distilling the copaiba with water the oil passes over and the resin remains as a brittle mass.

Specially Distinctive Tests.—All varieties of this substance are—1. Completely soluble in chloroform and entirely soluble in ether, the latter solution becoming turbid on the addition of alcohol.

2. An alcoholic solution gives no turbidity with an alcoholic solution of ferric chloride (1 in 10).

3. Entirely soluble in petroleum spirit boiling below 100° F.

After applying these general tests, Maranham and Para balsam may be distinguished from each other by dissolving in chloroform and adding a solution of bromine

in chloroform (1 in 20), which colours the former bluish but does not affect the latter.

Varieties and Adulteration.—The two chief varieties are *Maranham* and *Para*, the latter being usually a very fluid and light-coloured balsam. An inferior dark balsam is imported from Maracaibo. The most common adulteration is castor and other fixed oils. These may be most certainly detected and estimated by the process introduced by the present author. It consists in saponifying balsam with alcoholic soda, adding water and evaporating until the alcohol is expelled, then *exactly neutralising* the solution by a dilute acid and evaporating to dryness. The dry neutral soap is then treated with a mixture of 1 part of absolute alcohol and 5 parts of absolute ether, in which the copaiba soap dissolves easily, leaving any oil soap insoluble. The latter (if any) is then dissolved in water and boiled with excess of hydrochloric acid, which causes the fatty acids to separate and float on the surface, when they may be collected and weighed.*

A more rough and rapid way consists in simply boiling the copaiba with water until the volatile oil is

* For full particulars see "THE ANALYST," vol. i., p. 160.

expelled, and then letting the resin cool, when, if any fixed oil be present, it will make the resin soft or even remain floating in oily drops.

Another method is to heat the copaiba with 4 parts of rectified spirit and letting it cool, when it separates into two layers, and the upper will contain any castor oil which may be recovered by evaporation, and tested for and heating with soda lime (see Castor Oil).

The non-coagulation after heating to 270° F. proves the absence of *Gurjun Balsam*, for which see p. 71.

COPAIBÆ OLEUM.—Obtained by distilling *Copaiba* with water. A volatile oil, sp. gr. .88 to .9, colourless or pale yellow. Miscible with ether, and soluble in 8 parts of rectified spirit. It boils at 475° F., and is a solvent of sulphur, phosphorus, and iodine.

SUB-ORDER III.—MIMOSÆ.

GUMMI ACACIÆ, Gum Arabic.—A gummy exudation from the stems of one or more species of *Acacia*. Imported from Alexandria and elsewhere.

Botanical Source.—The plant which produces the

finest or *Kordofan* gum, is the *Acacia Verek*, a small tree growing in Eastern Africa, while the inferior sorts are produced by other species.

Varieties.—1. *Kordofan*, or “picked Turkey Gum.”—This is the finest colourless gum, in lumps of varying sizes and full of minute cracks. Sp. gr. 1.487, and if dried at 212° F. 1.525.

2. *Senegal Gum*.—Is in yellowish lumps, not having the minute cracks of the fine acacia. Imported from the French settlements in Senegal.

3. *Suakin or Talca Gum*.—Imported from Alexandria in very small fragments of colourless gum, mixed with many pieces of a deep red-brown colour. Usually known in the trade as *Acacia “sorts.”*

Besides these there are inferior brown gums imported from Morocco, Bombay, Australia (*Wattle gum*), and the Cape of Good Hope.

Uses.—Topically emollient and demulcent. Employed to suspend heavy powders, and to emulsify resins.

Chemistry.—Gum Arabic forms with 2 parts of water a mucilage having a sp. gr. of 1.149. It is slightly soluble in dilute spirit, but quite insoluble in spirit, containing

an amount of alcohol over 60 per cent. Its aqueous solution is not precipitated by normal plumbic acetate, but with *basic* plumbic acetate it forms a white jelly. When boiled with diluted sulphuric acid it is converted first into dextrin and then into sugar; while nitric acid transforms it into mucic and oxalic acids. It may be regarded as a compound of *Arabic acid* or *Arabin*, $C_{12}H_{22}O_{11}$, with calcium and potassium, and the formula $CaK_2C_{12}H_{22}O_{11} \cdot 6C_{12}H_{22}O_{11}$ has been assigned to it. The presence of calcium may be proved by adding ammonium oxalate to a solution, and *Arabic acid* may be isolated by the dialysis of a mucilage acidulated with hydrochloric acid, when the calcium and potassium will pass through as chlorides, leaving in the dialyser the *Arabic acid* which, when dried, is insoluble in water, but swells in it. Solutions of gum acacia are precipitated by borax and ferric salts.

Mucilago Acaciæ.

Gum Acacia, in small pieces.....	4 ounces
Distilled Water	6 fluid ounces

(Measures $8\frac{3}{4}$ ounces).

(3 drachms are required to emulsify 1 ounce of oil or resinous tinctures, and 8 drachms for 1 ounce of Copaiba.)

Incompatibles.—Rectified spirit; ferric salts; acetate of lead, and mercuric nitrate.

Gum Arabic also enters into *Mistura Cretæ* (1 in 34); *Mistura Guaiaci* (1 in 85); *Pulvis Amygdalæ Compositus* (1 in 13); *Pulvis Tragacanthæ Compositus* (1 in 13); in all the *Trochisci*, and in the French preparation *Sirop de Gomme*.

CATECHU NIGRUM, Black Catechu, *Cutch*.—An aqueous extract of the heart wood of *Acacia Catechu*. Imported from Pegu.

Botanical Source.—The article of commerce is obtained from two trees, the *A. Catechu* and the *A. Suma*, both growing in Southern India and in the tropical parts of Eastern Africa.

Collection.—The wood is cut into chips and boiled with water in earthen pots. The liquor is evaporated to a semi-solid state, and then run into clay moulds to harden.

Description.—In blackish brown masses, having an irregular shining but somewhat granular fracture. Frequently having fragments of brown leaves attached to the outside.

Uses.—Similar to pale Catechu, but much more astringent. (See *Cinchonaceæ*.)

Chemistry.—Cutch is only slightly soluble in cold water, but dissolves on boiling and crystallises out again on cooling. The portion soluble in cold water is *Catechu-tannic acid*, having the properties of ordinary tannic acid; and the portion insoluble is *Catechin* or *Catechuic acid*, $C_{13}H_{12}O_5$, which may be extracted from Cutch by ether.

NATURAL ORDER.—ROSACEÆ.

Trees, shrubs or herbs with alternate and usually stipulate leaves and regular flowers; calyx monosepalous and persistent; corolla rosaceous; stamens numerous and perigynous, with 2-celled innate anthers; pistil usually apocarpous and superior, sometimes becoming inferior in the fruit; fruit either a drupe, achæmium, follicle, etærio, cynarrhodum or pome; seeds exalbuminous.

SUB-ORDER I.—CHRYSOBALANEÆ.

Trees or shrubs with free stipules; style basilar; fruit a drupe.

SUB-ORDER II.—AMYGDALÆ.

Trees or shrubs, calyx deciduous; style terminal; fruit a drupe.

AMYGDALÆ DULCES, Sweet Almonds.—The seeds of *Amygdalus communis*, var. *dulcis*. Cultivated about Malaga.

Botanical Source.—The almond is a small tree with nearly sessile solitary flowers appearing before the leaves. It bears a drupe with a velvety mesocarp, which when ripe dries and splits off, leaving an oblong endocarp pitted with numerous holes, and containing the seeds, which are the almonds of commerce.

Varieties.—The finest are the Jordan almonds, from Malaga, easily known by their size and oblong form, and usually imported without the endocarp. The other qualities, in the order of their value, are those from Valencia, Sicily and Barbary, generally imported in their shells.

Uses.—To make Oleum Amygdalæ, and to suspend heavy powders in emulsion.

Chemistry.—Sweet almonds yield a fixed oil to the extent of 50 per cent. (see Ol. Amygd.) and also contain a nitrogenous body called *emulsin*, together with sugar and mucilage.

Mistura Amygdalæ.

Compound Powder of Almonds	2½ ounces
Distilled Water	20 fluid ounces

DOSE—1 to 2 ounces (54½ grains of powder in 1 fluid ounce).

Pulvis Amygdalæ Compositus.

Sweet Almonds	8 ounces
Refined Sugar	4 ounces
Gum Acacia	1 ounce

AMYGDALÆ AMARÆ, Bitter Almonds.—The seeds of *Amygdalis communis*, var. *amara*, imported from Mogadore.


Botanical Source.—Already described.

Description.—Smaller than sweet almonds, bitter to the taste, and giving off when rubbed with water an odour of hydrocyanic acid. The best are the French; the Sicilian and Barbary follow in order of value.

Uses.—Employed as a flavouring, and often with very deleterious effects.

Chemistry.—They yield by pressure 44 per cent. of

fixed oil, and the residual cake when boiled with alcohol yields a solution containing *Amygdalin*, which crystallises and gives the formula $C_{20}H_{27}NO_{11} \cdot 2H_2O$. It is a neutral bitter inodorous glucoside, and rarely amounts to more than $2\frac{1}{2}$ per cent. The residue from the alcohol if digested with water yields an albuminoid body called *emulsin*, which is precipitated by alcohol and rendered inert by boiling. It possesses the power of decomposing the amygdalin in presence of water, forming benzoic aldehyd (oil of bitter almonds), hydrocyanic acid, and glucose.



OLEUM AMYGDALÆ AMARÆ.—
Essential Oil of Bitter Almonds (*Benzoic aldehyd*) C_7H_6O , is prepared by making the residual bitter almond cake, after expression of the fixed oil, into an emulsion with cold water, letting it stand for 24 hours (for the emulsin to act on the amygdalin and develop the essential oil) and then distilling. The crude oil may be freed from hydrocyanic acid by agitating with a strong solution of sodium hydrosulphite, $NaHSO_3$, which combines with

the benzoic aldehyd and forms a crystalline precipitate. On separating this and treating with sodium carbonate, the pure oil is obtained and is then freed from moisture by distillation over calcium chloride.

Another process is to agitate the crude oil with milk of lime and ferrous sulphate, and then distil. The purified oil is more liable to oxidation than the crude oil, and much more expensive. Bitter almond cake yields about 1 per cent. of the crude oil. Pure benzoic aldehyd (often expressed in books as benzoyl hydride, $C_7H_5O.H$) is colourless, with a sp. gr. of 1.049, and boils at $356^{\circ} F$. Exposed to the air it oxidises into a crystalline mass of benzoic acid, $HC_7H_5O_2$. The crude oil is yellow, soluble in alcohol and ether, and forms a red liquid with sulphuric acid, which when poured into water produces a yellow emulsion.

Tests.—All samples should be carefully tested for hydrocyanic acid, as fresh oil may contain from 8 to 14 per cent., although it becomes lessened in quantity by age. If present in good quantity it may be detected by the ordinary *iron* test, and estimated if necessary by shaking up the oil with water, then adding a slight excess of sodium hydrate, and running in volumetric solution of silver.

A sure and delicate test paper for the examination of substances suspected to contain hydrocyanic acid is made by dipping bibulous paper in fresh tincture of guaiacum, then drying and dipping in an aqueous solution of cupric sulphate (1 in 2000), and once more drying. The paper will become intensely blue if moistened and afterwards dipped into oil containing hydrocyanic acid.

AMYGDALÆ OLEUM, Almond Oil.

—An oil obtained by expression from either bitter or sweet almonds.

Description.—A thin pale yellow oil, sp. gr. .92, and not coagulable at 32° F. like olive oil, but remaining fluid until cooled to 12.5° F. It gets rancid in contact with the air, but does not dry.

Uses.—An ingredient in Unguentum Cetacei, Unguentum Hydrargyri Oxidi Rubri, Unguentum Plumbi Subacetatis Compositum, Unguentum Simplex, and all other ointments containing the latter as an ingredient. With lard and wax it produces much whiter ointments than *Oleum Olivæ*.

Chemistry.—Almond oil is nearly all pure *olein*,
 $C_3H_5C_{18}H_{33}O_2$.

PRUNI FRUCTUS, Prunes.—The dried fruit of *Prunus domestica*. Imported from France.

Botanical Source.—The *P. domestica* is a small tree, destitute of spines, and having usually ovate lanceolate leaves. The flowers are in axillary cymes, having a white rosaceous corolla, numerous perigynous stamens, and a superior simple ovary producing a drupe well known as the common black plum. The dried drupe is imported from Southern Europe.

Varieties.—The best prunes are those from Bordeaux, which are dried by alternating the heat of a stove with that of the sun.

Uses.—Nutritious, demulcent and mildly laxative, especially for children.

Chemistry.—The same as most of the ordinary edible fruits, containing sugar, pectin, gummy matter, malic acid and some calcium oxalate.

An ingredient in *Confectio Sennæ*.

PRUNI VIRGINIANÆ CORTEX,
Wild Black Cherry bark.—The bark of *Prunus serotina*. Imported from North America.

Uses.—Of late this bark has been frequently prescribed in England; when obtained from the last-named plant it is official in the United States Pharmacopœia as “Cortex pruni serotinæ,” being used as a sedative and tonic, and in heart disease.

Chemistry.—Emulsin and amygdalin, which when brought into contact with water yield Benzoic Aldehyd and Hydrocyanic Acid. The bark also contains tannic acid and bitter extractive.

LAUROCERASI FOLIA, Cherry Laurel Leaves.—The fresh leaves of *Prunus Lauro-cerasus*. A native of Asia Minor, but now cultivated in Britain.

Botanical Source.—This plant is an evergreen shrub, having oblong smooth coriaceous shining leaves, with slightly serrate margin and shortly petiolate. Inflorescence axillary racemes. Corolla white rosaceous and spreading; stamens indefinite, perigynous; fruit a black drupe, resembling a cherry.

Uses.—Sedative and narcotic. Employed in heart disease, but its use is generally superseded by that of the more definite Hydrocyanic Acid.

Chemistry.—The leaves yield by distillation a volatile oil, similar to that of bitter almonds, *i.e.*, Benzoic Aldehyd, containing Hydrocyanic Acid. This is probably developed in the leaves, by the action of bodies similar to Emulsin and Amygdalin, although the latter has not been obtained from them.

Preparation.—*Aqua Laurocerasi*, 1 lb. to $2\frac{1}{2}$ pints, distil 1 pint. Dose—5 to 30 minims.

SUB-ORDER III.—ROSEÆ.

Shrubs or herbs with compound leaves and adnate stipules, fruit, follicles, etærio, or cynarrhodum.

CUSSO, Flores Koso, Kousso.—The flowers and tops of *Hagenia Abyssinica* (*Brayera anthelmintica*, B.P.). Collected in Abyssinia.

Botanical Source.—This is a tree growing in the highlands of Abyssinia. The inflorescence is in panicles of unisexual flowers, and the plant is monœcious. The female inflorescences, which are chiefly collected, have the calyx 5 partite in two rows, the corolla being very minute. Two carpels with hairy styles are contained in the calyx tube.

Description.—A mass of brownish flowers on panicles 10 inches long, with hairy pedicels, the points of the female flowers already noticed being discernible with care. It has an odour like tea, and a bitter acrid taste.

Uses.—Anthelmintic, especially in tapeworm, but by some thought to be comparatively useless. Dose, $\frac{1}{4}$ to $\frac{1}{2}$ ounce.

Chemistry.—Three per cent. of *koussin* or *kosin*, 24 per cent. of tannic acid, 6.5 per cent. of an acrid resin, gum, sugar, wax, and *koussin*, $C_{31}H_{38}O_{10}$, may be extracted by mixing the flowers with lime, exhausting them first with alcohol, then with water, concentrating the mixed solution and adding acetic acid which precipitates the *koussin*. It is freely soluble in boiling alcohol, in ether, chloroform, and solutions of the caustic alkalies, from which it is precipitated by acids.

Infusum Cusso $\frac{1}{2}$ ounce to 8 fluid ounces. To be taken unstrained.

ROSÆ GALLICÆ PETALA, Rose Leaves.—The fresh and dried unexpanded petals of *Rosa Gallica*, from plants cultivated in Britain.

Botanical Source.—The *R. Gallica* is a dwarfish

shrub covered with prickles and glandular hairs intermixed. Leaves compound imparipinnate, inflorescence axillary in a cluster, sepals ovate, leafy, compound; cynarrhodum oblong.

Uses.—Very slightly astringent and tonic, but chiefly used as a colouring matter. The acid infusion is given as a refrigerant drink in fevers, and as a vehicle for saline purgatives and gargles. The Confection is the excipient for the Pills of Aloes, Barbadoes Aloes and Assafoetida, Aloes and Iron, Aloes and Myrrh, Aloes Socotrina, Carbonate of Iron, Mercury, and Lead with Opium.

Chemistry.—A slight trace of volatile oil, tannic and gallic acids, and a glucoside called *quercitrin* ($C_{33}H_{30}O_{17}H_2O$), a little fat, glucose and colouring matter. *Quercitrin*, when boiled with a dilute acid, yields glucose and *quercetin* ($C_{27}H_{18}O_{12}$).

The colouring matter is turned bright red by mineral or vegetable acids; green by alkalies. It is entirely bleached by strong sulphuric acid.

Confectio Rosæ Gallicæ.

Fresh Red Rose Petals	16 ounces
Refined Sugar	48 ounces

Infusum Rosæ Acidum.

Dried Red Rose Petals, broken up	$\frac{1}{2}$ ounce
Dilute Sulphuric Acid.....	60 minims
Boiling Distilled Water	10 fluid ounces

Infuse $\frac{1}{2}$ hour and strain.

Incompatibles—Alkalies; earths; and all substances which act on, or combine with sulphuric acid. Ferri sulph. gives it a brown colour.

DOSE—I to 2 ounces.

Syrupus Rosæ Gallicæ.

Dried Red Rose Petals	2 ounces
Refined Sugar	30 ounces
Boiling Distilled Water	20 fluid ounces

Product, 46 ounces by weight, or 35 fluid ounces by measure; sp. gr. 1.335.

DOSE—I drachm.

ROSÆ CENTIFOLIÆ PETALA,
Cabbage Rose Leaves.—The fresh petals of *Rosa centifolia*, from plants cultivated in Britain.

Botanical Source.—A shrub closely covered with prickles and glandular hairs. Leaves compound imparipinnate, having ovate leaflets stipulate and hairy on the under surface. Inflorescence axillary in a drooping cluster; calyx divided in a pinnate manner but not reflexed; cynarrhodum ovate.

Chemistry.—The odour is due to the volatile oil which exists in the petals to about 0·04 per cent., and is the costly Attar or Otto of Roses. The petals also contain tannic, malic, and tartaric acids ; resin, fat and sugar.

Aqua Rosæ, 10 lbs. to 2 gallons ; distil 1 gallon.

Rose water enter into the composition of *Mistura Ferri Composita* and *Trochisci Bismuthi*.

OLEUM ROSÆ, *Attar or Otto of Roses*.—The chief supply of *Otto of Roses* comes from Turkey. It is distilled in copper stills from the fresh flowers. The first portion of the distillate is returned to the still ; that afterwards obtained is set aside for a few hours in order that the oil may separate, when it is skimmed off by means of a small tin funnel. Turkish attar consists of about 18 per cent. of odourless *stearopten*, $C_{20}H_{40}$, and of an oxygenated hydro-carbon, which is the odoriferous constituent. It should be of a bright straw colour, having a sp. gr. of 0·87 to 0·89, and becoming solid at between 60° and 68° F.

Inferior oils distilled from French flowers contain sometimes as much as 60 per cent. of *stearopten*, and congeal at about 86° F.

ROSÆ CANINÆ FRUCTUS, Hips.—

The ripe fruit of *Rosa canina* from indigenous plants.

Botanical Source.—A prickly plant with compound imparipinnate leaves, having adnate stipules. The calyx segments are pinnate and deciduous, the corolla rosaceous, and the fruit is apocarpous consisting of a number of achenes, enclosed in a red succulent tube produced by the union of the calyx tube and the thalamus; the whole forming an ovate cynarrhodum about an inch long.

Chemistry.—The pulp contains 3 per cent. of free citric acid, 7 per cent. of free malic acid, 25 per cent. of gum, 30 per cent. of uncrystallisable sugar, and the calcium and potassium citrates.

Preparation.—Pounded with twice their weight of refined sugar, they form *Confectio Rosæ Caninæ*, which in its turn is used as an excipient in *Pilula Quinæ*. The achenes are removed because the hairs with which they are surrounded are, like cowhage, exceedingly irritating.

SUB-ORDER IV.—SANGUISORBEÆ.

Usually herbs with the corolla absent and the fruit or achæmium enclosed in the calyx tube.

SUB-ORDER V.—POMEÆ.

Trees or shrubs with either simple or compound leaves ; stipules free ; fruit a *pome*.

CYDONIÆ SEMINA, Quince Seeds.—

The seeds of *Pyrus Cydonia*, a native of Western Asia, but cultivated in Southern Europe.

Botanical Source.—The *P. Cydonia* is a tree with ovate and entire leaves, and bearing a golden yellow *pome* resembling a pear, but differing by having 8 to 14 seeds in each cell embedded in a mucilaginous substance.

Description.—Mahogany-brown, ovoid, and three-sided seeds, generally agglutinated in masses.

Uses.—Demulcent, like linseed.

Chemistry.—The epidermis is so mucilaginous that it will absorb forty times its weight of water. The

formula of this mucilage is $C_{12}H_{20}O_{10}$, and it is identical with that of linseed. The entire seed yields 20 per cent. of the dry mucilage.

Decoctum Cydoniæ (not official), 120 grains to 1 pint.

SUB-DIVISION II.—EPIGYNÆ.

NATURAL ORDER.—CUCURBITACEÆ.

Succulent herbs with alternate scabrous leaves and unisexual flowers, with regular monopetalous corollas; male flowers having 5 stamens, frequently united and with sinuous anthers; female flowers having inferior 3-celled ovary with parietal placentation; fruit a *pepo*; seeds numerous and exalbuminous.

The order is divided into 3 sub-orders:—

1. *Nhandirobeæ*, anthers not sinuous.
2. *Siceæ*, only one seed in fruit.
3. *Cucurbiteæ*, anthers sinuous and fruit a many-seeded *pepo*.

ELATERIUM.—The sediment from the juice of the nearly ripe fruit of *Ecballium officinarum* (*Ecballium Elaterium*), cultivated in England and also imported from Malta.

Botanical Source.—The plant is a coarse, decumbent annual, with hispid and scabrous stems, which are without tendrils; leaves cordate with a crenate margin, and also hispid; flowers monœcious, stamens in male flowers triadelphous; ovary in female flowers, 3-celled and inferior. The fruit a *pepo*, which dehisces with elasticity and forcibly projects the juice and seeds from an orifice formed by the separation from the stalk.

Collection.—The juice is lightly pressed from the longitudinally-sliced fruit, strained through a hair sieve, and set aside to deposit. The supernatant liquor having been poured off, the sediment is collected on a linen filter and dried on porous tiles.

Description.—In greenish grey, light and friable fragments, about 1-10th of an inch thick, curved and marked on the convex surface by the impressions of the calico filter. The best is the English, and the inferior is the Maltese, which is in larger pieces with scraps of paper attached, on which it has been dried.

Uses.—Drastic, purgative, and hydragogue. It is the most powerful purgative known, even more so than Croton Oil. It is used in dropsy, in mania, and to

prevent apoplexy; but it requires great caution in its administration.

Its antidotes are, warm fomentations, demulcent drinks and opium, followed by stimulants if the circulation fail.

Dose, from 1-16th to $\frac{1}{2}$ grain.

Chemistry.—The juice of the fruit yields 0.123 per cent. of Elaterium, 3 per cent. of organic and 1.5 per cent. of inorganic matter, and 95 per cent. of water. Elaterium contains starchy and extractive matter and a glucoside called *Elaterin* ($C_{20}H_{28}O_5$), though in the purest samples both starch and cellular tissue appear to be absent. Elaterin may be extracted by treating Elaterium with rectified spirit, which dissolves about half its weight, filtering, and pouring the concentrated solution into warm solution of potassium hydrate, when the Elaterin separates on cooling in colourless crystals. The B.P. says that Elaterium thus treated should yield not less than 20 per cent. of *Elaterin*.

Elaterium should not effervesce with acids, showing absence of chalk.

A more complete method of analysis is to exhaust with chloroform and then add ether, which instantly

precipitates *elaterin*. The crystals may then be washed with ether and dried. By this process 36 and 27 per cent. of *elaterin* has been obtained from Mitcham and Maltese elaterium respectively.

Pulvis Elaterii Compositus.

Elaterium	10 grains
Sugar of Milk	90 grains

Rub them together in a mortar until they are reduced to a fine powder, and intimately mixed.

Dose— $\frac{1}{2}$ to 5 grains.

COLOCYNTHIDIS PULPA, *Colocynthis*

Pulp.—The dried decorticated fruit of *Citrullus Colocynthis* freed from seeds. Imported chiefly from Smyrna, Trieste, France and Spain.

Botanical Source.—The *C. Colocynthis* is an annual herb, somewhat similar in appearance to a cucumber plant, but with the leaves more divided. The pepo is rounded and about the size of a large orange, and has a thin yellow rind.

Description.—The best is the Levant or *peeled* colocynth, in white balls, light and spongy, with the seeds enclosed, which have to be removed before sale. The

second quality is the *unpeeled*, from Mogadore, which is the entire fruit covered with a thin brittle yellow rind.

100 parts of good fruits yield about 30 parts by weight of pulp, and 75 parts of seeds.

Uses.—In small doses an excellent purgative, especially in cases of long standing constipation; but in large doses it is a powerful drastic hydragogue and cathartic. It acts superficially upon the large intestines, and in extreme doses (100 grains) has proved fatal. It is useful in dropsy and uterine obstructions.

Dose, in powder, 2 to 8 grains.

Chemistry.—Colocynth pulp has been found to contain $\frac{1}{4}$ per cent. of Colocynthin, 13.2 per cent. of bitter resin, 14.4 per cent. of bitter principle, 10 per cent. of extractive, 30 per cent. of gummy matters, 5.7 per cent. of phosphates of lime and magnesia, and 19.2 per cent. of lignin.

Colocynthin, $C_{56}H_{84}O_{23}$, may be extracted by treating an alcoholic extract of colocynth with water, and mixing the solution first with neutral acetate of lead and afterwards with basic acetate of lead. From the filtered liquid the lead is separated by means of sulphuretted hydrogen, and then tannic acid is added which causes a precipitate. The precipitate is washed and dried, then

decomposed by lead oxide and the colocynthin dissolved out by means of ether. Colocynthin is thus produced in yellow tufts which, when boiled with a dilute acid, yield *colocynthein*, $C_{44}H_{64}O_{13}$, and glucose.

The seeds contain about 17 per cent. of fixed oil, and are eaten roasted and boiled by some African tribes.

Extractum Colocynthis Compositum.

Colocynth Pulp	6 ounces
Extract of Socotrine Aloes	12 ounces
Resin of Scammony	4 ounces
Hard Soap, in powder	3 ounces
Cardamom Seeds, fine in powder.....	1 ounce
Proof Spirit	1 gallon
(100 lbs. of Pulp yield about 65 lbs. of simple extract.)	

Dose—3 to 10 grains.

Pilula Colocynthis Composita.

Colocynth Pulp, in powder	1 ounce
Barbadoes Aloes, in powder	2 ounces
Scammony, in powder	2 ounces
Sulphate of Potash.....	$\frac{1}{2}$ ounce
Oil of Cloves.....	2 drachms (fluid)
Distilled Water	q.s.

1 grain of Colocynth and 2 grains each of Aloes and Scammony in 6 grains.

Dose—5 to 10 grains. Syrup or glycerin is the best excipient.

Pilula Colocynthis et Hyoscyami.

Compound Colocynth Pill.....	2 ounces
Extract of Henbane.....	1 ounce

Dose—5 to 10 grains.

NATURAL ORDER.—MYRTACEÆ.

CAJUPUTI OLEUM, Oil of Cajuput.

—The oil distilled from the leaves of *Melaleuca minor*, imported from Batavia and Singapore.

Description.—A light bluish green mobile liquid with a powerful odour, having a sp. gr. of $\cdot 914$ to $\cdot 925$, and consisting chiefly of a hydrated *terpene*, $C_{10}H_{16}H_2O$, sometimes called hydrate of cajuputene, with some organic colouring matter. To prevent the oil losing its fine green tint it is kept in copper vessels, which accounts for the traces of copper found in many samples.

Uses.—Cajuput oil is occasionally administered as a stimulant, antispasmodic, and diaphoretic; externally it is frequently used as a rubefacient. Once a celebrated remedy for cholera:

Preparations.—Spiritus Cajuputi (1 in 50 rectified spirit). Dose, $\frac{1}{2}$ to 1 drachm. Also contained in Linimentum Crotonis ($3\frac{1}{2}$ in 8).

Eucalyptus Globulus.—The leaves distilled in Australia with water yield a yellow fragrant oil, of which the main constituent is *Eucalyptol*, $C_{12}H_{20}O$. Various

preparations of *Eucalyptus* have been introduced for their febrifuge qualities. *Eucalyptus resinifera* yields the red gum called Botany Bay Kino.

CARYOPHYLLI, Cloves.—The dried unexpanded flower buds of *Caryophyllus aromaticus*, cultivated in Penang, Bencoolen, and Amboyna.

Botanical Source.—The *C. aromaticus* is an ever-green shrub with dark purple flowers.

Description.—The reddish brown ball on the top is formed by the imbricated unexpanded corolla, and is situated on the darker brown tubular and four-toothed calyx. Oil should exude from good cloves when pressed with the nail.

Uses.—An aromatic and carminative addition to purgatives, to prevent griping. Dose, 1 to 3 drops. It is useful to microscopists instead of oil of turpentine, to render animal tissues transparent before mounting them in balsam or varnish.

Chemistry.—Cloves contain 18 per cent. of volatile oil and three other volatile bodies called *eugenin* ($C_{10}H_{12}O_2$), *salicylic acid* ($C_7H_6O_3$), and *caryophyllin* ($C_{10}H_{18}O$); 16 per cent. of a peculiar tannic acid, 13

per cent. of gum, 18 per cent. of water and extractive matter. The oil is a yellowish coloured liquid of the sp. gr. of 1.046 to 1.058. It is composed of a light oil isomeric with turpentine ($C_{10}H_{16}$), and a heavy oil called *eugenol* or *eugenic acid* ($C_{10}H_{12}O_2$). The latter combines with alkalies to form crystalline salts, which yield a blue with iron salts. They also give a red with nitric acid, and thus great care must be exercised not to mistake them for morphia.

Infusum Caryophylli.

Cloves, bruised..... $\frac{1}{4}$ ounce
Boiling Distilled Water 10 fluid ounces

Infuse in a covered vessel $\frac{1}{2}$ hour, and strain.

Incompatibles—Salts of iron, zinc, lead, silver, and antimony; lime water and gelatin.

Dose—I to 4 ounces.

Also an ingredient in Infusum Aurantii Compositum, 60 grains to 1 pint; Mistura Ferri Aromatica, $\frac{1}{4}$ ounce to 16 fluid ounces; Vinum Opii, 75 grains to 1 pint.

The oil is used in Confectio Scammoniz, Pilula Colocynthis Composita, and Pilula Colocynthis et Hyoscyami, to prevent griping.

Adulterations.—1. *Mother Cloves* are the *fruits* of the clove-shrub, and are egg-shaped berries about an inch

long and not nearly so fragrant as the buds. They contain starch, and therefore are easily detected by the microscope.

2. The stalks of cloves are used to mix with ground cloves, and may be detected by showing thick walled prosenchymatous cells under the microscope.

PIMENTA, Pimento, *Allspice* or *Jamaica Pepper*.—The dried unripe berries of *Eugenia Pimenta*, grown in the West Indies and imported almost entirely from Jamaica.

Botanical Source.—The *E. Pimenta* is an elegant evergreen tree, growing on calcareous soil.

Description.—Dark brown berries about the size of a pea crowned with the remains of the calyx, and of a deep brown colour, aromatic odour, and spicy taste.

Uses.—Like cloves, aromatic and stimulant. The chief use of pimento is as a culinary spice.

Chemistry.—The berries yield from 3 to 4 per cent. of volatile oil. They are also rich in tannin, striking with persalts of iron an inky black. The decoction is

coloured deep blue with iodine, showing the presence of starch, and the berries also contain a little green fixed oil. The volatile oil is identical in all but odour with that of cloves. The fixed oil is acrid and forms a green solution with ether.

Aqua Pimentæ, 14 ounces to 2 gallons ; distil 1 gallon.
Dose, 1 to 2 ounces.

Oleum Pimentæ.—The oil distilled in Britain from pimento berries. It becomes reddish and afterwards brown by age. Its sp. gr. is 1.037. It is often termed essence of pimento.

GRANATI RADICIS CORTEX,
Pomegranate Root Bark.—The dried bark of the root of *Punica Granatum*, obtained from the South of Europe.

Botanical Source.—The pomegranate is a low tree, with small deciduous leaves and scarlet flowers. Its fruit is a *balausta* with a thick leathery rind, about the size of an apple, and crowned by very evident circular remains of the calyx.

Description.—In thin quills or fragments 3 or 4 inches long. The outer surface greyish and the inner

greyish yellow, finely striated and often with little strips of tough white wood attached. The fracture is short and granular, and the taste astringent.

Uses.—Powerfully astringent; sometimes used in diarrhœa, as an injection for leucorrhœa, and as a gargle. By some it is thought to be slightly anti-periodic. The Chinese and Hindoos use the fresh bark for the expulsion of *tœnia*. The flowers are occasionally prescribed in an infusion under the name of *Flores balaustini*.

Chemistry.—It contains 22 per cent. of a peculiar tannic acid, a little gallic acid, and some mannite. The rind of the fruit was formerly official in the London Pharmacopeia, but is now almost obsolete in British medicine. It is an inferior syncarpous fruit, and is called a *balausta*; very similar in composition and properties to the cortical portion of the root.

Decoctum Granati Radicis, 2 ounces to 2 pints; boil down to 1 pint. Dose, 1 to 2 ounces.

Adulterations.—The bark of the stem and branches (which is less corky), and the bark of the barberry (*Berberis vulgaris*) and box (*Buxus sempervirens*), the decoctions of which are not affected by iron salts.

NATURAL ORDR.—UMBELLIFERÆ

[APIACEÆ].

Herbs or shrubs with usually compound leaves sheathing at the base; flowers, in umbels; calyx, monosepalous and superior; corolla, 5 petals incurved; stamens, 5 epigynous and incurved; ovary, inferior two-celled, with one pendulous ovule in each cell; fruit, a cremocarp, consisting of two mericarps adhering by their face (*commissure*) to a common axis (*carpophore*); seeds, albuminous.

The order is divided into three sub-orders:—

1. *Orthospermeæ*, albumen flat on its face.

Ex. *Ænanthe*, *Daucus*.

2. *Campylospermeæ*, albumen rolled inwards at the edges. Ex. *Conium*.

3. *Cælospermeæ*, albumen curved inwards at the base and apex. Ex. *Coriandrum*.

CONII FRUCTUS, Hemlock Fruit.

The dried fruits of *Conium maculatum*, from indigenous plants.

Botanical Source.—The true hemlock is a biennial herb, having the stem covered with dull purple spots, and not distinctly swollen at the nodes. The leaves are compound and tripinnate with dilated and sheathing petioles. The inflorescence is a compound umbel with both a general and a partial involucre.

Description.—As usually found in commerce the fruit consists not of the entire cremocarp, but of the mericarps only, which have become separated from the carpophore. They are about 1-8th of an inch long, and have their dorsal surface marked with 5 crenate ridges with channels between, which latter are striated longitudinally but are *destitute of vittæ*.

Uses.—Narcotic and sedative, acting especially on the spinal cord; used to allay cough in phthisis and spasmodic asthma; sometimes to ease the pain in cancerous affections. It is a cerebro-spinal poison, paralysing the muscles, and causing death by asphyxia.

Antidotes.—Emetics, purges, and diffusible stimulants.

Chemistry.—They depend for their action on a volatile alkaloid called *conia*, $C_8H_{15}N$, supposed to be combined in the plant with malic acid. Conia is a

colourless oily fluid with strongly alkaline reaction, boiling at 340° F., and having an odour resembling that of mice. It may be isolated by exhausting the seeds with acidulated water, evaporating to a syrup, and adding excess of potassium hydrate. This liquid is then shaken up with ether, which when decanted and spontaneously evaporated leaves the alkaloid. Roughly, the presence of conia may be proved in any part of the hemlock plant, or in any of its preparations by rubbing and heating with a little solution of potassium hydrate. when the "mousey" smell will be distinctly perceptible.

Tinctura Conii.

Hemlock Fruit, bruised	2½ ounces
Proof Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—20 to 60 minims (1 grain of Hemlock fruit in 9 minims.)

According to recent medical statement, Hemlock, in all its preparations, may be given in much larger doses than those indicated in the British Pharmacopœia.

CONII FOLIA, Hemlock Leaves.—The fresh leaves and young branches; also the leaves separated from the branches and carefully dried. gathered from wild British plants when the fruit is beginning to form.

Cataplasma Conii.

Hemlock Leaf, in powder	1 ounce
Linseed Meal	3 ounces
Boiling Water	10 fluid ounces

Extractum Conii.

The expressed juice of the fresh leaves and young branches of Hemlock, evaporated to an extract. (100 lbs. yield about 5½ lbs. of extract.)

Dose—2 to 6 grains.

Pilula Conii Composita.

Extract of Hemlock	2½ ounces
Ipecacuanha, in powder.....	½ ounce
Treacle	q.s.

Dose—5 to 10 grains (2½ grains of Hemlock in 3 grains.)

Succus Conii.

Hemlock Juice.....	3 pints
Rectified Spirit.....	1 pint

Filter after standing 7 days.

Dose.—½ drachm to 1 drachm. (12 minims equal to 1 grain of extract.)

Vapor Coniæ.

Extract of Hemlock	60 grains
Solution of Potash	1 fluid drachm
Distilled Water	10 fluid drachms

Dose—20 minims placed on a sponge, in a suitable apparatus, that the vapour of hot water passing over it may be inhaled.

Adulterations.—1. The leaves of *Chærophyllum Anthriscus*, known by their more or less downy nature.

2. Those of *Æthusa Cynapium* or *Fools' Parsley*, altogether smaller as a whole, and not nearly so acute in the segments of the leaflets.

CARUI FRUCTUS, Carraway Fruits, *vulgo Seeds*.—The dried fruit of *Carum Carui*, cultivated in England and Germany.

Botanical Source.—A biennial herb with bipinnate leaves having linear segments and large vaginas. If any general involucre at all, it only consists of one leaf, while there are no partial involucre.

Description.—The mericarps, which either hang loosely to the carpophore or are altogether separated, are about 1-6th of an inch long and 1-20th in diameter. They are slightly arched, with 5 pale filiform ridges and dark brown channels between, with one vitta in each channel. A pair of vittæ are also visible on the commissure.

Uses.—Aromatic, stomachic, and carminative. Very useful in flatulence, and to prevent the griping of purgatives.

Chemistry.—The active principle is the volatile oil, of which good fruits yield from 4 to 6 per cent. It consists of 2-3rds of *carvol* ($C_{10}H_{14}O$) and 1-3rd of *carvene* ($C_{10}H_{16}$). The former is isomeric with menthol, myristicol, and thymol; and the latter with turpentine.

Aqua Carui.

Carraway Fruit, bruised.....	16 ounces
Water	2 gallons

Distil 1 gallon. (1 of Fruit in 10.)

Also contained in *Confectio Opii* (1 in 10); *Confectio Piperis* (3 in 20); *Pulvis Opii Compositus* (1 in $2\frac{1}{2}$); *Tinctura Cardamomi Composita*; and *Tinctura Sennæ*.

CARUI OLEUM, Oil of Carraway.—

The oil distilled in Britain from carraway fruit. It is colourless or pale yellow, sp. gr. .950, and is an ingredient in *Confectio Scammonii* and *Pilula Aloes Barbadensis*.

FÆNICULI FRUCTUS, Fennel

Fruits.—The ripe fruits of *Fœniculum dulce*, imported from Malta.

Botanical Source.—The fennel is a herb with perennial root, and decomposed leaves with fine fringe-

like segments ; the whole plant growing 3 to 5 feet high, and having a large compound umbel with 6 or 8 long thin rays.

Description.—When undivided the cremocarp is about 1-4th of an inch long and 1-10th in diameter. They are pale coloured, and have 8 brownish ridges with vittæ in the channels, surrounded by a dark brown tissue. The two lateral ridges are double, so that when split each mericarp appears to have 5 ridges.

Uses.—Carminative ; used to relieve flatulence.

Chemistry.—Fennel contains $3\frac{1}{2}$ per cent. of volatile oil, 12 per cent. of fixed oil, and sugar. The volatile oil is analagous to that of anise, and consists of *anethol* or *anise camphor*, $C_{10}H_{12}O$, and a variable portion of a liquid isomeric with turpentine. There are three varieties of oil of fennel in commerce :—Oil of sweet fennel, which has a sweet taste ; oil of bitter fennel, prepared in France, and having a bitter taste ; and Saxon oil of fennel, which resembles the oil obtained from the sweet fennel.

Aqua Fœniculi, 1 lb. to 2 gallons ; distil 1 gallon.

ANISI FRUCTUS, Aniseed.—The fruit of *Pimpinella Anisum*, indigenous to Asia Minor, but cultivated in France, Spain, Malta, and Greece.

Botanical Source.—The anise is a low annual, with cordate and divided radical leaves and biternate cauline leaves. The umbels are many rayed and have no involucre.

Description.—Generally entire cremocarps, about 2-10ths of an inch long, on slender pedicels, and crowned by a pair of short styles. The fruit has 10 pale-coloured ridges, so that it has a prismatic appearance. All the surface is covered with minute hairs.

Uses.—Similar to carraways.

Chemistry.—Aniseed yields about 2 per cent. of a volatile oil, concreting when cooled to 50° F. It consists of an elœopten isomeric with camphor, and a stearopten called *anethol*, or *anise camphor*, $C_{10}H_{12}O$, which fuses at 64.4° F., and when treated with diluted nitric acid forms oil of meadow-sweet, *anisyl hydride* (C_8H_7OH), together with oxalic and anisic acids.

Oleum Anisi, distilled in Europe from anise fruit.

Essentia Anisi (1 of oil in 4 of rectified spirit). Also contained in Tinctura Opii Ammoniata and Tinctura Camphoræ Composita.

The anise fruits are apt to get mixed with conium by mistake, and therefore the two should be carefully compared.

Adulteration.—Spermaceti, detected by shaking up with cold spirit, in which the oil is soluble, and spermaceti insoluble. Camphor may be detected by its odour.

ANETHI FRUCTUS, Dill Fruit.—The dried fruit of *Anethum graveolens*, cultivated in England and imported from Middle or Southern Europe.

Botanical Source.—The dill is an annual herb, with a smooth but finely striated stem, and not over 2 feet high. The leaves are tripinnate, with fringe-like segments, and very broad vaginas. The umbels have neither general nor partial involucre.

Description.—Oval and very flat in form, and usually split into mericarps, which have 5 filiform ridges—3 sharply keeled, and 2 losing themselves in the spreading margin. The vittæ are broad and solitary and fill the whole channels.

Uses.—Aromatic and carminative ; given to infants to relieve flatulence, and the water is used as a vehicle for other medicines.

Chemistry.—The fruit yields about 3 per cent. of volatile oil, a large proportion of which was found by Gladstone to be a hydro-carbon, $C_{10}H_{16}$, which he called Anethene. He also found oil of dill to contain an oxygenated oil, $C_{10}H_{14}O$, which he regarded as identical with carvol.

Aqua Anethi, 1 pound to 2 gallons ; distil 1 gallon.

CORIANDRI FRUCTUS, Coriander Fruit.—The dried ripe fruit of *Coriandrum sativum*, cultivated in Britain.

Botanical Source.—An annual herb, about 2 feet high, with bipinnate leaves, the under ones with broad and the upper with very fine segments. Umbels, with both involucre.

Description.—Always found entire as a pale yellow globular cremocarp, about the size of a white peppercorn, with the remains of style and calyx very distinct on the top. 10 zigzag primary ridges and 8 more prominent secondary ones, and no vittæ in the channels.

Uses.—Aromatic carminative. It is used to disguise nauseous medicines, and in order to prevent griping.

Chemistry.—The active principle is a volatile oil, $C_{10}H_{18}O$, of which the fruit yields 0·5 per cent. Its sp. gr. is 0·87, and it is soluble in ether and glacial acetic acid. The fruit also contains 13 per cent. of fixed oil. It is isomeric with *borneol*, $C_{10}H_{18}O$.

Coriander fruit is a constituent of *Confectio Sennæ*, *Mistura Gentianæ*, *Syrupus Rhei*, *Tinctura Rhei*, *Tinctura Sennæ*.

Oleum Coriandri (Oil of Coriander).—The oil distilled in Britain and described above, enters into the composition of *Syrupus Sennæ* B.P.

CUMINI FRUCTUS, Cumin Fruit.
—The dried fruit of *Cuminum Cyminum*, imported from Sicily, Malta, and Mogadore.

Botanical Source.—An annual umbellifer, with both involucre, growing about 2 feet high.

Description.—Usually unseparated cremocarps, of a brown colour, about $\frac{1}{4}$ of an inch long, 5 filiform primary ridges, and 4 secondary ones on each mericarp, the

latter being hairy; 1 vitta between the primary ridges and 2 on the commissure.

Uses.—A constituent in curry powder, and used in horse medicine.

Chemistry.—Yields a pale yellow volatile oil containing about equal parts of *cymol*, $C_{10}H_{14}$, and *cuminic aldehyd*, $C_{10}H_{12}O$. The former is the hydro-carbon of which turpentine is the hydride.

SUMBUL RADIX, Sumbul Root.—

The dried transverse sections of the root of *Euryangium Sumbul*, imported from Russia and also from India.

Botanical Source.—A perennial plant, about 10 feet high, with a very fragrant milky juice; discovered in 1869 near Samarkand.

Description.—Nearly round pieces, 1 to $1\frac{1}{2}$ inch thick and 3 to 5 inches long, dusky brown and covered with bristly fibres. Internally it is porous, and has an odour of musk.

Uses.—Nervous stimulant and antispasmodic, and prescribed in the form of tincture as a substitute for musk in hysteria and epilepsy.

Chemistry.—The root yields 9 per cent. of balsamic resin, about 0·75 per cent. of sumbulic, sumbulolic, or angelic acid, a small quantity of volatile oil, and a trace of valerianic acid. The resin is converted by solution of potassium hydrate into a crystalline potassium salt of sumbulamic acid, which has a strong musk-like odour, and appears to differ from the sumbulic acid before noticed.

Tinctura Sumbul.

Sumbul Root, in coarse powder 2½ ounces

Proof Spirit 1 pint

Dose—10 to 30 minims.

ASSAFŒTIDA (more correctly *Asafætida*).—A gum-resin, obtained in Affghanistan and the Punjaub by incision from the living root of *Narthex Asa-fætida*.

Botanical Source.—The *N. Asa-fætida* is a magnificent herb, rising to 10 feet in height, and with radical leaves bipinnate in form and 1½ feet long. It has flowered twice only since 1838 in the Edinburgh Botanical Gardens.

Collection.—A thin slice is cut from the crown of the root, and the juice which exudes is scraped off after a few days, the root being kept covered in the meantime

by twigs and stones. The finest gum-resin comes solely from the leaf bud at the centre of the root crown.

Description.—In irregular, softish masses of tears, white when first cut into, but becoming purplish pink by exposure, and having a very nauseous alliaceous odour. When rubbed with water it forms a milky emulsion, and when treated with sulphuric acid it yields a solution, which when diluted and neutralised, has a bluish fluorescence. A very fine quality in pure tears exists, but is rarely met with.

Uses.—Anti-spasmodic, expectorant, stimulant. It is of great use in hysteria, nervous debility, flatulency, spasmodic cough, and other affections of the chest, where there is no inflammation. But for the ill flavour of the drug, the *Spiritus Ammoniaë Fœtidus* is a most valuable medicine. Dose, 5 to 20 grains, best given in the form of emulsion. The gum dissolves in the water suspending the oil and resin.

Chemistry.—Contains 3 to 5 per cent. of volatile oil, 40 per cent. of gum, and 60 per cent. of resin. The oil is at first neutral, but by keeping it becomes acid and evolves sulphuretted hydrogen. Owing to its horrible odour it has not been properly investigated, but is

supposed to be a sulphur compound of *allyl*. The resin is not entirely soluble either in ether or chloroform, but is soluble in warm nitric acid. It contains *ferulaic acid*, $C_{10}H_{10}O_4$. From the presence of sulphur in the oil, pills containing asafœtida should never be silvered, as they become black. The best coating is ethereal tincture of tolu.

Enema Asafœtidæ.

Asafœtida	30 grains
Distilled Water	4 fluid ounces

Rub the Asafœtida with the water, added gradually, for one enema.

Pilula Asafœtidæ Composita. (Pil. Galban. Co. P.L.)

Asafœtida	2 ounces
Galbanum	2 ounces
Myrrh	2 ounces
Treacle, by weight	1 ounce

(3 grains each Aloes, Galbanum, and Myrrh, in 10½ grains.)

Dose—5 to 10 grains.

Spiritus Ammoniaë Fœtidus.

Asafœtida	1½ ounce
Strong Solution of Ammonia	2 fluid ounces
Rectified Spirit	20 fluid ounces

Macerate 24 hours, distil, and add the Ammonia to the product.

Dose—½ to 1 drachm (4 grains of Asafœtida in 1 fluid drachm.)

Tinctura Asafœtidæ.

Asafœtida, in small fragments 2½ ounces
Rectified Spiritad. 20 fluid ounces

Macerate 7 days, then filter.

Dose—¼ to 2 drachms (7 grains of Asafœtida in 1 fluid drachm.)

Also contained in *Pilula Aloes et Asafœtidæ* (1 in 4).

GALBANUM.—A gum-resin obtained from an unascertained umbelliferous plant. Imported from India and the Levant.

Botanical Source.—Although not perfectly certain, the source of galbanum is generally believed to be the *Ferula Galbaniflua*, a plant growing on the slopes of the mountains of Northern Persia.

Collection.—So far as is known, the gum-resin appears to exude spontaneously from the bases of the leaves.

Description.—In irregular tears about the size of a hazel nut, having a tendency to agglutinate and form a greenish yellow to a brown colour. In one variety the tears are translucent, and in another dull, waxy, and light yellow—the latter quality being that rarely found

in agglutinated masses. Galbanum has a heavy peculiar odour, and an acrid and alliaceous taste.

Uses.—Resembles asafoetida in action, but is less energetic. It is employed principally in the form of plaister, as an external stimulant to chronic tumours. Dose, in the form of an emulsion, 10 to 30 grains.

Chemistry.—Galbanum contains about 7 per cent. of volatile oil, 32 per cent. of gum, and 60 per cent. of resin. The oil, obtained by distillation with water, is isomeric with turpentine, boils at 325° F., and smells of the drug. The resin is soft and easily soluble in ether and alkalies, but only partially in carbon disulphide. When heated to 212° F. with hydrochloric acid, and the acid solution shaken with chloroform, the latter leaves on evaporation colourless needle-shaped crystals of *umbelliferone*, $C_9H_6O_3$. This body is yielded not only by galbanum, but also by all umbelliferous resins. It is soluble in water, and on adding an alkali the solution shows a bright blue fluorescence, immediately destroyed by an acid.

Emplastrum Galbani.

Galbanum	1 ounce
Ammoniacum	1 ounce
Yellow Wax	1 ounce
Lead Plaister	8 ounces

Also contained in Pilula Asafoetida Composita (1 in 3½ nearly).

AMMONIACUM.—A gum resin, obtained by exudation from *Dorema Ammoniacum*, and collected in Persia and the Punjaub.

Botanical Source.—The *D. Ammoniacum* is a herb with a perennial root, and having a flower-stem which grows 6 to 8 feet high perfectly leafless, and bears numerous umbels of white downy flowers.

Collection.—The stem becomes about June so gorged with juice that it exudes at the slightest puncture, and just then the plant is attacked by a beetle which pierces it, and the juice running out hardens in tears, which are collected.

Description.—There are two qualities:—1. *Tear Ammoniacum*, which is the best, and is found in roundish tears from the size of a pea to that of a hazel nut. They have a creamy yellow to reddish brown colour externally, and are brittle, opaque and milky white inside the fracture, having a waxy lustre. 2. *Lump Ammoniacum* (the inferior quality) is in brittle masses of agglutinated tears, usually mixed

with fragments of the mericarp of the fruit of the plant and other impurities.

Both varieties have a faint odour and nauseous taste, and when rubbed with water they form white emulsions. They are also coloured yellow by solution of potassium hydrate, and bright orange by chlorinated lime.

Uses.—Expectorant, and generally stimulating to the mucous membranes. Externally it is a resolvent to scirrhus tumours, the B.P. plaister being an excellent application for “housemaid’s knee.”

Dose, 10 to 20 grains in emulsion with yolk of egg.

Chemistry.—Contains 2 per cent. of volatile oil, 28 per cent. of gum, and 70 per cent. of resin. It however yields no umbelliferone. The researches on its oil and resin are not satisfactory as yet.

Test to distinguish between Ammoniacum, Asafœtida, and Galbanum.—Warm the gum-resin (or, better, the resin extracted by alcohol or ether) with *strong* hydrochloric acid, and then slowly add spirit of wine. Ammoniacum is not affected; Asafœtida turns dirty green; and Galbanum becomes bluish violet.

Emplastrum Ammoniaci c Hydrargyro.

Ammoniacum	12 ounces
Mercury	3 ounces
Olive Oil	1 fluid drachm
Sublimed Sulphur	8 grains.

Mistura Ammoniaci.

Ammoniacum, in coarse powder	$\frac{1}{2}$ ounce
Distilled Water	8 fluid ounces

Dose— $\frac{1}{2}$ to 1 ounce (14 grains of Gum in 1 fluid ounce.)

Also contained in Emplastrum Galbani (1 in 11); Pilula Ipecacuanhæ cum Scillâ (1 in 7); Pilula Scillæ Composita (1 in $6\frac{1}{4}$).

Table for the Detection of the Umbelliferous Gum-Resins.*—All are imperfectly attacked both by chloroform and ether; the alcoholic solutions are rendered turbid by ammonium hydrate, and give no precipitate with alcoholic ferric chloride (1 in 10).

I. *The drugs contain sulphur* (by fusion with nitre and testing with BaCl_2).

A. Yields umbelliferone by dry distillation.

1. Hydrochloric acid colours the petroleum spirit extract residue reddish yellow; the chloral test colours it green.—*Persian Sagapenum*.

* E. HIRSCHSOHN, "Pharmaceutical Journal," Nov. 17, 1877.

2. Hydrochloric acid colours the residue blue-violet; the chloral test colours it rose-colour to raspberry-red and violet.—*Levant Sagapenum*.

3. Not coloured by hydrochloric acid. The solution of the drug in sulphuric acid is yellow-brown with a blue fluorescence. Potassium nitrate colours the gum-resin malachite green.—*Ordinary Asafætida*.

B. Yields no umbelliferone by dry distillation.

1. Sodium carbonate solution colours the drug light brown, and the extract is not altered by acetic acid or lead acetate.—*Asafætida from Ferula alliacea*.

II. *The drugs contain no sulphur.*

A. Yields umbelliferone by dry distillation.

1. The petroleum spirit extract residue is coloured by hydrochloric acid and the chloral test.

a. Hydrochloric acid colours reddish yellow; the chloral test colours green.—*Persian Galbanum*.

b. Hydrochloric acid colours red-violet; the chloral test colours greenish.—*Levant Galbanum, as at present in commerce*.

c. Hydrochloric acid colours violet-blue; the chloral test carmine-red.—*Older specimens of Levant Galbanum*.

2. Hydrochloric acid gives no colour; the chloral test colours light brown.—*African Ammoniacum*.

B. Yields no umbelliferone by dry distillation.

1. Chloride of lime solution colours the gum-resin orange-yellow.—*Persian Ammoniacum*.

2. Chloride of lime solution produces no colour; lead acetate gives no precipitate.

a. Iodine solution is not altered; the chloral test colours greenish.—*Olibanum*.

3. Chloride of lime solution produces no colour; lead acetate gives a precipitate; bromine solution produces no colour, or only yellowish; ferric chloride, green.—*Opoponax*.

SUB-CLASS III.—COROLLIFLORÆ.

Plants having both calyx and corolla united and the stamens epipetalous; rarely epigynous or hypogynous.

This sub-class has 3 sub-divisions:—

1. Epigynæ, having the ovary inferior.

2. Hypostamineæ, having the ovary superior and the stamens hypogynous.

3. Epipetalæ, having the ovary superior and the stamens epipetalous.

SUB-DIVISION I.—EPIGYNÆ.

NATURAL ORDER.—CAPRIFOLIACEÆ.

SAMBUCI FLORES, Elder Flowers, *Fleurs de Sureau*.—The fresh flowers of the *Sambucus nigra*, indigenous to Central and Southern Europe, and also a native of England and Ireland, but not of Scotland.

Botanical Source.—An arborescent plant with opposite compound and imparipinnate leaves having ovate, serrate, and cuspidate leaflets. Inflorescence a broad flat cyme of small flowers which have a monopetalous rotate, cream coloured corolla; stamens, 5: stigma 3-lobed and sessile.

Collection.—The bunches of flowers are laid in heaps for some hours, when a little heat being developed the corollas become detached, and are then rubbed or shaken off from the stalks and dried rapidly.

Uses.—Simply employed as a flavouring agent, and as a cooling external application.

Chemistry.—The flowers yield by distillation slight traces of some volatile acids, and a small amount of an essential oil. This latter is of a buttery consistence, has the characteristic smell of elder, and is easily decomposed by exposure to the air.

Aqua Sambuci (1 in 1).

The flowers may also be digested with lard to form an ointment (1 in 1), and the fruit, which is a globular black berry, is used for making elder wine.

NATURAL ORDER.—CINCHONACEÆ.

[*This is regarded by many botanists as a Sub-order of the Natural Order RUBIACEÆ.*]

CATECHU PALLIDUM, Pale Catechu ; Gambier, Gambir or Terra Japonica.—An extract from the leaves and young shoots of *Uncaria Gambier*, chiefly imported from Singapore.

Botanical Source.—The *Uncaria Gambier* is a strong climbing shrub and is a native specially of the Islands at the eastern end of the Straits of Malacca ;

of the Eastern Archipelago generally, and of Ceylon. Another plant known as *U. acida* is also sometimes employed.

Collection.—The leaves and shoots are boiled with water in an iron pan and then squeezed. The decoction is evaporated to a syrupy consistence, and then poured into buckets and kept mixed while cooling by pushing a stick up and down. When cooled to the consistence of clay, the extract is cut into cubes and dried.

Description.—In cubes, either free or coherent, each about an inch square, which are externally brownish and internally pale, dull, reddish yellow. They are odourless, friable, and have a dull fracture and an astringent taste, at first slightly bitter, but subsequently decidedly sweetish. Examined microscopically the cubes are found to be built up of small needle-shaped crystals.

Uses.—An astringent in diarrhœa and Pyrosis, but chiefly consumed by the dyeing and tanning trades, owing to the tannic acid.

Chemistry.—Very similar to Cutch. (See *Acacia Catechu*, page 146.) Active principle, tannic acid.

Infusum Catechu.

Pale Catechu, in coarse powder	160 grains
Cinnamon, bruised	30 grains
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel $\frac{1}{2}$ hour, and strain.

Incompatibles.—The alkalies and alkaline earths; most metallic salts; gelatin; most vegetable infusions containing an alkaloid, as Decoctum Cinchonæ.

Dose—I to 2 ounces.

Pulvis Catechu Compositus.

Pale Catechu, in fine powder	4 ounces
Kino do.	2 ounces
Rhatany Root do.	2 ounces
Cinnamon Bark do.	1 ounce
Nutmeg do.	1 ounce

Dose—20 to 40 grains (8 grains of Catechu and 4 grains each of Kino and Rhatany in 20 grains).

Tinctura Catechu.

Pale Catechu, in coarse powder	2½ ounces
Cinnamon Bark, bruised	1 ounce
Proof Spirit	ad. 20 fluid ounces

Macerate 7 days, strain, press, and filter.

Dose— $\frac{1}{2}$ to 2 drachms (7 grains Catechu in 1 fluid drachm.)

Trochisci Catechu.

Pale Catechu, in powder	720 grains
Refined Sugar, in powder	25 ounces
Gum Acacia, in powder	1 ounce
Mucilage of Gum Acacia	2 fluid ounces
Distilled Water	q.s.

Divide into 720 lozenges.

Dose—I or more. (Each lozenge contains 1 grain of Catechu.)

CINCHONÆ FLAVÆ CORTEX,
Yellow Cinchona Bark.—The bark of *Cinchona*
Calisaya, collected in Bolivia and Southern Peru.

Botanical Source.—The *C. Calisaya* is a tall tree generally overtopping all others in the forests of Bolivia. It is evergreen, and bears fine paniced cymes of pink flowers.

Collection.—By preference the tree is cut down just above the ground, but so as to leave a “stool” from which young shoots may once more spring. The bark is loosened and denuded of its *periderm* (i.e., *epiphylæum* and a portion of the *mesophylæum*, which has become dry and useless with age) by beating with a mallet, and is then cut off in slips from 15 to 18 inches long and about 4 wide. If these are from the younger branches they are at once placed in the sun to dry and curl up into “quill bark”; but if from the older stem they are dried under slight pressure into “flat bark.” When dry, the pieces are packed in *serons* or bales of hides.

Description.—There are two varieties of this bark:
—1. *Flat yellow bark*, in pieces deprived of the *periderm* about 3 lines thick and 1 to 3 inches wide, but varying

much in length. The bark is compact, heavy, and marked externally by longitudinal depressions, which are broad, shallow, and irregular, of which the internal or liber surface is tawny yellow, and most distinctly fibrous in structure. When broken the true bark has a very short and finely fibrous fracture. The odour is slightly aromatic, and the taste bitter.

2. *Quill yellow bark*, in double quills (*i.e.*, rolled in at both edges) from $\frac{3}{4}$ to $1\frac{1}{2}$ inch in diameter. The *periderm* is greyish or silvery white externally, and is thick, rugged, and marked with deep cracks with slightly elevated edges, and running both along and across. The inner surface is dark brown and fibrous, and the fracture is very shortly fibrous.

Uses.—Anti-periodic, tonic and astringent. Administered in all kinds of intermittent fevers and intermittent neuralgic affections. The doses of the bark, or quinine, should always be taken *in the interval between the paroxysms*, and never while the fit is on. Either a full dose may be administered just before the expected return of the paroxysm, or a series of small and separate doses may be kept up during the whole time of intermission. *As a tonic* it is useful in all cases of convalescence from

acute diseases, in dyspepsia and scrofula; and as an *astringent* it is employed in gargles.

The dose of sulphate of quinine is from 1 to 10 grains, and of yellow bark in powder, as a tonic, 15 to 30 grains, and as an anti-periodic, 1 to 2 drachms.

Tests.—1. *Qualitative.*—All true cinchona barks, really containing one or other of the cinchona alkaloids, will give off deep crimson fumes when heated in a test tube.

2. *Quantitative.*—The official process is as follows:—Boil 100 grains of the bark, reduced to very fine powder, for a quarter of an hour in one fluid ounce of distilled water acidulated with 10 minims of hydrochloric acid, and allow it to macerate for 24 hours. Transfer the whole to a small displacement tube, and after the fluid has ceased to percolate, add at intervals about 1½ ounces of similarly acidulated water, or until the fluid, which passes through, is free from colour. Add to the percolated fluid, solution of subacetate of lead, until the whole of the colouring matter has been removed, taking care that the fluid remains acid in re-action. Filter and wash it with a little distilled water. To the filtrate add about 35 grains of caustic potash, or as much as will

cause the precipitate which is first formed to be nearly re-dissolved ; and afterwards six fluid drachms of pure ether. Then shake briskly ; and having removed the ether, repeat the process twice, with three fluid drachms of ether, or until a drop of the ether employed, leaves on evaporation, scarcely any perceptible residue. Lastly, evaporate the mixed ethereal solutions in a capsule. The residue, which consists of nearly pure quinia, when dried should weigh not less than two grains, and should be readily soluble in diluted sulphuric acid.

NOTE.—This process being for the estimation of Quinine only, is not applicable to the pale or red barks in which the value depends, not on the Quinine only, but on the entire mixed alkaloid which the bark contains. When examining the pale or red barks, therefore, the B.P. orders the substitution of chloroform for ether in the above process. The residue weighed is considered to be “mixed alkaloids.”

Chemistry.—Yellow cinchona bark contains a series of chief alkaloids—viz., quinia, cinchonina, quinidia, cinchonidia, and the amorphous alkaloids all in combination with kinic, quinovic, cincho-tannic, and cincho-fulvic acids, together with a trace of volatile oil and the usual woody fibre.

The following are short descriptions of the more important constituents:—

QUINIA, $C_{20}H_{24}N_2O_2 \cdot 3H_2O$, may be prepared by dissolving sulphate of quinine in water by the aid of a little sulphuric acid, and then precipitating with the slightest possible excess of ammonia. The precipitate, dissolved in strong alcohol, crystallises in silky prisms, having three molecules of water of crystallisation. It is slightly soluble in water (1 in 350), and very soluble in ether, alcohol, and chloroform. Its aqueous solution is slightly alkaline to test paper. It dissolves readily in acids to form salts, and is re-precipitated by ammonia soluble in excess.

TESTS.—1. Chlorine water, added to a solution of quinine and followed by the addition of a drop or two of ammonia, produces a bright green colour.

2. If a drop or two of potassium ferro-cyanide in solution be added *between* the chlorine water and ammonia, an evanescent red is produced.

3. By dissolving the salt of quinine in proof spirit containing 5 per cent. of sulphuric acid, and then adding tincture of iodine, a precipitate is formed which consists of herapathite. It is in brilliant crystals of greenish tint

and metallic lustre. Its formula has been defined as $2C_{20}H_{24}N_2O_2 \cdot I_6 \cdot 3(H_2SO_4) \cdot 3H_2O$; different formulæ are however given.

Quinidia, $C_{20}H_{24}N_2O_2 \cdot 2H_2O$, is isomeric with quinia. It agrees with that alkaloid in bitterness, fluorescence, and in the chlorine water and ammonia re-action. It may, however, be separated by precipitation by potassium iodide, with which it forms an insoluble iodide of quinidia. It crystallises from ether in long rhombic prisms.

Cinchonia, $C_{20}H_{24}N_2O$, crystallises in four-sided prisms, fusible at 320 F. It is much more soluble in water than quinia (1 in 250); it is soluble in alcohol, but only very slightly soluble in ether, this latter point forming a simple method of separating it from quinia, which is freely soluble. Its salts do not show fluorescence in solution, and when treated with chlorine water, and ammonia, they yield a white precipitate instead of a green colour. The precipitated solutions of its salts by ammonia are practically insoluble in excess.

Cinchonidia crystallises from alcohol in anhydrous, rhombic prisms. It is isomeric with cinchonia, from which it may be separated by precipitation of the mixed

sulphates by Rochelle salt, when it is precipitated as tartrate, leaving the cinchonia in solution.

Quinicine and **Cinchonicine** are amorphous alkaloïds, produced in the making of the quinine. They are not crystallisable, and are doubtless formed by the action of the mineral acids; possessing about the same relation to quinine and cinchonine, as treacle and sugar bear to each other. They are sometimes classed together under the name of quinoïdine.

Kinic (Quinic) acid, $C_7H_{12}O_6$, crystallises in bold plates, soluble in two pints of water; less soluble in alcohol, and almost insoluble in ether. It may be precipitated by heat, and yields when treated with manganese peroxide and sulphuric acid, *Quinone*, $C_6H_4O_2$, together with carbonic anhydride and water. The quinone collects in deep yellow prisms on the cool part of the apparatus. By destructive distillation, *per se*, kinic acid yields benzol, benzoic, and salicylic acids, with hydro-quinone, $C_6H_6O_2$, and carbonic anhydride.

Quinovic acid, $C_{24}H_{38}O_4$, only exists in small quantity. It is soluble in boiling alcohol.

Cincho-tannic acid, $C_{14}H_{12}O_7H_2O$, exists to a con-

siderable extent. It may be precipitated from a decoction of cinchona bark (which has been previously freed by magnesia from the presence of cinchona-red) by the addition of plumbic acetate; and the precipitate so produced, having been suspended, is treated with sulphuretted hydrogen, which removes the lead, leaving the cinchotannic acid in solution. It is very soluble in water, alcohol, and ether. Its solutions are turned green by ferric-chloride; and, in contact with excess of alkalies, it undergoes decomposition, yielding cinchona-red. It also yields precipitates with tartar emetic and gelatin.

Cinchona-red, $C_{12}H_{14}O_7$, is called "cincho-fulvic acid," and is the natural colouring matter of the bark, from which it may be extracted by the action of alkalies with which it forms a deep red solution. It is precipitated from its ammoniacal solution by hydrochloric acid.

Decoctum Cinchonæ Flavæ.

Yellow Cinchona Bark, in coarse powder. $1\frac{1}{4}$ ounce
 Distilled Water 1 pint

Boil 10 minutes in a covered vessel, and strain when cold;
 make up product to 20 fluid ounces.

Incompatibles—The alkalies and their carbonates, lime water,
 most metallic salts, and gelatin.

DOSE—1 to 2 ounces.

In the decoction of yellow bark many of the alkaloids which have been extracted by the boiling water are precipitated on cooling by the cincho-tannic acid, and are consequently filtered out.

Extractum Cinchonæ Flavæ Liquidum.

Yellow Cinchona Bark, in coarse powder. 16 ounces
 Rectified Spirit 1 fluid ounce
 Distilled Water q.s.

Product, 4 ounces; sp. gr. 1.100. (1 drachm is equivalent to $\frac{1}{2}$ ounce of Bark, or 10 fluid ounces of Infusum Cinchonæ Flavæ.)

Dose—10 to 30 minims.

As usually met with, this preparation contains scarcely any alkaloids at all, and must simply be viewed as an astringent, and even then one of weak strength.

Infusum Cinchonæ Flavæ.

Yellow Cinchona Bark, in coarse powder. $\frac{1}{2}$ ounce
 Boiling Distilled Water 10 fluid ounces

Infuse in a covered vessel for 2 hours, and strain.

Incompatibles—The alkalies and their carbonates, lime water, most metallic salts, and gelatin.

Dose—1 to 2 ounces.


Tinctura Cinchonæ Flavæ.

Yellow Cinchona Bark, in moderately fine powder 4 ounces
 Proof Spirit ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose— $\frac{1}{2}$ to 2 drachms. (11 grains of Bark in 1 fluid drachm.)

QUINIÆ SULPHAS



is prepared according to the B.P. by boiling three fluid ounces of hydrochloric acid with ten pints of water, and placing one pound of yellow cinchona bark in coarse powder in a porcelain basin, and adding to it as much of the diluted hydrochloric acid as will render it thoroughly moist. After maceration, with occasional stirring for 24 hours, place the bark in a displacement apparatus and percolate with the diluted hydrochloric acid, until the solution which drops through is nearly destitute of bitter taste. Into this liquid pour 4 pints of solution of soda, agitate well, let the precipitate completely subside, decant the supernatant fluid, collect the precipitate on a filter and wash it with cold water until the washings cease to have colour. Transfer the precipitate to a porcelain dish containing a pint of water; and applying to this a steam heat, gradually add diluted sulphuric acid until very nearly the whole precipitate has been dissolved, and a neutral liquid has been obtained. Filter the solution while hot, wash the filter with boiling hot distilled water, concentrate until a film remains on the surface of the solution and set aside to crystallise.

The crystals should be dried on filtering paper without the application of heat.

In this process the hydrochloric acid dissolves out the quinia and cinchonia as hydrochlorates, setting free kinic acid. The soda precipitates the alkaloids by combining with the acids. The precipitated alkaloids are then carefully treated with just enough of diluted sulphuric acid to obtain a perfectly neutral sulphate, when the sulphate of quinine crystallises out, being being much less soluble than the sulphate of cinchonine which remains in solution.

In making quinine on a larger scale this process is modified by the use of excess of lime instead of solution of soda. The precipitated alkaloids, together with the excess of lime, are washed and dried by pressure between calico, and the quinine is dissolved out from the cake by rectified spirit.

Sulphate of quinine is in silky crystals, slightly soluble in cold water (1 in 740), much more so in boiling water (1 in 30). It is also soluble in rectified spirit. It is exceedingly soluble in water acidulated with sulphuric acid, being thereby converted into the acid sulphate, $C_{20}H_{24}N_2O_2 \cdot H_2SO_4 \cdot 7H_2O$.

Owing to its costliness sulphate of quinine is subjected

to many adulterations:—1. Test for the presence of salicin by moistening a little with cold strong sulphuric acid, when it should not become red. The solution should then be heated gently, and if sugar, starch, gum, or other organic impurities be present, it will darken.

2. It should be tested for the presence of cinchonine by dissolving 10 grains in $\frac{1}{2}$ ounce of water by the aid of 10 minims of B.P. diluted sulphuric acid, and adding a few drops of ammonia to precipitate the quinine; the whole should then be shaken up with $\frac{1}{2}$ fluid ounce of pure ether and allowed to stand at rest, when the quinia dissolves, and any cinchonia present is seen as a crystalline matter floating on the surface of the lower stratum of fluid. If the upper stratum of ether be now carefully and entirely removed by a pipette and evaporated spontaneously in a weighed capsule, and, when dried, heated for a few minutes in the water bath, the residue of pure quinine should weigh 8·6 grains.

Sulphate of quinia is used in making the following:—

Pilula Quiniæ.

Sulphate of Quinia	60 grains
Confection of Hips	20 grains

Dose—2 to 10 grains (3 grains of Quinine in 4 grains).

Tinctura Quiniæ.

Sulphate of Quinia	160 grains
Tincture of Orange Peel	20 fluid ounces

Dissolve with a gentle heat, and after 3 days filter.

Dose— $\frac{1}{2}$ to 2 drachms (1 grain of Quinine in 1 fluid drachm).

Tinctura Quiniæ Ammoniata.

Sulphate of Quinia	160 grains
Solution of Ammonia	2½ fluid ounces
Proof Spirit	17½ fluid ounces

Dissolve the Quinia in the Spirit by a gentle heat, and add the ammonia.

Dose— $\frac{1}{2}$ to 2 drachms (1 grain of Quinine in 1 fluid drachm).

NOTE.—This deposits on keeping, owing to loss of ammonia, the *excess* of which alone retains the quinia in solution.

Vinum Quiniæ.

Sulphate of Quinia	20 grains
Citric Acid	30 grains
Orange Wine	20 fluid ounces

Dissolve first the acid and then the Quinine in the wine. allow it to remain 3 days, shake occasionally, and filter.

Dose— $\frac{1}{2}$ to 1 ounce (1 grain of Quinine in 1 fluid ounce).

Quinia is also contained in *Ferri et Quiniæ Citras*, in which it exists as a slightly acid citrate to an amount equivalent to 16 per cent. of pure quinia.

CINCHONÆ PALLIDÆ CORTEX,
Pale Cinchona Bark.—The bark of *Cinchona officinalis* (var. *Condaminea*), collected about Loxa, in Ecuador.

Description.—The finest, or “pale crown” bark, is in thin quills from 1-8th to $\frac{3}{4}$ inch in diameter, twisted and often double. The thickness of the bark varies from that of paper up to about 1-10th of an inch. Externally it is dark grey splashed with silver grey, and on fine pieces are numerous large lichens, while the whole outside is cracked transversely at close intervals. Internally it is bright yellowish brown, and very fibrous. When broken the fibres are seen to be in the endophlœum only. Besides this we meet with “ashy crown” bark (the produce of *C. Macrocalyx*), which, together with several other varieties, should be carefully studied.

Uses.—As yellow bark, but not so rich in anti-periodic properties, although more astringent, owing to its containing more quino-tannic acid.

Dose, of the powder, 10 to 60 grains.

Chemistry.—Contains very little quinia, but from

about .5 to 1 per cent. of cinchonia and quinidia, chiefly the latter.

Official Test.—To be performed exactly as for yellow bark, only starting with 200 grains, and using chloroform instead of ether, when the residue should weigh at least one grain.

Tinctura Cinchonæ Composita.

(*Huxham's Tincture of Bark.*)

Pale Cinchona Bark, in moderately fine powder	2 ounces
Bitter Orange Peel, bruised	1 ounce
Serpentary Root, bruised	$\frac{1}{2}$ ounce
Saffron	60 grains
Cochineal, in powder	30 grains
Proof Spirit.....	20 fluid ounces

Macerate 48 hours, then percolate.

Dose— $\frac{1}{2}$ to 2 drachms. ($5\frac{1}{2}$ grains of bark in 1 fluid drachm).

This tincture, under its popular name, was formerly in extensive demand. It may still be regarded as an admirable preparation. Pale cinchona bark is also contained in *Mistura Ferri Aromatica*.

CINCHONÆ RUBRÆ CORTEX,
Red Cinchona Bark.—The bark of *Cinchona succirubra*, collected on the western slopes of the Chimborazo.

Description.—In flat or channelled pieces coated with their periderm, and sometimes reaching $\frac{1}{2}$ inch in thickness. The inner surface is brick-red and fibrous, and the whole breaks with a short fibrous fracture. At present it is not extensively used, but it is being rapidly planted in India, and may soon be expected to be in larger demand.

Uses.—For quinine manufacture.

Chemistry.—Contains chiefly quinia and quinidia, with some cinchonia. According to the official test performed on 100 grains, and working with chloroform as for pale bark, it should yield not less than 1.5 per cent. of mixed alkaloids; but it usually contains much more than that, and about one-third of the whole is frequently quinia.

CINCHONÆ LANCIFOLIÆ
CORTEX, Columbian or Carthagena
Bark.—The bark of *C. lancifolia*, imported *via* Santa
Fè and Carthagena. *Not official.*

Uses.—One of the chief, if not actually the principal source of the quinine manufacture in England. It is

rich in alkaloids, but variable. It is imported both in "quills" and "flats."

The remaining bark which is used for quinine manufacture is that of *C. Pitayensis*, known in commerce as Pitayo bark. It comes from Columbia, and is usually imported in much broken fragments. It is also rich in alkaloids, quinine prevailing.

Complete Analysis of Cinchona Bark for Alkaloids.—The following is a more complete process than the official one, and has the additional advantage of rapidity when only pursued as far as the obtaining of the total mixed alkaloids. The second part involving separation is more tedious.

Mix 300 grains of powdered bark, dried at 212° F., with milk of lime (75 grains of slaked lime to 750 grains of water). Dry the mixture slowly and then boil it with 7 fluid ounces of rectified spirit. Pour off the clear liquid; boil again with half as much more alcohol; filter, and wash the powder with 3½ ounces more alcohol. From the mixed liquors precipitate the calcium as sulphate by a few drops of diluted sulphuric acid. Filter; distil off the spirit, and pour the residual liquid into a capsule, and heat it on a

water bath, until the spirit is wholly expelled. Filter the remaining liquor, which contains all the alkaloids in the form of acid sulphates, and wash the residue (quinovic acid and fatty matter) with water, slightly acidulated with sulphuric acid. The filtrate and washings, reduced to about $\frac{1}{2}$ fluid ounce, are now treated, while still warm, with caustic soda in slight excess. Wash the precipitate with a very little water; press it between folds of blotting paper, and dry on a water-bath. The weight, divided by three, gives the per-centage of the total alkaloids.

The dried alkaloids are then shaken up with ten times their weight of ether, which dissolves quinine and quinicine (A) and leaves cinchonine, cinchonidine, and quinidine (B), which may be treated as follows :—

The portion soluble in ether (A) is evaporated to dryness, and the residue dissolved in ten parts of proof spirit, acidulated with 1-20th of diluted sulphuric acid. The whole is warmed, and tincture of iodine added in excess, and the precipitate collected, washed, and dried at 100° F. One part of this precipitate equals .565 parts of quinia. To the filtrate add a few drops of sulphurous acid; evaporate to remove the alcohol, and then add

caustic soda, which will precipitate the quinicine, and any other amorphous alkaloid present.

The portion (B) is dissolved in 40 parts of hot water by the aid of a little diluted sulphuric acid, so as to produce perfectly neutral sulphates. On the addition of potassium iodide, the quinidine is precipitated, and the precipitate may be collected, dried, and weighed. To the filtrate Rochelle salt is added, and the precipitated tartrate of cinchonidine collected after 12 hours, dried and weighed; and, finally, cinchonine is precipitated from the filtrate by a slight excess of sodium hydrate, and washed and dried.

IPECACUANHÆ RADIX,
Ipecacuanha Root, *Ipecacuan.*—The dried root of *Cephaëlis Ipecacuanha*, imported from Brazil.

Botanical Source.—The *Cephaëlis Ipecacuanha* is a small shrub with a creeping root, growing in the shady forests of tropical South America, from Rio de Janeiro to Pernambuco. The root of commerce is obtained chiefly from the Province of Matto Grosso.

Collection.—The root is collected by men called Poayeros, and afterwards dried in the sun. The collec-

tion goes on continuously, except during the rainy season.

Description.—In pieces of 3 or 4 inches in length, of a dusky grey or dull brownish colour. It is hard, breaks readily with a granular fracture, and it is white or greyish inside. When broken it is found to consist of a cortical portion, which constitutes 75 to 80 per cent. of the whole root, and a medullium, which is tough and slender. The active portion is the cortical, and therefore the best quality is that showing the greatest amount of this portion. This being so easily separable, the root is often found with fragments of it detached. As met with in commerce, ipecacuanha root contains the maximum diameter of about 2-10ths of an inch, and its surface is covered with rings which penetrate in some cases almost to the wood, from which circumstance its popular name of *annulated* is derived. The inferior quality, distinguished as “New Granada Ipecacuanha,” is sometimes imported, which is larger in size and has a distinctly radiate arrangement of the medullium.

Uses.—In small doses it is an expectorant and diaphoretic; in large doses an emetic. It is found useful

in chronic dysentery. When given as an emetic it is not so depressing as *antimonium tartaratum*, but it is not so active an emetic as zinc sulphate.

Dose as an expectorant $\frac{1}{2}$ to 2 grains; as an emetic 15 to 30 grains.

Chemistry.—Ipecacuanha contains an alkaloid called *emetine*, combined with ipecacuanhic acid. It also yields a small portion of a fetid volatile oil, to which its faint nauseous odour is due. According to Lefort the formula of emetine is $C_{30}H_{44}N_2O_8$. It is a colourless and amorphous alkaloid, melting at 150° F., very soluble in chloroform and alcohol, but slightly in ether and water; dissolves readily in acids, forming soluble salts, but its nitrate is characterised by its great insolubility, and its tannate is still more insoluble. Emetine may be obtained by extracting the powdered root with alcohol, evaporating the spirit, taking up the residue with water slightly acidulated with hydrochloric acid, then adding an excess of potassium hydrate and shaking up with chloroform. According to Reich the formula of this alkaloid should be $C_{20}H_{30}N_2O_5$, but that of Lefort, above given, is more modern. Ipecacuanhic acid is a reddish brown bitter substance, closely allied to kinic

acid, and usually regarded as being a glucoside. Besides the above important principles the root is found to contain small quantities of resin, sugar, gum, pectin, and starch.

Pilula Ipecacuanhæ cum Scillâ. (*Vide* p. 53.)

Pulvis Ipecacuanhæ Compositus (1 in 10). (*Vide* p. 53.)

Trochisci Ipecacuanhæ.

Ipecacuanha, in powder.....	180 grains
Refined sugar, in powder	25 ounces
Gum Acacia, in powder	1 ounce
Mucilage of Gum Acacia	2 fluid ounces
Distilled Water.....q.s.	1 fluid ounce.

Divide into 720 lozenges. ($\frac{1}{4}$ grain in each.)

Dose—I to 3.

Vinum Ipecacuanhæ.

Ipecacuanha, bruised	1 ounce
Sherry	ad. 20 fluid ounces

Macerate 7 days, and filter.

Dose—Expectorant, 5 to 40 minims; emetic, 3 to 6 drachms.
(1 grain of Ipecacuanha in 22 minims.)

Ipecacuanha wine loses power on keeping; much of the emetine being deposited along with potassium bitartrate. An *Acetum Ipecacuanhæ* has been recommended as a more stable preparation.

Ipecacuanha also enters into the composition of the following:—*Pilula Conii Composita* (1 in 6).

Trochisci Morphiæ et Ipecacuanhæ (1-12th grain in each). (*Vide* page 45.)

Substitutions.—1. *Striated Ipecacuanha* is the root of *Psychotria emetica*, and is thicker than the Ipecacuanha, and has marked longitudinal furrows instead of rings on its sides. It is not easily broken but remains moist and tough, and when cut exhibits a dull violet colour.

2. *Undulated Ipecacuanha* is the root of *Richardsonia scabra*. It is rough and knotted, and deep iron-grey in colour. Its bark is thick and brittle, and noticeably light coloured and farinaceous; the medullium is slender, but strong and flexible. It is never marked with rings.

NOTE.—For a ready method of estimating the strength of a sample of Ipecacuanha, see the Author's work on Pharmaceutical Chemistry, page 623.

NATURAL ORDER.—VALERIANACEÆ.

Usually herbs with opposite leaves. Calyx, superior, 4-toothed or pappose; corolla, tubular and epigynous, often gibbous at the base; stamens, 1 to 5, free and epipetalous; ovary, inferior 3-celled, with 2 cells usually abortive; stigma, 2 or 3 lobed; fruit, dry indehiscent; seeds, exalbuminous.

VALERIANÆ RADIX, Valerian Root.—The dried root of *Valeriana officinalis* from indigenous plants, and others cultivated in Britain.

Botanical Source.—The *V. officinalis* is a perennial herb with imparipinnate leaves and lanceolate leaflets; inflorescence a corymbose cyme; calyx persistent; corolla hypocrateriform and gibbous at the base; fruit 1-celled and 1-seeded, crowned by a feathery pappus.


Collection.—The root is derived principally from plants cultivated near Chesterfield, in Derbyshire.

Description.—A short rhizome, yellowish white or brown in colour, with numerous dark brown fibrous roots about 2 or 3 inches long. The odour is penetrating and fetid, and the taste bitter. The rootlets are generally 3 to 4 inches long, and slender, rarely attaining 1-10th of an inch in diameter.

Uses.—A stimulant, anti-spasmodic, and nervine tonic. Useful in hysteria and its allies, chorea and epilepsy, but its action is rather feeble, and the result is not always satisfactory.

Chemistry.—The active principle is a volatile oil, of which the root yields from $\frac{1}{2}$ to 2 per cent. This oil

contains 5 per cent. of valerianic acid, $\text{HC}_5\text{H}_9\text{O}_2$. It also contains 25 per cent. of a hydrocarbon resembling oil of turpentine, called *valerene*, C_5H_8 , as well as *valerol*, $\text{C}_6\text{H}_{10}\text{O}$, and *stearopten*, $\text{C}_{10}\text{H}_{18}\text{O}$, indetical with Dryobalanops camphor. Exposed to the air, valerol is slowly converted into valerianic acid, a reaction which is instantly produced by potassium hydrate.



Besides the volatile oil, Valerian Root contains 6 per cent. of resin, 12 of extractive matter, and a little sugar and malic acid.

Infusum Valerianæ.

Valerian Root, bruised	120 grains
Boiling Distilled Water	10 fluid ounces

Infuse 1 hour.

DOSE—I to 2 ounces.

Tinctura Valerianæ.

Valerian Root, in coarse powder	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

DOSE—I to 2 drachms.

Tinctura Valerianæ Ammoniata.

Valerian Root, in coarse powder	2½ ounces
Aromatic Spirit of Ammonia	ad. 20 fluid ounces

DOSE—½ to 1 drachm.

NATURAL ORDER.—COMPOSITÆ.

Herbs or shrubs with exstipulate leaves. Flowers (florets) in capitula surrounded by an involucre of bracts. Calyx superior, frequently pappose; corolla, tubular or ligulate; stamens, 5 epipetalous, and having syngenesious anthers; ovary, inferior; fruit, a dry indehiscent, 1-celled 1-seeded cypsela (an achæmium, inferior and compound); seed, solitary and exalbuminous.

Of this order, De Candolle makes three sub-orders:—

1. *Tubulifloræ*, with florets of the ray usually ligulate, and either unisexual or neuter, and the florets of the disc tubular and hermaphrodite. Examples:—*Anacyclus*, *Anthemis*, *Arnica*, *Artemisia*, *Carthamus*, *Inula*, *Matricaria* and *Tussilago*.

2. *Labiatifloræ*, with the unisexual florets labiate. No medicinal examples.

3. *Ligulifloræ*, with all florets ligulate and perfect. Examples:—*Cichorium*, *Lactuca* and *Taraxacum*.

PYRETHRI RADIX, Pellitory Root, *Pellitory of Spain*.—The dried root of *Anacyclus Pyrethrum*, imported from the Levant.

Botanical Source.—A low perennial, with small divided leaves and capitulum resembling that of the ox-eye daisy.

Collection.—The root is chiefly collected in Algeria and exported from Tunis *via* Leghorn and Egypt.

Description.—In pieces about the length and thickness of the little finger, cylindrical or tapering off, having a thick brown bark studded with shining points, and only a few hair-like rootlets. Breaks with a resinous fracture, and exhibits internally a radiate structure; has an aromatic smell with a pungent taste, exciting a tingling sensation of the lips and tongue and a flow of saliva. It is very liable to perforations, caused by the attacks of insects.

Use.—As a sialogogue for the relief of tooth-ache.

Chemistry.—It contains a little volatile oil, sugar, gum, tannic acid, and a resinous matter called *pyrethrin*, which has not yet been fully examined.

Tinctura Pyrethri.

Pellitory Root, in coarse powder 4 ounces
Rectified Spiritad. 20 fluid ounces

Macerate 48 hours, and percolate.

(12 grains of Pellitory Root to 1 fluid drachm.)

Substitutions.—In Germany and Russia the root of *Anacyclus officinarum*, which is an annual herb cultivated in Russia, is used instead of the pellitory already described. It is light grey in colour, and only half the thickness of our bark, being besides abundantly covered with the remains of sticks and leaves.

ANTHEMIDIS FLORES, Chamomile Flowers.—The dried single or double flower heads of *Anthemis nobilis*, from plants wild and cultivated in England.

Botanical Source.—The *Anthemis nobilis* is a perennial herb growing not more than a foot high, and much branched and leafy. The leaves are compound bipinnate with linear subulate leaflets. The capitula are terminal and *solitary*, with a convex yellow disc and membranous scales; fruit without a pappus.

Collection.—The flowers are gathered from plants cultivated chiefly at Mitcham, and dried by gentle artificial heat.

Description.—The single flower heads contain both white ligulate and yellow tubular florets, while the double have ligulate and female florets only. The receptacle in both cases is solid, conical, and about a $\frac{1}{4}$ inch high, and covered with narrow chaffy scales. The flowers possess a powerful aroma and have a bitter taste.

Uses.—A stimulant, aromatic tonic. In large doses diaphoretic and even emetic. Useful as a stomachic in dyspepsia and flatulence.

Chemistry.—Chamomiles yield about a half per cent. of essential oil, which when fresh is pale blue, but becomes brownish by keeping. This may be viewed as a mixture of butylic and amylic angelate and valerate. These bodies are easily decomposed by the action of potassium hydrate forming potassium angelate. In practice the oil is distilled from the entire plant. Besides the oil, chamomiles contain a small quantity of a bitter acrid principle, apparently a glucoside.

Extractum Anthemidis.

Chamomile Flowers	16 ounces
Oil of Chamomile	15 minims
Distilled Water	1 gallon

(100 lbs. of flowers yield about 42 lbs. of extract.)

Dose—2 to 10 grains.

Infusum Anthemidis.

Chamomile Flowers	$\frac{1}{2}$ ounce
Boiling Distilled Water	10 fluid ounces

Infuse 15 minutes, in a covered vessel, and strain.

Incompatibles—Salts of iron, mercury, silver, and lead.

Dose—1 to 4 ounces.

Oleum Anthemidis.

The oil distilled in Britain from the flowers.

Substitutions.—1. The flower heads of *Chrysanthemum Parthenium*, or feverfew, which have flat or only slightly convex receptacles and all the florets ligulate. The scales on the receptacle are not membranous.

2. The flower heads of *Matricaria Chamomilla* used in Germany under the name of common chamomile. The flower heads are single; not bitter, and the receptacle is very conical, hollow, and devoid of scales.

ARNICÆ RADIX, Arnica Root.—The dried rhizome and rootlets of *Arnica montana*, collected in mountainous parts of Middle and Southern Europe.

Botanical Source.—*Arnica montana* is a perennial herb with stems about a foot high. The radical leaves are simple, obovate, and entire; the cauline leaves rarely more than two small pairs; inflorescence in bold capitula, and never more than three on one stalk; florets of the ray, yellow, ligulate, and in one whorl only; florets of the disc, tubular, yellow, and hermaphrodite; fruit, cylindrical, with a pappus consisting of one whorl and rigid hairs.

Description. — Slender, contorted; dark brown rhizome, 1 to 3 inches long, and 2 to 3 inches thick, having on its under side an abundance of wiry roots, 3 or 4 inches long, and on its upper side the remains of coriaceous leaves. It has an acrid taste and faint aromatic odour.

Uses. — Externally applied it is feebly irritant; internally, stimulant and diaphoretic. The tincture is used as a popular application to bruises, but its efficacy for this purpose is somewhat hypothetical.

Chemistry.—The root contains about $\frac{1}{2}$ per cent. of volatile oil, having the formula, C_6H_9O , but its active principle is *arnicin*, $C_{20}H_{30}O_4$, a yellow amorphous body with an acrid taste, slightly soluble in water, freely in alcohol and ether, and in alkaline solutions. It is presumed to be a glucoside as it is decomposable by dilute acids, but this has not yet been definitely proved. It may be precipitated from its spirituous solution both by tannic acid and water.

Tinctura Arnicæ.

Arnica Root, in coarse powder..... 1 ounce
Rectified Spiritad. 20 fluid ounces

Macerate 48 hours, then percolate. •

DOSE—1 to 2 drachms. ($2\frac{3}{4}$ grains of Arnica in 1 fluid drachm.)

Substitutions.—The root of *Geum urbanum*, a herb belonging to the Rosaceæ. It is thicker than the rhizome of arnica, and it is, moreover, not a rhizome, but a true root covered on all sides by rootlets.

ARNICÆ FLORES.—The flowers of the same plant, formerly used in making the tincture. They are said to be richer in *arnicin* than the root now employed.

A stronger and better preparation is a tincture made with equal parts of the flowers as well as the root of arnica.

SANTONICA, Wormseed, *Semen-contra*.
—The unexpanded heads of a species of *Artemisia*, imported from Russia.

Botanical Source.—This has lately been described as *Artemisia Cina*, but was formerly considered to be *Artemisia maritima*. It is a low, shrubby, aromatic plant, having for its inflorescence a number of very minute capitula, arranged in a close panicle.

Collection.—The unexpanded flower heads are collected on the Steppes of Kirghiz, in the northern part of Turkestan, and imported *via* Moscow and St. Petersburg.

Description.—In good samples, this drug should consist of entire, unexpanded capitula, very minute in size, and requiring about 90 to weigh one grain. A mixture of stalks and leaves is found in inferior samples. The flower heads are of an oblong form, 1-10th of an

inch long, yellowish when new, and brownish when old. Rubbed in the hand they exhale a powerful odour, somewhat between oil of cajuput and camphor.

Uses.—Santonin is a vermicide, being active against the round and thread worms, but not against *tænia*. The dose is 2 to 6 grains in powder, or dissolved in oil and borax.

Chemistry.—Santonica yields about 1 per cent. of volatile oil, and an active principle called *santonin*, which is a glucoside having the formula $C_{15}H_{18}O_3$.

Santonin B.P. is prepared by boiling 1 lb. of santonica with 1 gallon of water and 5 ounces of slaked lime for $\frac{1}{2}$ hour; strain and boil the flowers again with $\frac{1}{2}$ gallon of water and 2 ounces of slaked lime for $\frac{1}{2}$ hour; strain, mix the two liquors, filter and evaporate to $2\frac{1}{2}$ pints. Add to the hot liquid, hydrochloric acid in slight excess, and set aside for 5 days. Remove any floating oily matter, decant the fluid, and collect the precipitate on a filter and wash it with cold water until it is free from acid, and then with a mixture of $\frac{1}{2}$ fluid ounce of solution of ammonia and 5 ounces of water, and lastly with cold water, until the washings pass colourless. Press and dry the residuc. Boil for 10 minutes with 60 grains of purified

animal charcoal, and 9 ounces of rectified spirit. Filter : wash the charcoal with an ounce of boiling spirit, and set aside for two days to crystallise. Protect the crystals from the light. In this process the santonin is extracted as a soluble calcium santonate ; this is decomposed and the santonin precipitated by hydrochloric acid, the resinous matters being washed away by the ammonia water. It is often re-dissolved in boiling spirit decolorised by animal charcoal, and on cooling the santonin crystallises.

Characters and Tests. — In colourless tubular crystals, slightly bitter, fusible and subliming at a gentle heat. It is dimorphous, and on exposure to light it undergoes a crystalline change, which renders it yellow. It is very slightly soluble in cold and even boiling water, but readily soluble in chloroform and boiling rectified spirit. *It is insoluble in dilute acids.* Santonin has been viewed as an anhydride of santonic acid, $C_{15}H_{20}O_4$.

INULÆ RADIX, Elecampane Root.
—The dried root of *Inula Helenium*, from Central and Southern Europe, cultivated in England.

Botanical Source.—It is a perennial herb with large ovate-cordate, acute leaves, downy on the under surface; inflorescence, large bright yellow capitulum, having small florets of the ray, and hermaphrodite florets of the disc.

Collection.—The roots should be collected from plants not younger than two, nor older than three years. They are sometimes dried entire, but when large they are variously sliced.

Uses.—Aromatic tonic, diaphoretic and expectorant.

Chemistry.—Elecampane contains 36 per cent. of bitter extractive, and a peculiar kind of starch termed inulin. This body has the same composition as starch, $C_6H_{10}O_5$, and is usually found in the roots of the Compositæ. It is a white tasteless powder, turned *yellow* on the addition of tincture of iodine. It is almost insoluble in cold but readily soluble in boiling water, from which it partly separates again on cooling. With dilute acids it is converted into dextrin and sugar, and by the action of nitric acid it is converted into oxalic and acetic acids. It may be precipitated from its solution by alcohol. Elecampane root also contains a neutral crystalline principle called *helenin*.

Not official, but made into a decoction of $\frac{3}{4}$ ounce to the pint.

Dose, 1 to 2 ounces.

TARAXACI RADIX.—The fresh and dried roots of *Taraxacum Dens Leonis*, from meadows and pastures in Britain and elsewhere.

Botanical Source.—*Taraxacum officinale*, *Leontodon Taraxacum*, or *Taraxacum Dens Leonis*, is a herb with a milky juice; radical, pinnatifid, and runcinate leaves: inflorescence a bold yellow capitulum, supported on a perfectly glabrous scape, well known as the common dandelion.

Collection.—The roots are to be dug up, according to the British Pharmacopœia, between September and February, but they are generally considered to be in perfection for making extract in the month of November. Some discrepancy of opinion however exists upon this point.

Description.—The dried root is dark brown, with shrivelled wrinkles running longitudinally often in a spiral manner. It is $\frac{1}{2}$ an inch or less in thickness, and breaks easily with a short corky fracture, showing

a thick white cortical portion containing a white medullium, which is very porous, without rays. It is inodorous and bitter in taste. It is very liable to the attacks of maggots.

Uses.—Employed as a mild laxative in liver disorders, and it is popularly supposed to be diuretic in action. Its roasted root is sometimes used in the adulteration of coffee.

Chemistry.—The dried root contains 24 per cent. of inulin, and about 20 per cent. of sugar. It also contains an acrid resin, a colourless amorphous bitter substance called *taraxacin*, which may be obtained by boiling the milky juice of the fresh root with water and evaporating, when it deposits warty crystals. The fresh root also contains a substance called *taraxacerin*, $C_8H_{16}O$, a substance resembling *lactucerin*. The leaves and stocks of dandelion yield a peculiar sugar called *inosite*, $C_6H_{12}O_6 \cdot 2H_2O$.

Decoctum Taraxaci.

Dried Dandelion Root, sliced and bruised 1 ounce
Distilled Water 1 pint

Boil 10 minutes; make up product to 20 fluid ounces.

Dose—2 to 4 ounces.

Extractum Taraxaci.

The expressed juice of fresh Dandelion Root, evaporated at 160° to an extract, after the feculence has deposited.

100 lbs. of root yield 25 lbs. dried ; 30 lbs. of juice, or 8 lbs. of extract.

Dose—5 to 30 grains.

Succus Taraxaci.

Dandelion Juice 3 pints

Rectified Spirit..... 1 pint

Filter after standing 7 days.

Dose—1 drachm to 2 drachms. (2 drachms are equal to $\frac{1}{2}$ drachm of extract, or 6 drachms of fresh root.)

Substitutions.—The roots of *Leontodon hispidus* (common hawkbit). The leaves of the hawkbit are pilose, while those of the dandelion are glabrous. The flower stalk of the hawkbit is branched and the fresh root does not show any quantity of milky juice.

LACTUCA VIROSA.—The dried flowering herb of *Lactuca virosa*, indigenous.

Botanical Source.—It is a biennial herb with an acrid milky juice. Radical leaves, obovate and undivided, with a prickly keel, but otherwise nearly smooth, and finely dented ; cauline leaves, often auricled and half amplexicaul ; inflorescence, a panicle

of capitula; flowers, pale yellow. The plant has a rank smell, and red spots on the stem.

Uses.—A soporific, but not active.

Chemistry.—The plant when wounded yields a white milky juice, which when exposed to the air rapidly hardens to small yellow-brown masses, which are known as *lactucarium* or lettuce-opium. This substance, which is not official, is specially collected in Rhenish Prussia, by cutting the stem just before the plant flowers, and collecting the juice from the wounded top by the finger and transferring it to earthen cups. Formerly the *annual* plant was cultivated near Edinburgh and the juice collected in tin vessels. *Lactucarium* contains several substances of which the following are the principal:—

1. *Lactucerin*, $C_{16}H_{26}O$, this may be obtained by extraction with boiling alcohol in the form of colourless needles insoluble in water, but soluble in ether and in oils. Good *lactucarium* yields 59 per cent. of this body.

2. *Lactucin*, $C_{11}H_{12}O_3 \cdot H_2O$, in pearly scales, soluble in acetic acid but insoluble in ether.

In addition to those we find albumen, gum, oxalic,

citric, malic and succinic acids, sugar, mannite and metallic solids.

Extractum Lactucæ.

The expressed juice of the flowering herb of lettuce, evaporated at 160° to an extract.

(100 lbs. of the plant yield 52 lbs. of juice, or 5¼ lbs. of extract.)

Dose—5 to 15 grains.

NATURAL ORDER.—LOBELIACEÆ.

Herbs or shrubs with a milky juice; corolla irregular; stamens, 5 syngenesious; ovary inferior, 3-celled; fruit capsular.

LOBELIA.—The dried flowering herb of *Lobelia inflata* from North America, from Canada to Carolina.

Botanical Source.—The *Lobelia inflata* is an annual herb, growing along the roadsides in the eastern part of North America.

Collection.—The herb is collected by the sect called Shakers in August and September; dried, and chopped before it is quite dry, and neatly trimmed in the form of rectangular cakes. The drying is then finished, and the cakes are usually wrapped in blue paper.

Description.—As met with dry, lobelia has an angular stem, alternate ovate and dented leaves, hairy beneath. The fruits in racemes, consisting of membranous capsules, ovoid in shape, inflated and 10-ribbed. Inside the fruits are numerous minute seeds about 1-50th of an inch long, dark brown in colour, ovate-oblong in shape, with a reticulated testa. The characters of the seeds under a microscope form an excellent guide to recognition.

Uses. — A powerful anti-spasmodic and depressive remedy, and useful in spasmodic asthma and tetanus. It produces nausea, coupled with vomiting, cold sweats, muscular relaxations, and causes death by collapse and finally coma.

Chemistry.—The active principle is due to a volatile alkaloid of an oily consistence, smelling like tobacco. It is called *lobelina*, and is soluble in water and more so in alcohol and ether. It may be isolated by making a decoction of the plant with acidulated water, precipitating with sodium phospho-molybdate, and distilling the precipitate with baryta water. It forms crystallisable salts with acids. Lobelia also contains a little volatile oil, resin and gum, and an acrid principle called

lobelacrin, which may be isolated by making a tincture and distilling it with animal charcoal. The charcoal retains the acrid principle, and after washing with water it is distilled out by boiling spirit. It crystallises in warty tufts, and is freely soluble in ether and chloroform.

Tinctura Lobeliæ.

Lobelia, in coarse powder 2½ ounces
 Proof Spiritad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—10 to 30 minims. (7 grains of Lobelia in 1 fluid drachm.)

Antidotes—Internal and external stimulants.

Tinctura Lobeliæ Ætherea.

Lobelia, in coarse powder 2½ ounces
 Spirit of Etherad. 20 fluid ounces

Macerate 7 days, then strain, press, and filter.

Dose—10 to 30 minims. (7 grains of Lobelia in 1 fluid drachm.)

SUB-DIVISION II.—HYPOSTAMINEÆ.

NATURAL ORDER.—ERICACEÆ.

Evergreen shrubby plants; leaves opposite or whorled. exstipulate; calyx, persistent; corolla, monopetalous and hypogynous; stamens, equal in number to or double

the divisions of the corolla; anthers, appendiculate; ovary, superior; placentation axile; fruit, capsular; seeds, numerous and albuminous.

UVÆ URSI FOLIA, Bearberry Leaves

—The dried leaves of *Arctostaphylos Uva Ursi* from Northern Europe, North Britain, and North America.

Botanical Source.—The bearberry is a small ever-green shrub bearing terminal clusters of flowers and scarlet globose berries.

Collection.—The leaves are taken in September and October usually from indigenous plants.

Description.—Small dark green coriaceous and glabrous leaves, not exceeding 1 inch in length by 3-8th of an inch in breadth, obovate in shape, with short petiole. The upper surface is shining and deeply impressed with a network of veins and much reticulated beneath.

Uses.—As a simple astringent in diarrhœa; also in some affections of the bladder and kidneys.

Chemistry.—*Uva Ursi* leaves contain 36 per cent. of tannic acid and $1\frac{1}{2}$ per cent. of gallic acid. They

also yield a bitter neutral substance called *arbutin*, $C_{24}H_{32}O_{14} \cdot H_2O$. This is very soluble in water and alcohol, but only slightly so in ether. It may be obtained by treating a decoction with basic acetate of lead, filtering out the precipitated tannate and gallate of lead, removing the excess of lead from the filtrate by sulphuretted hydrogen and then concentrating and crystallising the subsidiary glucoside, convertible by boiling with a dilute acid into hydro-quinone, $C_6H_6O_2$, and glucose. By oxidation it is convertible into *quinone*, $C_6H_4O_2$, which when heated evolves its characteristic odour and a solution stains the skin yellow. Another bitter amorphous substance called *ericolin*, may be obtained from the mother liquor after the crystallisation of the arbutin. Its formula is $C_{34}H_{56}O_{21}$. As the leaves contain tannic acid the decoction may be precipitated by salts of iron.

Infusum Uvæ Ursi.

Bearberry Leaves, bruised.....	$\frac{1}{2}$ ounce
Boiling Distilled Water.....	10 fluid ounces

Infuse in a covered vessel for 2 hours, and strain.

Incompatibles—Alkalies and their carbonates; lime water.
and most metallic salts; Decoctum Cinchonæ.

Dose—1 to 2 ounces.

Substitutions.—The leaves of the red whortleberry, *Vaccinium Vitis-Idoea*, which are dotted on the under surface, and crenate towards the apex; also the leaves of the common box (*Buxus sempervirens*).

SUB-DIVISION III.—EPIPETALÆ.

NATURAL ORDER.—SAPOTACEÆ.

GUTTA-PERCHA.—The concrete juice of *Isonandra gutta* from the East Indian Archipelago.

Botanical Source.—The *I. gutta* is a bold ever-green tree, a native of Borneo, Sumatra and Singapore, at the latter of which it is now extinct, owing to the wasteful manner of its collection.

Collection.—The tree is simply felled by the Malays and the juice collected in cocoa-nut shells and then pressed into blocks of about 5 lbs. weight each. When imported to this country it is softened in hot water, torn into shreds by machinery, and then while still soft pressed together again.

Description.—In light brown or chocolate flexible

but inelastic pieces. Insoluble in water, alcohol, dilute acids and alkalies or fixed oils, but soluble in ether, benzol, chloroform, carbon disulphide and volatile oils. Sp. gr. .979.

Uses.—Employed in making the official preparation *Charta Sinapis*.

Chemistry.—Gutta-percha softens at 113° F., and at 212° can be easily moulded, and is very adhesive when heated dry. It is a non-conductor of electricity and becomes negatively electrified when rubbed. Crude gutta contains about 75 to 82 per cent. of a terpene called *caoutchin*, $C_{20}H_{32}$, which is solid, white, soluble in ether, benzol and chloroform, but insoluble in alcohol and only fusible at 300° F. The remainder is built up of two resins: viz., a white one having the formula $C_{10}H_{16}O$ or $C_{20}H_{32}O_2$, which is crystalline; and a yellow one $C_{20}H_{32}O$, which is amorphous. Both are soluble in boiling alcohol. Crude gutta by exposure to the light gradually oxidises to these resins and becomes brittle, but keeps well under water in the dark.

Liquor Gutta Percha.

Gutta Percha, in thin slices	1 ounce
Chloroform	8 fluid ounces
Carbonate of Lead, in fine powder	1 ounce

Dissolve the Gutta-percha in 6 ounces of Chloroform in a stoppered bottle, add the Carbonate of Lead, previously mixed with the remaining Chloroform, and after shaking several times, allow the insoluble matter to subside. Lastly, decant the clear liquid, and keep it in a stoppered bottle.

The action of the plumbic carbonate is merely mechanical to separate the small portion of flocculent matter insoluble in the chloroform.

NATURAL ORDER.—STYRACEÆ.

BENZOINUM, Benzoin, *Gum Benjamin*.—

A balsamic resin from the stem of *Styrax Benzoin*, imported from Siam and Sumatra.

Botanical Source.—*Styrax Benzoin* is a tree with a thick stem, indigenous to Java and Sumatra.

Collection. — The resin is obtained by making incisions in the trees, when seven years old. It exudes and is scraped off with a knife. This process is carried on for about ten years, when the tree is cut down, and an inferior variety is obtained by splitting and scraping. Siamese benzoin is obtained by cutting the bark and allowing the resin to concrete between it and the wood and then stripping the tree.

Description.—Siamese benzoin consists of a mass of flattened tears of an opaque milk-white resin, loosely agglutinated. It is very brittle, and softens in the mouth. It fuses at 167° and evolves fumes of benzoic acid. Sumatra benzoin is in cubic blocks of greyish tint, containing opaque tears of a greyish resin, mixed with fragments of wood. In inferior samples the tears are almost entirely absent. Its odour is more feeble than the Siamese. It melts between 185° and 203° , and is much inferior in value to the Siamese.

Uses.—Chiefly for making incense. It acts medicinally as an active stimulant. It also lessens the tendency of fats to become rancid.

Chemistry.—The active principle of benzoin is benzoic acid, which exists to the extent of from 14 to 20 per cent. It may be obtained by sublimation and also by boiling with calcium hydrate and water, and precipitating benzoic acid by concentrating the solution, adding hydrochloric acid. An alcoholic solution of benzoin is coloured a dark brownish green by the addition of ferric chloride, but a watery decoction is unaffected. Cold, strong sulphuric acid dissolves benzoin, producing a beautiful carmine colour, and on

the addition of water characteristic crystals of benzoic acid separate.

Adeps Benzoatus.

10 grains to 1 ounce. Used in the preparation of the four B.P. suppositories, and some pharmacopœial ointments.

Tinctura Benzoini Composita. (*Friar's Balsam.*)

Benzoin, in coarse powder.....	2 ounces
Prepared Storax	1½ ounce
Balsam of Tolu	½ ounce
Socotrine Aloes	160 grains
Rectified Spirit	ad. 20 fluid ounces

Macerate 7 days, then filter.

Dose—½ to 1 drachm, triturated with mucilage. As a lotion, 1 drachm to 5 ounces of rose water, adding the water to the tincture.

Specially Distinctive Tests.—*Siamese* benzoin is—

1. Completely soluble in chloroform, but imperfectly in ether.

2. Extracted by petroleum spirit (boiling under 100° F.) and the filtered solution evaporated, a residue is obtained which is coloured cherry-red by sulphuric acid.

3. Heated with sulphuric acid and potassium dichromate, it evolves no odour of benzoic aldehyd.

Sumatra benzoin sometimes does give a little odour with the last test.

Acidum Benzoicum, B.P. ($\text{HC}_7\text{H}_5\text{O}_2$), is prepared as already stated, by sublimation or by the wet process with lime. It is also manufactured artificially from *naphthalin*, C_{10}H_8 , which is treated with nitric acid, and thus converted into *phthalic acid*, $\text{H}_2\text{C}_8\text{H}_4\text{O}_4$. By heating with calcium hydrate to about 650°F . the calcium phthalate formed is converted into calcium carbonate and calcium benzoate, from which latter the acid is liberated by the action of hydrochloric acid.

Benzoic acid is in light feathery crystals, having usually a slight odour of benzoin. It is slightly soluble in water (1 in 200) and more so in boiling water, and readily in rectified spirit and alkalies, forming salts which are freely soluble. It melts at 248°F ., boils at 462°F ., and then volatilises. It gives a flesh-colour precipitate with ferric chloride, which when filtered and washed, and treated in the filter with diluted ammonia, yields a solution of ammonium benzoate, which deposits crystals of benzoic acid on the addition of hydrochloric acid.

The dose is 10 to 30 grains.

Benzoic acid enters into the composition of *Tinctura Camphoræ Composita*, *Tinctura Opii Ammoniata*, and is used in the preparation of Ammonium Benzoate.

NATURAL ORDER.—LOGANIACEÆ.

NUX VOMICA, Poison Nut.—The seeds of *Strychnos Nux vomica*, from Coromandel and other parts of India.

Botanical Source.—The *Strychnos Nux vomica* is a moderately sized tree, indigenous especially to the Coast districts of India.

Description.—It is a greyish disc-like seed, covered with very short satiny hairs about 1 inch in diameter and a $\frac{1}{4}$ inch in thickness. It is convex on the ventral surface and slightly concave on the dorsal. In the centre there is a scar, which is the hilum, from which the raphè runs as a faintly projecting line to a small protuberance on the edge.

Uses.—Owing to the presence of strychnia, it is, like that alkaloid, a nervine tonic, only in a smaller degree; but, from the variation in the amount of alkaloids present in different samples, its preparations are uncertain in action compared with that of *liquor strychniæ*.

Chemistry. — The seeds contain two principal alkaloids—*Strychnia* ($\frac{1}{4}$ to $\frac{1}{2}$ per cent.), and *brucia* 2 1-10ths to 1 per cent.), and one of lesser importance called *igasuria*.

Strychnia, $C_{21}H_{22}N_2O_2$, is obtained, according to the B.P. by subjecting 1 lb. of nux vomica for two hours to steam, then drying rapidly and grinding in a coffee mill. Digest the powder at a gentle heat for twelve hours with two pints of rectified spirit and one of water: strain through linen, express strongly, and repeat the process twice. Distil off the spirit from the mixed fluids; evaporate the watery residue to about 16 ounces and filter when cold. Add now 180 grains of acetate of lead previously dissolved in distilled water, so long as it occasions any precipitate; filter; wash the precipitate with 10 ounces of cold water, adding the washings to the filtrate; evaporate the clear liquid to 8 ounces, and when it has cooled add solution of ammonia in slight excess, stirring freely. Let the mixture stand at the ordinary temperature for twelve hours; collect the precipitate on a filter, wash it once with a few ounces of cold distilled water, dry it in the vapour bath and boil it with successive portions of rectified spirit till the fluid scarcely tastes bitter. Distil off most of the spirit,

evaporate the residue to the bulk of about $\frac{1}{2}$ ounce, and set it aside to cool. Cautiously pour off the yellowish mother liquor (which contains the brucia and igasuria) from the white crust of strychnia which adheres to the vessel. Throw the crust on a paper filter, wash it with a mixture of two parts of rectified spirit and one of water, till the washings cease to become red on the addition of nitric acid (absence of brucia); finally dissolve it by boiling it with an ounce of rectified spirit and set aside to crystallise. More crystals may be obtained by evaporating the mother liquor.

The alcohol extracts the alkaloids as igasurates. The addition of plumbic acetate forms plumbic igasurate and liberates acetates of strychnia and brucia. The ammonia precipitates both alkaloids, which are afterwards separated by fractional crystallisation from boiling rectified spirit, the strychnia crystallising out and leaving the brucia in solution.

Strychnia is in rectangular prisms, colourless and inodorous, slightly soluble in cold water (1 in 6700) more so in boiling water (1 in 2500). It is soluble in boiling rectified spirit, but not in absolute alcohol or ether, and it is very soluble in chloroform. Fragments of strychnia dissolve in cold strong sulphuric acid

without colour, but the addition of a crystal of potassium dichromate causes a beautiful evanescent violet. A similar effect may be produced by the use of potassium ferricyanide, manganese peroxide and several other re-agents, but the potassium dichromate is the one usually employed.

In doses of 1-36th of a grain, increased under certain circumstances to 1-12th, strychnia is a powerful nervine tonic. In larger doses it poisons by affecting the whole of the contractile muscular tissue and producing tetanic convulsions, with well marked *opisthotonos*, and finally killing by suffocation.

Antidotes.—Tobacco enemata, chloroform, chloral hydrate, Indian hemp, sal. volatile, animal charcoal.

Brucia, $C_{23}H_{26}N_2O_4 \cdot 4H_2O$, may be obtained by crystallisation from the alcoholic mother liquors left after the preparation of strychnia; it crystallises in colourless oblique four-sided prisms or laminæ. It is soluble in cold water (1 in 850), and in boiling water (1 in 150), also freely in chloroform, but almost insoluble in ether. It is freely soluble in alcohol. Its salts are bitter and have an action similar to strychnia,

but less powerful. Treated with nitric acid it strikes a red colour, which when heated turns yellowish, and stannous chloride added to the solution causes it to turn violet. Brucia gives a dirty orange colour precipitate with potassium iodide.

Extractum Nucis Vomicae.

Nux Vomica..... 16 ounces

Rectified Spirit q.s.

Dose— $\frac{1}{2}$ to 2 grains, (16 ounces yield 1 ounce of extract.)

Liquor Strychniæ.

Strychnia, in crystals 4 grains

Dilute Hydrochloric Acid 6 minims

Rectified Spirit..... 2 fluid drachms

Distilled Water 6 fluid drachms

(Contains $\frac{1}{2}$ grain of Strychnia in 1 fluid drachm.)

Dose—5 to 10 minims.

Tinctura Nucis Vomicae.

Nux Vomica..... 2 ounces

Rectified Spiritad. 20 fluid ounces

Soften by steam, dry and powder. Macerate 48 hours,
then percolate. (2-5ths the strength of P.D.)

Dose—10 to 20 minims. (1 grain of Nux Vomica in 11 minims.)

Antidotes—Tobacco enemata, chloroform, sal. volatile, animal charcoal.

Strychnia being a cumulative remedy requires to be watched with care. The use of *liquor strychniæ* is

much to be preferred to the exhibition of the alkaloid in the solid state.

IGNATII SEMINA, St. Ignatius's Beans.—The seeds of *Strychnos Ignatia*, from the Philippine Islands. Not official.

Description.—Dark coloured beans about 1 inch in length, ovoid in form and very irregular in shape owing to pressure, with well marked hilum at one end and slightly hairy on the surface.

Uses. — For the manufacture of strychnia. An extract of the seeds has likewise been introduced into medicine.

Chemistry.—The seeds contain 1·5 per cent. of strychnia and ·5 of brucia.

SPIGELIÆ RADIX, Carolina or Indian Pink Root.—The dried rhizome, with rootlets and portions of the herb attached, of *Spigelia Marilandica*, from southern parts of North America. Not official.

Botanical Source.—It is a herb with a perennial and quadrangular stem, and sessile, lanceolate and decussate leaves. The flowers are sessile and in spikes.

Description.—A short knotty dark brown rhizome with knotty roots, having portions of the quadrangular stem attached, sometimes almost the entire plant. It must be carefully distinguished from Serpentry Root to which it bears a resemblance.

Uses. — Anthelmintic. In large doses an acrid narcotic poison.

Chemistry.—It has not been properly examined, but appears to contain both a fixed and volatile oil, and a bitter extractive matter, which is possibly the active principle.

NATURAL ORDER.—GENTIANACEÆ.

Smooth herbs with opposite, entire, sessile and ribbed leaves; flowers regular; calyx and corolla persistent; stamens equal in number to, and alternate with, the lobes of the corolla; ovary superior 1-celled, with 2 parietal placentæ meeting in the centre; fruit, usually a capsule with septicidal dehiscence; seeds, numerous and albuminous.

GENTIANÆ RADIX, Gentian. — The dried root of *Gentiana lutea*, collected in the mountainous districts of Central and Southern Europe.

Botanical Source.—The *Gentiana lutea* is a herb growing 2 to 3 feet high, with a perennial root, and having opposite sessile, ovate, and ribbed leaves: inflorescence, verticillasters of dark bluish flowers, with irregular corolla.

Description.—The twisted root from $\frac{1}{2}$ to 1 inch in thickness, brownish externally, yellowish within. The roots are much wrinkled longitudinally, and marked transversely with many rings.

Uses.—A pure simple bitter tonic.

Chemistry.—The active principle of gentian is called *gentiopicrin*, $C_{20}H_{30}O_{12}$, of which the root contains about .1 per cent. This substance is a neutral, crystallisable glucoside, soluble in water and spirit. By boiling with a dilute acid it is converted into *gentiogenin* and glucose. It forms a yellowish liquid with potassium iodide, which rapidly loses its bitterness. It is insoluble in ether. Gentian also

contains *gentianic* acid, $C_{14}H_{10}O_5$, which is in pale yellow silky crystals, tasteless and volatile without decomposition. This constituent is inert. Besides these we find in gentian, sugar, pectin, and oil, but no starch.

Extractum Gentianæ.

Gentian Root	16 ounces
Boiling Distilled Water	1 gallon

(100 lbs. of root yield about 50 lbs. of extract.)

Dose—2 to 10 grains.

Infusum Gentianæ Compositum.

Gentian Root, sliced	1 drachm
Bitter Orange Peel, cut small	1 drachm
Fresh Lemon Peel, ,,	$\frac{1}{4}$ ounce
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel 1 hour, and strain.

Incompatibles—Salts of iron, lead, and analogous salts.

Dose—1 to 2 ounces.

Mistura Gentianæ.

Gentian Root, sliced	$\frac{1}{4}$ ounce
Bitter Orange Peel, cut small	30 grains
Coriander Fruit, bruised	30 grains
Proof Spirit	2 fluid ounces
Distilled Water	8 fluid ounces

Dose— $\frac{1}{2}$ to 1 ounce. (Infusion, 1 in 80; mixture, 1 in 40; tincture, 1 in 13 $\frac{1}{2}$.)

Tinctura Gentianæ Composita.

Gentian Root, bruised	1½ ounce
Bitter Orange Peel, cut small	$\frac{3}{4}$ ounce
Cardamom Seeds, freed from the pericarps and bruised	$\frac{1}{4}$ ounce
Proof Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE— $\frac{1}{2}$ to 2 drachms. (4 grains of Gentian Root in 1 fluid drachm.)

Substitutions.—Owing to careless collection, gentian frequently contains the root of *Gentiana purpurea*, which is somewhat thicker having a peculiar branched appearance at the top never seen in gentian. Two rarer mixtures of the roots are *Gentiana punctata* and *G. Pannonica*, the root much resembling the *purpurea* but not so thick.

CHIRATA, Chiretta (variously spelt).—The entire plant of *Ophelia Chirata*, collected in Northern India.

Botanical Source.—It is an annual herb with orange brown stem and a tapering root. The stem is cylindrical beneath, but bluntly quadrangular towards its apex, having a thin bark and very large pith. The

leaves are entire, sessile, ovate-acuminate, opposite and decussate; inflorescence, a paniced cyme of numerous flowers with two bracts at each division. The corolla persistent and rotate, and the capsules 1-celled, but bi-valved with numerous seeds.

Collection.—The entire plant is pulled up when in flower, with some of the lower fruits beginning to ripen, and is tied into bundles weighing about 2 lbs.

Uses.—A pure bitter tonic, more powerful than gentian.

Chemistry.—Chirata contains two bitter principles: *Ophelic* acid $C_{13}H_{20}O_{10}$, and *Chiratin*, $C_{26}H_{48}O_{15}$, the former being the most abundant. *Ophelic* acid is an amorphous, yellowish, viscid substance, with a bitter taste and faint odour. It is soluble in water, alcohol, and ether. Its watery solution reduces Fehling's solution of copper, and may be precipitated by basic plumbic acid, but not by tannic acid. *Chiratin* is a neutral, crystalline, pale yellow glucoside, soluble in water, alcohol and ether, and is precipitated by tannic acid.

Infusum Chiratæ.

Chiretta, cut small	$\frac{1}{4}$ ounce
Distilled Water at 120°.....	10 fluid ounces

Infuse in a covered vessel $\frac{1}{2}$ hour, and strain.

Incompatibles—Most metallic salts.

Dose—1 to 2 ounces.

Tinctura Chiratæ.

Chiretta, cut small, bruised	2 $\frac{1}{2}$ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

Dose— $\frac{1}{2}$ to 2 drachms. (7 grains of Chiretta in 1 fluid drachm.)

Substitutions.—The chiretta imported up to the present has been found perfectly pure. There are, however, two British plants belonging to the same order which may be used as substitutes, and which indeed have been employed for many years by the herbalist. They are :—

1. *Menyanthes trifoliata*, or Buckbean, sometimes called Marsh trefoil. It possesses very characteristic large ternate leaves, with long petioles.

2. *Erythræa Centaurium*, or small centaury.

Both these plants are pure bitters.

NATURAL ORDER.—ASCLEPIADACEÆ.

HEMIDESMI RADIX,

Hemidesmus Root, *Indian Sarsaparilla*.—

The dried root of *Hemidesmus Indicus*, from India.

Botanical Source.—The *Hemidesmus Indicus* is a twining shrub growing both in India and Ceylon.

Description.—It is a brownish cylindrical tortuous root, longitudinally furrowed, from .2 to .7 of an inch in thickness, with a few thin rootlets and a number of annular cracks, having an odour resembling that of the tonquin bean.

Uses.—Tonic, alterative, diuretic. Employed as a substitute for sarsaparilla.

Chemistry.—Hemidesmus has not been satisfactorily examined.

Syrupus Hemidesmi.

Hemidesmus Root, bruised	4 ounces
Refined Sugar	28 ounces
Distilled Water	20 fluid ounces

Product, 2 lbs. 10 ounces by weight ; or 2 lbs. by measure.

Dose—1 drachm. Sp. gr. 1.335.

NATURAL ORDER.—CONVOLVULACEÆ.

Herbs or shrubs with a twining habit, and alternate exstipulate leaves ; calyx much imbricated and persistent ; corolla 5-plaited and deciduous ; stamens 5 alternate with the lobes of the corolla, and rising from near its base ; ovary superior ; fruit capsular 1—4-celled with septifragal dehiscence.

JALAPA, Jalap. —The dried tubercules of *Exogonium purga*, imported from Mexico.

Botanical Source.—Jalap is a twining plant with a tuberous root, deeply auricled cordate leaves and fine pink flowers, which grows wild on the Eastern slopes of the Andes.

Collection.—The root is dug up at any time. The larger tubercules are gashed to assist the drying. They are then placed in a net inside the house until quite dry.

Description.—Ovoid tubercules, pointed at one end, very dense and hard, dark brown and wrinkled. They are frequently met with gashed or cut into segments.

They form a light greyish brown powder, with a peculiar odour and a sickly taste.

Uses.—A drastic purgative, producing copious watery stools, not followed by after-irritation. As a rule, jalap does not gripe and is less irritant than either gamboge or podophyllin. It is useful for the removal of round and thread worms. Dose, of the powdered root, 10 to 20 grains ; of the resin, 1 to 3 grains.

Chemistry.—Jalap root contains about 12 to 18 per cent. of resin, which is the active principle, together with sugar, gum and cellulose.

Jalapæ Resina, resin of jalap, is prepared officially as follows :—Digest 8 ounces of jalap in coarse powder with 16 fluid ounces of rectified spirit in a covered vessel, at a gentle heat for 24 hours ; then transfer to a percolator, and when the tincture ceases to pass, pour into the percolator successive portions of rectified spirit, until the jalap is exhausted. Add to the tincture 4 fluid ounces of water, and distil off the spirit by the water-bath. Remove the residue while hot to an open dish and allow it to become cold. Pour off the supernatant fluid from the resin, wash it two or three times with hot

water, and dry it in a porcelain plate by the heat of a stove or water bath.

As thus prepared the resin is in dark brown, opaque fragments, brittle, breaking with a resinous fracture and easily reduced to a pale brown powder. It is freely soluble in rectified spirit, but only very slightly soluble in ether, and insoluble in oil and turpentine. The portion of the official resin insoluble in ether consists of *convolvulin* $C_{31}H_{50}O_{16}$. It melts at $300^{\circ}F.$, and is colourless, and insoluble in ammonia. It is soluble in caustic potash and soda, which converts it into convolvulic acid. When oxidised by nitric acid, it is converted into oxalic and ipomœic acids. The portion of the official resin which is soluble in ether is called jalapin ($C_{34}H_{56}O_{16}$), amounts to about 7 per cent. and appears to be active.

Jalap resin enters into the composition of *Pilula Scammonii Composita*.

Extractum Jalapæ.

Jalap, in coarse powder	16 ounces
Rectified Spirit	4 pints
Distilled Water	1 gallon

(100 lbs. of Root yield 50 lbs. of extract.)

DOSE—5 to 15 grains.

Pulvis Jalapæ Compositus.

Jalap, in powder	5 ounces
Acid Tartrate of Potash.....	9 ounces
Ginger, in powder	1 ounce
DOSE—20 to 60 grains. (10 grains of Jalap in 30 grains.)	

Tinctura Jalapæ.

Jalap, in coarse powder	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—½ to 2 drachms. (7 grains of Jalap in 1 fluid drachm.)

Jalap enters into the composition of Pulvis Scammonii Compositus.

Adulterations.—1. The root of *Ipomœa Orizabensis* or Mexican male jalap. Is usually found in spindle-shape pieces of a lighter colour than true jalap, and very deeply wrinkled. It is compact, but not so heavy as true jalap. When broken it exhibits a radiate section, and a number of woody fibres project from the broken surface. The resin yielded by this root is soluble in ether, and apparently identical with the jalapin of scammony resin. It is similar in re-actions and exists to the extent of about 12 per cent.

2. The root of *Ipomœa simulans*, or Tampico jalap, is very similar to true jalap, but more elongated and shrivelled, and destitute of the transverse scars found on the true tubercules. It yields 10 per cent. of a resin entirely insoluble in ether, and, therefore, probably *convolvulin*.

SCAMMONIÆ RADIX, Scammony Root.—The dried root of *Convolvulus Scammonia*, imported from Syria and Asia Minor.

Botanical Source.—The *Convolvulus Scammonia* is a twining plant, much resembling the wild convolvulus which grows in waste places in Syria and Asia Minor.

Description.—Long tough roots, somewhat like a carrot in shape, and spirally twisted; sometimes attaining 3 inches in diameter at the crown. It is greyish brown without and very pale brown internally with a faint odour.

Chemistry.—The root yields from 5 to 5½ per cent. of scammony resin, besides gum, starch, sugar and extractive matter.

Scammoniæ Resina (B.P.), resin of Scammony, is

prepared, according to the B.P., in a manner similar to that described for the manufacture of jalap resin. It is in dark brown fragments, translucent at the edges, breaking with a resinous fracture, and it is completely soluble in alcohol. It is also entirely soluble in ether, and does not form an emulsion when rubbed with water (a distinction from *scammonium*). It should be tested for possible adulteration with guaiacum resin by placing a drop of its tincture on a freshly cut potato peeling, which ought not to turn blue.

Extractum Colocynthis Compositum. (*Vide* p. 167.)

Mistura Scammonii.

Resin of Scammony 4 grains

Milk 2 fluid ounces

Dose— $\frac{1}{2}$ to 2 ounces (for a child). Virgin Scammony makes a better emulsion (*Squire*.)

Pilula Scammonii Composita.

Resin of Scammony 1 ounce

Resin of Jalap 1 ounce

Curd Soap, in powder 1 ounce

Strong Tincture of Ginger..... 1 fluid ounce

Rectified Spirit..... 2 fluid ounces

Add the tincture and spirit to the soap and resins, and dissolve with the aid of a gentle heat, then evaporate in a water bath to a pilular consistence.

Dose—5 to 15 grains.

Scammonium, Scammony.—A gum-resin obtained by incision from the living root of *Convolvulus Scammonia*, imported from Smyrna and Asia Minor.

Collection.—The head of the root is cut off in a slanting direction 2 or 3 inches below the crown, and the milky juice which immediately flows out is collected in a mussel-shell. Sometimes the juice is allowed to dry in the shell, but as a rule it is emptied into a receptacle, and after a sufficient quantity has been collected, the whole is well mixed together and allowed to dry. In this way the purest scammony is obtained, known as *Virgin Scammony*.

Description.—The finest virgin scammony is in flat pieces or lumps, of a dark grey colour. It is easily broken with a lustrous fracture, and when powdered it becomes ashy grey, with a peculiar cheese-like odour. When rubbed with water it forms a white emulsion.

Uses.—Scammony is similar in action to jalap.

Chemistry.—Scammony is a gum-resin yielding from 88 to 90 per cent. of resin to ether, and leaving a fairly colourless residue. The active portion is the

resin called *jalapin*, $C_{34}H_{56}O_{16}$, which is readily soluble in ether, alcohol, and benzol, but insoluble in carbon disulphide.

Confectio Scammonii.

Scammony, in fine powder	3 ounces
Ginger, in fine powder	1½ ounce
Oil of Carraway	60 minims
Oil of Cloves.....	30 minims
Syrup.....	3 fluid ounces
Clarified Honey	1½ ounce

Dose—10 to 30 grains (10 grains of Scammony in 30 grains).

Pulvis Scammonii Compositus.

Scammony, in powder	4 ounces
Jalap, „	3 ounces
Ginger, „	1 ounce

Dose—10 to 20 grains (4 grains of Scammony in 8 grains).

Scammony is an ingredient in *Pilula Colocynthis Composita* and *Pilula Colocynthis et Hyoscyami*.

Adulterations.—Owing to its cost, scammony is extensively adulterated, most frequently by starch and chalk. Generally, scammony that is heavy, clayey-looking, not easily broken, and which when fractured does not exhibit a clean lustrous surface, should be rejected; and more especially so if it does not yield at least 80 per cent. of resin to ether.

A simple method of examining scammony is by taking a weighed portion and drying it at 212° F., when the loss equals moisture. The dried residue is then to be extracted with ether, and the ether evaporated in a weighed capsule spontaneously, and when dried, the capsule and its contents are heated for a time in a water-bath and weighed, the weight giving the amount of resin. The insoluble residue from the ether may then be treated with cold water to dissolve out the natural gummy constituents, and any residue left may be boiled with water, and the cold decoction tested with iodine for starch. Lastly, any residue insoluble in boiling water may be tested for chalk by effervescing with hydrochloric acid, neutralising the solution and testing for lime with ammonium oxalate.

NATURAL ORDER.—SOLANACEÆ.

Herbs or shrubs with alternate and often geminate leaves; inflorescence axillary or extra axillary; calyx persistent—5 partite; corolla regular with valvate æstivation; stamens equal in number to and alternate

with the corolla lobes ; anthers introse ; ovary superior 2-celled, with axile polyspermous placentæ ; fruit capsular or baccate ; seeds numerous and albuminous.

It is divided into two sub-orders :—

1. *Rectembryeæ*, embryo straight and short.

2. *Curvembryeæ*, embryo slender and curved.

Ex. *Solanum* and *Capsicum*.

DULCAMARA, Woody Nightshade.

—The dried young branches of *Solanum Dulcamara*, from indigenous plants.

Botanical Source.—*Solanum Dulcamara*, or Bitter Sweet, is a shrubby plant, found in hedges, with ovate, acute, entire and auriculate leaves ; inflorescence extra axillary cymes ; corolla rotate and purple ; anthers, connivant and standing out from the centre of the flower in a bright yellow mass ; fruit, baccate and bright scarlet.

Collection.—Branches of the first or second year should be chosen and collected in the autumn, when the leaves have fallen off.

Description.—Light, cylindrical, hollow stems, about the thickness of a quill, sometimes slightly warty or furrowed longitudinally.

Uses.—Supposed to be beneficial in rheumatism and skin diseases, but its action is very doubtful.

Chemistry.—The stems are said to contain 2-16ths per cent. of an alkaloid called *dulcamarine*, which is amorphous. They also appear to yield a bitter principle decomposable into sugar and solanine, but the quantity of this latter substance is very minute.

Infusum Dulcamaræ.

Dulcamara, bruised.....	1 ounce
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Incompatibles—The alkalies and their carbonates.

Dose—1 to 2 ounces.

CAPSICI FRUCTUS, *Capsicum*,
Chillies, *Cayenne Pepper*, *Guinea Pepper*.

The dried fruit of *Capsicum fastigiatum*, imported from Zanzibar.

Botanical Source.—The *capsicum* is a small shrub,

bearing oblong deep red capsules containing seeds of a pale colour.

Description.—Small membranous pods, orange-red in colour, with a very burning taste, conically pointed, smooth, shining, and somewhat shrivelled. From $\frac{1}{2}$ to $\frac{3}{4}$ inch long.

Uses.—Capsicum is a stimulant to the nerves of sensation, frequently administered in atonic dyspepsia, and to restore the function of the stomach after overdoses of opium. The Tincture of Capsicum added to Infusion of Roses in the proportion of $\frac{1}{2}$ drachm to 8 ounces, forms an excellent gargle for sore throat.

Chemistry.—The active principle is *capsicin*. It is obtained by making a tincture, evaporating to dryness, and extracting the residue with ether. It is a crystalline alkaloid, volatile by heat, producing extremely acrid and irritating fumes. When distilled with potash, it yields a substance very similar in smell to conia. It may be precipitated by tannic acid, and is very soluble in petroleum ether. From the pericarp of the fruit a pungent active principle, called *capsaicin*, has been isolated.

Tinctura Capsici.

Capsicum Fruit, bruised	$\frac{3}{4}$ ounce
Rectified Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—10 to 20 minims. As a gargle $\frac{1}{2}$ drachm to 8 minims.
(1 grain of Capsicum in 60 minims.)

Substitutions.—The fruits of *Capsicum annum* the ordinary Chillies of commerce. They are easily distinguished from the official capsicum by being three or more inches in length, and much broader.

NATURAL ORDER.—ATROPACEÆ.

Similar in characters to the Solanaceæ, but distinguished by the æstivation of the corolla being imbricate and in having one or more of the stamens sterile, with either extrorse or introrse anthers.

BELLADONNÆ RADIX, Belladonna Root. — The dried root of *Atropa Belladonna*, cultivated in England, but chiefly imported from Germany.

Botanical Source.—The *Atropa Belladonna* is a perennial herb, with ovate, acute, entire leaves, having an unequal base. The flowers are axillary, the calyx persistent, and the corolla campanulate, purple above, but greenish at the base. The fruit is baccate, bluish black in colour, and about the size of a small cherry.

Description.—In rough irregular pieces, of a dirty greyish colour, branched and wrinkled, and whitish internally. The roots are sometimes 2 inches thick, but those not thicker than the finger should be preferred as they yield nearly $\frac{1}{2}$ per cent. of atropia. The older roots are not by far so rich in that alkaloid.

Uses.—A powerful vasculo-cardiac stimulant, useful in syncope. It is also beneficial in neuralgia, rheumatism and acute inflammation generally. It is a powerful cerebro-spinal poison. Belladonna has been long used in ophthalmic practice to dilate the pupil of the eye: acting in a directly opposite manner to Calabar bean which contracts the pupil. It is also employed to check the secretion of milk.

Antidotes. — Emetics and afterwards astringents. Opium may be used, carefully avoiding narcotism.

Linimentum Belladonnæ.

Belladonna Root, in coarse powder.....	20 ounces
Camphor	1 ounce
Rectified Spirit	ad. 20 fluid ounces

(1 fluid ounce is equal to 1 ounce of Root.)

Moisten the Root with spirit for 3 days, then percolate.
(7 parts and 1 of chloroform form an excellent liniment
for lumbago, applied on spongio-piline).—*Squire.*

Atropia, $C_{17}H_{23}NO_3$, is prepared by macerating 2 lbs. of belladonna root, recently dried and in coarse powder, in 4 pints of rectified spirit for 24 hours, with frequent stirring. Transfer to a displacement apparatus and exhaust with 6 pints more of spirit by slow percolation. Add 1 ounce of slaked lime to the tincture placed in a bottle, and shake occasionally several times. Filter; add diluted sulphuric acid in very feeble excess, and filter again. Distil off 3-4ths of the spirit; add to the residue 10 fluid ounces of distilled water; evaporate at a gentle heat, but as rapidly as possible, until the liquid is reduced to 1-3rd of its volume and no longer smells of alcohol; then let it cool. Add very cautiously, with constant stirring, a solution of potassium carbonate, so as nearly to neutralise the acid, care, however, being taken that an excess be not used. Set to rest for 6 hours; then filter and add potassium carbonate, in

such quantity that the liquid shall acquire, on testing, alkaline re-action. Place it in a bottle with 3 fluid ounces of chloroform ; mix well by a frequently repeated brisk agitation, and pour the mixed liquids into a funnel furnished with a glass stop-cock. When the chloroform has subsided, draw it off and distil it in the water-bath from the retort, connected with the condenser. Dissolve the residue in warm rectified spirit ; digest the solution with a little animal charcoal ; filter, evaporate and cool, until colourless crystals are obtained.

In this process the natural bimalate of atropia is extracted by the alcohol, and is then decomposed by the lime, forming calcium malate, and setting free atropia, which is immediately converted into sulphate, as if left in contact with heated water the alkaloid decomposes. The first addition of potassium carbonate separates the excess of sulphuric acid and resinous matters ; while the second addition (this time in excess) sets free the alkaloid, which is then dissolved out by the chloroform. As thus prepared atropia is in colourless acicular prisms, very slightly soluble in water, but soluble in alcohol, ether and chloroform. Its watery solution is alkaline ; gives a precipitate with auric chloride, and powerfully dilates the pupil of the eye. When left with caustic

alkalies, atropia is decomposed into tropic acid, $C_9H_{10}O_3$, and tropinia, $C_8H_{15}NO$.

Liquor Atropiæ.

Atropia, in crystals.....	4 grains
Rectified Spirit.....	1 fluid drachm
Distilled Water	7 fluid drachms

($\frac{1}{2}$ grain of Atropia in 1 fluid drachm.)

Unguentum Atropiæ.

Atropia	8 grains
Rectified Spirit.....	30 minims
Prepared Lard	1 ounce

$\frac{1}{2}$ drachm containing $\frac{1}{2}$ grain of Atropia is sufficient for one application.

Atropiæ Sulphas, sulphate of atropia, prepared by mixing 120 grains of atropia with 4 fluid drachms of water, and gradually adding diluted sulphuric acid, until a perfectly neutral solution is obtained, and evaporating to dryness at a temperature not exceeding 100° F. It is a colourless powder, freely soluble in water.

Liquor Atropiæ Sulphatis.

Sulphate of Atropia	4 grains
Distilled Water	1 fluid ounce

($\frac{1}{2}$ grain of Sulphate of Atropia in 1 fluid drachm.)

BELLADONNÆ FOLIA.—The leaves of
Atropa Belladonna.

Collection.—They should be gathered when the plant is in flower.

Chemistry.—Belladonna leaves contain atropia.

Extractum Belladonnæ.

The expressed juice of the fresh leaves and young branches of Belladonna evaporated to an extract.

(100 lbs. of herb yield 16 lbs. dried, or 56 lbs. of juice, or 63 ounces of extract.)

Dose— $\frac{1}{4}$ to 1 grain.

Emplastrum Belladonnæ.

Extract of Belladonna	3 ounces
Resin Plaster	3 ounces
Rectified Spirit.....	6 fluid ounces

Rub the extract and spirit in a mortar, and when the insoluble matter has subsided, decant the clear solution. Remove the spirit by distillation, and mix the extract obtained with the resin plaster, by heating in a water-bath to a proper consistence.

Succus Belladonnæ.

Belladonna, fresh leaves, and young branches	7 lbs.
Rectified Spirit.....	q.s.

Bruise the Belladonna in a stone mortar, press out the juice, and to every 3 measures of juice, add one of spirit. Set aside 7 days, and filter. Keep it in a cool place.

Dose—5 to 15 minims.

Tinctura Belladonnæ.

Belladonna Leaves, in coarse powder.... 1 ounce
 Proof Spirit.....ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—5 to 20 minims. (1 grain of Belladonna in 22 minims
 —half the strength of London and Dublin.)

Antidotes—An emetic of 10 grains of sulphate of copper,
 afterwards opiates.

Unguentum Belladonnæ.

Extract of Belladonna 80 grains
 Prepared Lard 1 ounce

Rub down the Extract with a few drops of water before
 mixing with the lard.

STRAMONII FOLIA, *Stramonium*
Leaves.—The dried leaves of *Datura Stramonium*,
 (thorn apple), cultivated in Britain.

Botanical Source.—The *Datura* is an annual herb
 with a smooth dichotomous stem. The leaves, which
 arise from the forks of the stem, are ovate, acuminate,
 deeply dentate, and markedly petiolate. The flowers
 are axillary, and have a white infundibuliform corolla,
 about 3 inches long. The fruit is a capsule, covered
 externally with prickles and dehiscing septifragally,
 containing numerous dark coloured seeds.

Uses.—For making cigarettes for the relief of spasmodic asthma.

Chemistry.—The leaves contain, in addition to fixed oil, resin, wax, and about .03 per cent. of daturia—an alkaloid resembling atropia.

STRAMONII SEMINA, Stramonium Seeds.—The ripe seeds of the *Datura Stramonium*.

Description.—Dark brownish grey, reniform seeds, flat and very reticulated in the testa.

Uses.—Similar to belladonna.

Chemistry.—They yield about .1 per cent. of daturia, $C_{17}H_{23}NO_3$, isomeric with, and possessing the same properties as atropia. They also contain 25 per cent. of fixed oil.

Extractum Stramonii.

Stramonium Seeds, in coarse powder	16 ounces
Ether	q.s.
Proof Spirit	q.s.
Distilled Water	q.s.

100 lbs. of the Seeds yield from 12 lbs. to 14 lbs. of Extract.

DOSE— $\frac{1}{4}$ to $\frac{1}{2}$ grain.

Tinctura Stramonii.

Stramonium Seeds, bruised 2½ ounces
 Proof Spiritad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—10 to 30 minims (1 grain of Stramonium Seeds in 9 minims).

Antidotes—An emetic of 10 grains of sulphate of copper. afterwards opiates.

Datura Tatula.

The leaves have been strongly recommended to be used as cigarettes, which have been found useful in asthma.

HYOSCYAMI FOLIA, Henbane. —

The fresh and dried leaves of *Hyoscyamus niger*, from indigenous plants.

Botanical Source.—The henbane is a biennial or annual herb, of a pale, dull green colour, and feeling clammy to the touch. Both stem and leaves are pilose. The leaves are simple, sessile and ovate; with an acute apex and a sinuate dented margin. The flowers are axillary, and the corolla is infundibuliform; of a dirty straw colour, with violet veins. The fruit is a capsule, dehiscing transversely by a convex lid, from which circumstance it is called a pyxis or pyxidium.

Collection.—The biennial henbane is the more valuable, and is cultivated extensively at Hitchin. The first year's leaves gathered in August are usually dried; while the second year's leaves gathered in July are employed fresh for making the extract.

Uses.—The action of henbane is similar to belladonna, producing dilation of the pupil and a tendency to sleep. It is employed generally as a sedative, and to correct the griping effect of purgatives. In excessive doses it is a poison.

Chemistry.—The active principle of henbane is an alkaloid called *hyoscyamia*, $C_{15}H_{23}NO_3$. It is an oily liquid, which is soluble in water, ether, alcohol and chloroform, and smells and tastes somewhat of tobacco. It is decomposed by boiling with caustic alkalies, being resolved into hyoscinic acid, $C_9H_{10}O_3$, and hyoscina, $C_6H_{13}N$, the latter being a volatile alkaloid. The sulphate of hyoscyamia, which is a suitable and permanent form for the administration of this alkaloid, may be prepared by macerating henbane seeds in water, acidulated with sulphuric acid and then percolating to exhaustion. The fluid thus obtained is placed in flat

dishes and allowed to evaporate spontaneously, when it deposits a large group of crystals of potassium sulphate. The mother liquor is neutralised with barium carbonate, and then evaporated to dryness, and treated with ether, containing 10 per cent. of alcohol, which dissolves the alkaloid. The ethereal solution is then evaporated to dryness and the residue taken up with water acidulated with sulphuric acid, filtered to remove any resinous matter undissolved. Lastly, a solution of hyoscyamia thus obtained is evaporated to dryness over sulphuric acid in a dessicator.

Extractum Hyoscyami.

The expressed juice of the fresh leaves and young branches of Hyoscyamus, evaporated to an extract.

(100 lbs. yield 50 lbs. of juice, 5 lbs. of extract, or 15½ lbs. of dried leaves.)

Dose—5 to 10 grains.

Succus Hyoscyami.

Hyoscyamus, fresh leaves and young	
branches	7 lbs.
Rectified Spirit	q.s.

Bruise the Hyoscyamus in a mortar, press out the juice. and to every 3 measures of juice, add one of spirit. Set aside 7 days, and filter. Keep in a cool place.

Dose—½ to 1 drachm.

Tinctura Hyoscyami.

Hyoscyamus Leaves, in coarse powder .. 2½ ounces
Proof Spirit.....ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—½ to 1 drachm (7 grains of Hyoscyamus Leaves
in 1 fluid drachm.)

TABACI FOLIA, Tobacco.—The dried leaves of *Nicotiana Tabacum*, cultivated in America.

Botanical Source.—Tobacco is a herb with simple, sessile, oblong, lanceolate, and entire leaves, the lower ones being very large, often attaining the length of 2 feet. Inflorescence terminal and in panicles; corolla, tubular infundibuliform in shape, and a pale pink colour.

Uses.—As an injection for the relief of spasmodic cholera; but its employment is dangerous, as it is really a powerful poison, causing intense vomiting and purging, with cold sweats and entire muscular relaxation. The best antidotes are external warmth and diffusible stimulants.

Chemistry.—When tobacco is distilled with potassium hydrate it yields from 6 to 7 per cent. of a volatile

alkaloid called *nicotia*, $C_{10}H_{14}N_2$. It is an inflammable oily liquid, boils at $480^{\circ}F.$, and has an odour of tobacco. It is soluble in water and fixed oils, also in alcohol and ether. Its aqueous solution is strongly alkaline, and its salts are very deliquescent. It gives a precipitate with platinic chloride, tannic acid, mercuric chloride, and plumbic acetate. To obtain perfectly pure *nicotia*, a strong aqueous solution of tobacco (in which the alkaloid exists combined with citric and malic acids) is mixed with an excess of potassium hydrate, and shaken up with ether until nothing more can be extracted. Oxalic acid is then added to the ethereal solution, which causes the separation of oxalate of *nicotia* as a syrupy substratum. This is washed with ether until perfectly clean, and the pure oxalate thus obtained is then treated with excess of potash and shaken up with ether, and finally the alkaloid taken up with the ether is distilled in a current of hydrogen.

Enema Tabaci.

Leaf Tobacco	20 grains
Boiling Water	8 fluid ounces

Infuse half an hour and strain—for one enema.

NATURAL ORDER.—OLEACEÆ.

Trees or shrubs with opposite leaves; flowers regular and quaternary in calyx and corolla; stamens 2 (rarely 4) epipetalous; ovary superior 2-celled, with 2 suspended ovules in each cell; fruit frequently drupaceous; seeds albuminous.

MANNA.—The concrete saccharine exudation of *Fraxinus Ornus* and *Fraxinus rotundifolia*, imported from Sicily.

Botanical Source.—Manna ash is a small tree, apparently indigenous to Italy, having compound imparipinnate leaves, and bearing panicles of feathery flowers in early summer.

Collection.—Manna is obtained by making incisions into the bark, and placing leaves and sticks below them in the middle of summer. The clear juice flows out, concretes upon the leaves and sticks and produces what is known as flake manna, which is the article the most

valued in this country, although the manna in tears is usually purer.

Description.—In even, friable, and porous stalactiform pieces, of a yellowish white colour, curved on one side, with a faint odour and a sweet taste. The pieces are usually from 1 to 6 inches long, and from 1 to 2 inches wide.

Uses.—A mild laxative for children. Dose, 1 drachm and upwards.

Chemistry.—Manna contains from 60 to 80 per cent. of mannite and about 15 per cent. of dextrin and glucose. **Mannite**, $C_6H_{14}O_6$, crystallises in shining rhombic prisms, soluble in six parts of water, sparingly soluble in alcohol and insoluble in ether. Its solution is not affected by boiling with dilute acids or alkalies, and it does not reduce Fehling's solution of copper. It may be prepared artificially by treating glucose with sodium amalgam. When heated it melts at 330° F. Mannite is not a true sugar but a hexatomic alcohol, its rational formula being $C_6H_8O_6$. Mannite may be extracted from manna by treatment with boiling rectified spirit. Manna also contains a bitterish, astringent substance called **fraxin**, $C_{16}H_{18}O_{10}$, soluble in water and alcohol.

Substitutions.—Inferior mannas are converted by means of glucose into what is apparently fine flake manna. This adulteration may be detected by boiling with four parts of rectified spirit, when good manna will leave a hard, and the artificial manna a viscid residue.

OLIVÆ OLEUM, Olive Oil, Salad Oil.—The oil expressed in Europe from the fruit of *Olea Europæa*, from Italy, Sicily, Spain, and North America; a native of Asia.

Botanical Source.—The *Olea Europæa* is a fragrant tree cultivated in countries bordering on the Mediterranean. It bears a purple oval drupe about 1 inch in length, which, when preserved for table, becomes of a greenish brown colour.

Collection.—The oil exudes in the mesocarp of the fruit, which contains nearly 70 per cent. The fruits are crushed, put into boxes and submitted to pressure, and the oil thus extracted is called “virgin oil.” The contents of the boxes are then mixed with boiling water and once more pressed, when the second quality of oil is

produced. Lastly, if the residue be left in heaps until it becomes decomposed, the last and worst variety of oil is obtained.

Description.—Olive oil is very pale yellow in colour, with a faint odour and bland taste. Its sp. gr. is '916 at 63·5° F., and '910 at 77°. At 36° it begins to become turbid, and at 20° it separates into a solid and liquid portion. It is a non-drying oil, comparatively insoluble in alcohol, but miscible with twice its bulk of ether. The finest varieties of olive oil are the Florence and Provence; the Lucca and Genoa varieties are very good, while the Gallipoli, imported in casks, is inferior.

Uses.—Olive oil is a nutritive emollient, and an excellent antidote in cases of irritant poison.

Chemistry.—It contains about 72 per cent. of olein, and 28 per cent. of palmitin and stearin. It is readily saponifiable by alkalies and plumbic oxide.

Preparations.—Hard soap, soft soap, curd soap, soap liniment, soap plaster, lead plaster, &c.—(Refer to B.P.) Olive oil is also contained in the following preparations:—Cataplasma Lini, Charta Epispastica.

Emplastrum Ammoniaci cum Hydrargyro, Emplastrum Cerati Saponis, Emplastrum Hydrargyri, Emplastrum Picis, Enema Magnesiae Sulphatis, Linimentum Camphoræ, Unguentum Cantharidis, Unguentum Hydrargyri Compositum, Unguentum Hydrargyri Nitratis, and Unguentum Veratriæ.

Adulterations.—The mixture of olive oil with cheaper oils may be detected by the following tests:—


1. Olive oil is completely solidified at 32° F.
2. Mixed with 1-12th of its volume of acid solution of nitrate of mercury it becomes in four hours perfectly solid; any foreign oil present will float upon the surface.
3. When mixed with strong sulphuric acid and the mixture is stirred with a thermometer, the raising of the temperature effected is less in the case of pure olive oil than when adulterated.

The plan adopted is to take the temperature of 50 c.c. of the oil placed in a long tube; then add 10 c.c. of strong sulphuric acid, stir the whole up with a Centigrade thermometer, and note the rise in temperature which takes place. Thus treated—

Pure olive oil gives a rise of 42 degrees

Colza oil	„	„	58	„
Sesame oil	„	„	68	„
Hempseed oil	„	„	98	„
Linseed oil	„	„	130	„

Soaps are compounds produced by boiling olive or other oils or fats with alkalies or oxides and water. The olive oil may be looked upon as *glyceryl oleate*, $(C_3H_5)3C_{18}H_{33}O_2$, and when saponified it is converted into an oleate of the metal used and free glycerin (glyceryl hydrate C_3H_53HO). For example, in making Emplastrum Plumbi we have the following re-action



In the soaps of the B.P. the glycerin has been separated by adding salt to increase the density of the liquid, and so causing the soap to float, leaving the glycerin in the solution. In Emplastrum Plumbi the glycerin is driven off by the steam-bath heat employed ($230^\circ F.$), but in the various liniments both the soap and the glycerin are retained.

The following are the official soaps and preparations in which saponification takes place during manufacture :

A. Soaps ready formed, or made with soap, and containing no glycerin, unless purposely ordered, as in *Linimentum Potassii Iodidi c. Sapone*.

Sapo Durus. (*Hard Soap.*)

Composition—Chiefly sodium oleate ; entirely soluble in rectified spirit.

Uses—Diuretic in large doses, purgative combined with rhubarb for habitual constipation.

Dose—5 to 20 grains. In mineral acid poisoning, $\frac{1}{2}$ pint of the strong solution.

Sapo Mollis. (*Soft Soap.*)

Composition—Chiefly potassium oleate ; entirely soluble in rectified spirit, and not imparting an oily stain to paper.

Preparation—The oil is boiled with a solution of potassium hydrate, until the whole has become saponified ; it is then separated from the glycerin and water by means of evaporation, or by a strong solution of sodium chloride, which causes the soap to float to the surface by increasing the density of the solution.

Emplastrum Saponis.

Hard Soap	6 ounces
Lead Plaster	36 ounces
Resin, in powder	1 ounce

Linimentum Saponis.

	<i>British.</i>	<i>London.</i>
Hard Soap	2½ ounces	2½ ounces
Camphor	1¼ ounce	1¼ ounce
Oil of Rosemary	3 fluid drachms	13½ minims
Rectified Spirit.....	18 fluid ounces	18 fluid ounces
Distilled Water.....	2 fluid ounces	2 fluid ounces

Digest at a temperature not exceeding 70°, for 7 days, and filter.

Linimentum Potassii Iodidi c. Sapone.

Hard Soap, cut small.....	
Iodide of Potassium	of each 1½ ounce
Glycerin	1 fluid ounce
Oil of Lemon	1 fluid drachm
Distilled Water	10 fluid ounces

Pilula Saponis Composita.

Hard Soap	2 ounces
Opium	½ ounce
Water	q.s.

Dose—3 to 5 grains (4 of soap in 5).

B. Preparations in which soaps are formed during preparation, and containing all the natural glycerin from the oil used, except Emplastrum Plumbi.

Emplastrum Plumbi.

Oxide of Lead	64 ounces
Olive Oil	1 gallon
Water	3½ pints

Boiled down at 230° F. to a suitable consistence with frequent stirring; consists of plumbic oleate, the glycerin being evaporated off by the temperature employed.

Linimentum Ammoniæ.

Solution of Ammonia	1 fluid ounce
Olive Oil	3 fluid ounces

Contains ammonium oleate and glycerin.

Linimentum Calcis.

Solution of Lime.....	2 fluid ounces
Olive Oil	2 fluid ounces

Contains calcium oleate and glycerin.

Linimentum Hydrargyri.

Ointment of Mercury	1 ounce
Solution of Ammonia	1 fluid ounce
Liniment of Camphor.....	1 fluid ounce

(1 of ointment of mercury in 3.)

Contains ammonium oleate and glycerin.

NATURAL ORDER.—BORAGINACEÆ.

Similar to the Solanaceæ, but distinguished by their scorpioid or helicoid inflorescence and deeply 4-lobed ovary, with a solitary ovule in each lobe.

No plants of this order are now official.

ALKANET ROOT.—The dried root of *Anchusa tinctoria*, imported from Southern Europe and Asia Minor.

Alkanet root is easily recognisable from its deep red colour and characteristic marking, and is used as a dye and in colouring oil for perfumery.

SYMPHYTUM OFFICINALE,
Comfrey.—An indigenous herb with ovate-lanceolate, entire, decurrent, and pilose leaves; inflorescence a scorpioid cyme (of which it is a typical illustration), and regular flowers of a pale pink colour.

It is reputed as a vulnerary by herbalists.

NATURAL ORDER.—LABIATÆ.

Usually herbs with rectangular stems, and opposite, simple, and exstipulate leaves; flowers, irregular; calyx, persistent; corolla, labiate; stamens, 4 didynamous and epipetalous; ovary, deeply 4-lobed, with a bifid stigma and basilar style; fruit, 1—4 achænia, enclosed in persistent calyx.

LAVANDULÆ OLEUM.—The oil distilled in Britain from the flowers of *Lavandula vera*.

Botanical Source.—It is a hoary branched shrub,

about 4 feet high, with small linear oblong leaves and a rectangular stem. Inflorescence, a spike of verticillasters; the corolla labiate, and purple coloured, with a powerful and agreeable odour.

Collection.—The entire flowering tops are cut off and distilled with water, when the oil comes over, floating on the surface of the distillate. As a rule, the flowering tops yield $1\frac{1}{2}$ per cent. of essential oil. In a good season, at Mitcham, as much as 2 lbs. of oil have been obtained from 100 lbs. of flowering tops.

Description.—The oil of *Lavandula* is pale yellow, when first drawn over nearly colourless, with a sp. gr. of .87 to .94. It boils at 397° F. It is a mixture of eleopten and stearopten, the former having the formula $C_{10}H_{16}$, and therefore isomeric with turpentine, while the stearopten is identical with camphor. Oil of lavender is very soluble in rectified spirit, and less so in proof spirit. The oil is mellowed and remarkably improved in character by keeping in a cool place and apart from the light: but by exposure to the air it absorbs oxygen and becomes acid.

Spiritus Lavandulæ.

Oil of Lavender	1 fluid ounce
Rectified Spirit.....	49 fluid ounces
(9½ minims of Ol. Lavand. in 1 fluid ounce.)	

Dose—½ drachm to 1 drachm (1-5th the strength of 1864).

Tinctura Lavandulæ Composita.

Oil of Lavender	1½ fluid drachm
Oil of Rosemary	10 minims
Cinnamon Bark, bruised	150 grains
Nutmeg, bruised	150 grains
Red Sandalwood	300 grains
Rectified Spirit	ad. 40 fluid ounces

Macerate 7 days, then add the oils to the strained tincture.

Dose—½ to 2 drachms.

This tincture is used to colour the official arsenical solution.

Oleum Lavandulæ is also a constituent of Linimentum Camphoræ Compositum.

Adulteration.—The oil distilled from *Lavandula spica*, commonly called “oil of spike.” It is a French oil of inferior flavour and value, readily recognised by its inferiority in odour.

MENTHÆ PIPERITÆ OLEUM,
Oil of Peppermint.—The oil distilled in Britain from the fresh flowering tops of *Mentha piperita*.

Botanical Source.—It is a perennial herb, with a procumbent and ascending stem. The leaves are ovate, lanceolate, acute and decussate; inflorescence, a short spike of verticillasters (or clustered axillary cymes).

Collection.—Usually cut in August, and allowed to dry on the ground before being submitted to distillation with water.

Description.—Very pale yellow, with clean aromatic smell, sp. gr. from .84 to .92.

Uses.—Stimulant and carminative. Useful in flatulence.

Chemistry.—Contains a liquid portion called *menthene*, $C_{10}H_{18}$, and a stearopten (*menthol*, $C_{10}H_{18}H_2O$), which is the odoriferous portion, and constitutes, when crystallised, the whole Chinese oil of peppermint. One ton of the plant yields about 5 lbs. of oil.

Aqua Menthæ Piperitæ.

Oil of Peppermint	$1\frac{1}{2}$ fluid drachm
Water	$1\frac{1}{2}$ gallon

Distil 1 gallon. (About $\frac{1}{2}$ gtt. of Oil in 1 fluid ounce.)

Essentia Menthæ Piperitæ.

English Oil of Peppermint.....	1 fluid ounce
Rectified Spirit	4 fluid ounces
DOSE—10 to 20 minims. (1 of Oil in 5.)	

Spiritus Menthæ Piperitæ.

Oil of Peppermint	1 fluid ounce
Rectified Spirit	49 fluid ounces
(9½ minims of Ol. Menth. Pip. in 1 fluid ounce.	
DOSE—½ drachm to 1 drachm. (1-5th the strength of 1864.)	

Also used in *Pilula Rhei Composita*.

MENTHÆ VIRIDIS OLEUM, Oil of Spearmint.—The oil distilled in Britain from the fresh flowering plant of *Mentha viridis*.

Botanical Source.—Spearmint is an annual herb, with a smooth, erect stem; and lanceolate, acute and equally serrated, sessile, opposite, decussate leaves; inflorescence, a spike of verticillasters.

Collection.—The herb is cut and distilled with water when in full flower, usually in the month of August.

Description.—Colourless or pale yellow. Sp.gr. '914; boiling point 320° F.

Uses.—Stimulant carminative, and as a flavour in aperient draughts.

Chemistry.—It consists of a fluid portion or elæopten, isomeric with turpentine. When cooled to 25° F. it deposits a stearopten, called *menthol*, $C_{10}H_{18}H_2O$.

Aqua Menthæ Viridis.

Oil of Spearmint $1\frac{1}{2}$ fluid drachm

Water $1\frac{1}{2}$ gallon

Distil 1 gallon. (About $\frac{1}{2}$ gtt. of Oil in 1 fluid ounce.)

MENTHÆ PULEGII OLEUM.—

Oil of Pennyroyal.—The oil distilled from the fresh flowering herb of *Mentha Pulegium* cultivated in Britain. Not official.

Botanical Source.—Perennial herb, with branched and prostrate stem; leaves not exceeding $\frac{1}{2}$ an inch in length, ovate, crenate, petiolate, opposite, decussate and slightly pilose; inflorescence, axillary verticillasters; corolla, labiate and pale purple.

Collection.—The herb is collected when commencing to flower.

Description.—Very pale yellow. Sp. gr. .92; boiling point 395° F.

Uses.—Pennyroyal is stimulant and carminative.

Chemistry. — The odoriferous principle consists chiefly of a body isomeric with camphor, $C_{10}H_{16}O$.

ROSMARINI OLEUM, Oil of Rosemary.—The essential oil distilled from the flowering tops of *Rosmarinus officinalis*, cultivated in England.

Botanical Source.—Rosemary is a leafy shrub with simple, sessile, linear leaves, thick, revolute and hoary beneath. The flowers are greyish blue, labiate and axillary.

Collection.—The tops of rosemary should be cut when the flower commences to expand, and distilled with water.

Description.—A colourless oil, having a variable sp. gr. about .88; boiling at 365° F., and consisting chiefly of hydrocarbon, isomeric with turpentine. One cwt. of rosemary tops yields, as a rule, $4\frac{1}{2}$ ounces of oil.

Uses.—Stimulant and carminative. Used in perfumery and hair washes.

Spiritus Rosmarini.

Oil of Rosemary	1 fluid ounce
Rectified Spirit	49 fluid ounces
(1-5th the strength of 1864.)	

Also contained in Linimentum Saponis Compositum and Tinctura Lavandulæ Composita.

Among other plants of the *Labiata* which are used in the culinary and medicinal arts, may be mentioned :—

Marrubium vulgare, horehound.

Melissa officinalis, common balm.

Origanum vulgare, common marjoram.

Salvia officinalis, garden sage.

Thymus vulgaris, or garden thyme.

All of these should be studied from the plants themselves both in the fresh and dry state.

Thymol, which is yielded by several plants of this order, is said to have powerful antiseptic properties. Its sp. gr. is 1.028, and it is represented by the formula $C_{10}H_{14}O$. Thymol is permanently soluble in 1000 parts of water; also in rectified spirit, fats and oils.

Used in the form of spray, and as an ointment (1 in 8).

NATURAL ORDER.—SCROPHULARIACEÆ.

Herbs with usually opposite leaves and irregular anisomerous flowers ; calyx, persistent 4—5 partite ; corolla irregular, usually personate, calceolate or rotate ; stamens, 2 or 4, always the reverse of the number of the corolla lobes, and if 4 didynamous ; ovary, 2-celled with axile polyspermous placentæ ; fruit, capsular or rarely baccate ; seeds, albuminous.

DIGITALIS FOLIA, Foxglove. — The dried leaves of *Digitalis purpurea* from wild indigenous plants.

Botanical Source. — The *Digitalis* is a biennial herb, with a slightly pubescent stem and large, simple, ovate leaves having an acute apex, crenate margin and sheathing petiole. They are nearly smooth on the upper surface and downy beneath, the under surface being marked with prominent and reticulated veins. The inflorescence is a raceme of pale purple flowers. The calyx is persistent ; the corolla is campanulate-labiate ; the stamens are didynamous, and the fruit is a capsule with septicidal dehiscence.

Collection.—The leaves ought to be collected while the plant is in full flower; *i.e.*, when 2-3rds of the flowers are expanded.

Description.—The dried leaves exhibit the characteristics of those of the fresh plant described above.

Uses.—*Digitalis* is a sedative, given in cardiac and arterial diseases. In large doses it is an active poison, producing nausea, dizziness, dilated pupils and syncope, accompanied usually by great pain in the abdomen and purging.

Chemistry.—The active principle of *digitalis* is a neutral substance called *digitalin* (*Digitalinum B.P.*), which is extracted by an official process as follows:—Digest 40 ounces of *digitalis* leaf in coarse powder with 1 gallon of rectified spirit for 24 hours, at a temperature of 120° F.; then put them into a percolator, and when the tincture has ceased to drop exhaust the leaf by the slow percolation of another gallon of rectified spirit. Distil off the greater part of the spirit, and evaporate the remainder over a water-bath, until the whole of the alcohol has been dissipated. Mix the residual extract with 5 ounces of distilled water (to which $\frac{1}{2}$ ounce of acetic acid has been previously added), and digest the

solution thus formed with $\frac{1}{4}$ ounce of purified animal charcoal; then filter and dilute the filtrate with water until it measures 1 pint. Add solution of ammonia nearly to neutralisation, and afterwards 160 grains of tannic acid dissolved in 3 ounces of water. Wash the precipitate formed with a little water, mix it with a small quantity of rectified spirit and $\frac{1}{4}$ ounce of oxide of lead, and rub them together in a mortar. Place the mixture in a flask, and add to it 4 fluid ounces of rectified spirit; raise the temperature to 160° F., and keep it at this heat for about an hour; then add $\frac{1}{4}$ ounce of purified animal charcoal; filter; from the filtrate carefully drive off the spirit by the heat of a water-bath; and, lastly, wash the residue repeatedly with pure ether.

In this process the *digitalin* is dissolved out from the alcoholic extract by diluted acetic acid, and the solution partially decolorised and nearly neutralised. It is precipitated by tannic acid, which forms a tannate of digitalinum. On rubbing this with plumbic oxide and alcohol, plumbic tannate is formed, and the digitalinum set free, which dissolves in the alcohol.

Digitalinum is met with in porous scales, inodorous and intensely bitter; soluble in spirit, but almost insoluble in ether and water; soluble in acids, but does not neutralise

them. With hydrochloric acid it becomes first faint yellow, and afterwards green. It is precipitated by tannic acid. The dose is 1-60th to 1-30th of a grain made into pills, digitalinum being previously dissolved in spirit to assure thorough admixture.

Infusum Digitalis.

Digitalis Leaves, dried 30 grains
Boiling Distilled Water 10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Incompatibles—Most metallic salts, and preparations of cinchona.

Dose— $\frac{1}{4}$ to $\frac{1}{2}$ ounce.

Succus Digitalis. (Prepared like *Succus Hyoscyami*.)

Tinctura Digitalis.

Digitalis Leaves, in coarse powder $2\frac{1}{2}$ ounces
Proof Spirit ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—10 to 30 minims (1 grain of Digitalis in 9 minims).

Antidotes.—Friction and diffusible stimulants, with the external warmth well kept up.

Substitutions.—The leaves of *Verbascum Thapsus* (Mullein) may be at once distinguished from those of digitalis by being woolly on both surfaces.

SUB-CLASS IV.—MONOCHLAMYDEÆ.

[APETALÆ.]

Plants either monochlamydeous or achlamydeous.

NATURAL ORDER.—POLYGONACEÆ.

Usually herbs with alternate leaves and ochreate stipules; flowers, usually hermaphrodite or rarely unisexual; calyx, inferior; ovary, superior and 1-celled; fruit, a triangular nut.

RHEI RADIX, Rhubarb Root.—

The dried decorticated root of *Rheum officinale*, from China, Chinese Tartary and Thibet, and imported from Shanghai and Canton; and also brought overland by way of Moscow.

Collection.—The root is dug up early in autumn, its cortical layer pared off, and the remainder cut into suitable pieces, which after partially drying on a stove, are strung upon a cord and the drying completed spontaneously.

Description.—Sometimes in cylindrical and sometimes in plano-convex pieces, but very variable in shape and frequently pierced with a hole, in which may be found the remains of the string used to suspend the root while drying. The outer surface slightly shrivelled, and having here and there portions of the dark bark not thoroughly cut away. When broken transversely it is found to have medullary rays, in the shape of a number of short, broken, deep brown lines, traversing a whitish ground and forming a sort of internal ring of starlike spots. The interior of good rhubarb should be compact gritty, and finely veined, and there should be no decolorisation or sponginess. The yellow, dirty matter on the outside should not be turned red by a solution of boracic acid.

The source of the so-called Russian or Turkey Rhubarb has also been assigned to *Rheum palmatum* var. *Tanguticum*.

Uses.—Mild purgative, tonic and astringent; slowly empties the bowels, and afterwards acts as an astringent to confine them. Owing to this secondary action, it is exhibited in the first stage of diarrhœa, but it should not be given for constipation, unless combined with other

purgatives. The dose in powder as a stomachic is 1 to 5 grains, and as a purgative 10 to 30 grains. The most useful way of administering rhubarb with the latter object is, however, undoubtedly in the form of *Pulvis Rhei Compositus*, stirred up in water with a little aromatic spirit of ammonia.

Chemistry.—The root of rhubarb contains starch, and is loaded with raphides of calcium oxalate. In good samples these constitute 35 to 40 per cent. of the root. The yellow colour is due to *chrysophanic acid*, $C_{14}H_8O_4$. It is crystalline yellow, slightly soluble in water; very soluble in alcohol and ether, and is turned dark red by alkalies. The red solution caused by caustic potash, if evaporated to dryness, turns first violet and finally blue during that process. It is volatilised by heat with yellow fumes.

Rhubarb also contains three resins, named Aporetin, Phæoretin and Erythretein. Besides these it contains a considerable quantity of rheo-tannic acid, $C_{26}H_{26}O_{14}$, which is a yellow substance, soluble in water and alcohol, but insoluble in ether. It produces a greenish black precipitate with ferric sulphate and pale blue with ferrous salts. It is not really known to which

of these substances the medicinal action of rhubarb is due.

Extractum Rhei.

Rhubarb, sliced or bruised.....	16 ounces
Rectified Spirit.....	10 fluid ounces
Distilled Water	5 pints

100 lbs. of sound root yield 39 lbs. of extract.

DOSE—5 to 15 grains. (1 grain is equal to about 3 grains of Rhubarb.

Infusum Rhei.

Rhubarb, in thin slices	$\frac{1}{4}$ ounce
Boiling Distilled Water.....	10 fluid ounces

Infuse in a covered vessel for 1 hour, and strain.

Incompatibles—The mineral acids, metallic solutions, some astringent infusions. Alkalies darken the colour, but do not decompose it.

DOSE—1 to 2 ounces.

Pilula Rhei Composita.

Rhubarb, in fine powder	3 ounces
Socotrine Aloes, in fine powder	$2\frac{1}{4}$ ounces
Myrrh, in fine powder.....	$1\frac{1}{2}$ ounce
Hard Soap, in powder	$1\frac{1}{2}$ ounce
Oil of Peppermint	$1\frac{1}{2}$ fluid drachm
Treacle, by weight	4 ounces

DOSE—5 to 10 grains. ($2\frac{1}{2}$ grains of Rhubarb and 2 grains of Aloes in 10 grains.)

Pulvis Rhei Compositus. (*Dr. Gregory's Powder.*)

Rhubarb, in powder	2 ounces
Light Magnesia	6 ounces
Ginger, in powder	1 ounce

Dose—20 grains to 1 drachm. (13 grains of Rhubarb in 60 grains.)

Syrupus Rhei.

Rhubarb Root, in coarse powder	2 ounces
Coriander Fruit ,,	2 ounces
Refined Sugar	24 ounces
Rectified Spirit	8 fluid ounces
Distilled Water	24 fluid ounces

Product, 37 ounces by weight, or 28 ounces by measure.

Dose—I to 4 drachms.

Tinctura Rhei.

Rhubarb, in coarse powder	2 ounces
Cardamom Seeds, bruised.....	$\frac{1}{4}$ ounce
Coriander, bruised	$\frac{1}{4}$ ounce
Saffron	$\frac{1}{4}$ ounce
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

Dose—Purgative, 4 drachms to 1 ounce. Stomachic, 1 to 2 drachms. (22 grains of Rhubarb in $\frac{1}{2}$ fluid ounce.)

Vinum Rhei.

Rhubarb Root, in coarse powder	1 $\frac{1}{2}$ ounce
Canella Alba Bark, in coarse powder	60 grains
Sherry	ad. 20 fluid ounce

Dose—I to 2 drachms.

Substitutions.—1. East Indian rhubarb, an inferior variety, which is supposed to be the residue from which the refined rhubarb has been selected. It is hard and knotty, imperfectly trimmed, and has often brown or blackish stains.

2. Batavian, or Dutch trimmed rhubarb, is found in flattish pieces prepared in imitation of the so-called Turkey rhubarb.

3. English rhubarb is the root of *Rheum Rhaponticum*, cultivated in England at Banbury, in Oxfordshire. It is usually in sticks, or semi-cylindrical pieces, and more spongy in texture; differing somewhat in odour from true rhubarb. It may be distinguished by the entire absence of the star-like spots above mentioned; or, if they be present, they are never arranged in an irregular ring.

Adulterations.—Powdered rhubarb is frequently adulterated with starch and turmeric. The former may be discovered by the microscope, and the latter by a solution of boracic acid.

NATURAL ORDER.—PIPERACEÆ.

PIPER NIGRUM, Black Pepper.—The dried unripe berries of the *Piper nigrum*, from the East Indies.

Botanical Source.—The *Piper nigrum* is a perennial plant with a trailing or climbing stem, bearing spikes of sessile fruits about the size of a pea, at first red, but black when ripe.

Collection.—The berries are gathered before they are ripe, and dried in the sun. The proper stage for picking is when one or two berries at the base of the spike turn red.

Description.—Small round wrinkled berries, without stalks, brownish black externally, with a characteristic odour and taste of pepper.

Chemistry.—Pepper contains from 1 to 2 per cent. of a volatile oil, and an acrid resin which is soluble in alcohol and ether, and appears to be the pungent principle. Pepper also contains from 2 to 3 per cent. of a substance isomeric with morphia, called *piperia* or

piperine; its formula is $C_{17}H_{19}NO_3$. It is a neutral crystalline body, insoluble in water, fusible at $212^{\circ} F.$, volatile and soluble in alcohol, acetic acid, and ether. By the action of nitric acid it is decomposed into piperic acid and *piperidia*, $C_5H_{11}N$, which latter is an oily volatile alkaloid, having the smell of pepper and ammonia combined. An odour of benzoic aldehyd is given off during the process of its formation.

Confectio Piperis.

Black Pepper, in fine powder	2 ounces
Clarified Honey	15 ounces
Carraway Fruit, in fine powder	3 ounces

Dose—1 to 2 drachms (6 grains of Pepper in 1 drachm).

Also an ingredient in Confectio Opii (1 in 31) and Pulvis Opii Compositus (1 in $7\frac{1}{2}$).

Adulterations.—Whole pepper is tolerably clean when imported, but when ground is subject to many adulterations—such as starch, linseed and pea-meal, or sand, all of which may be detected by the microscope.

PIPER ALBUM, White Pepper.—The produce of the same plant as black pepper, only in this case the fruit has been allowed to ripen, and

the pulpy portion removed by soaking in water and rubbing. In all other points white pepper agrees with black.

PIPER LONGUM, Long Pepper.—The dried spadices of *Piper officinarum* and *Piper longum*, imported from India, Singapore, and Batavia.

Not official.

Description.—A spike containing a multitude of minute baccate fruits, closely packed, about $1\frac{1}{2}$ inch long and about $\frac{1}{4}$ inch thick, greyish white in colour, becoming when wetted deep reddish brown.

Uses.—Employed as a spice, and in veterinary medicine.

Chemistry.—Similar to that of ordinary pepper.

CUBEBA, Cubebs.—The dried and ripe fruit of *Cubeba officinalis*, wild and cultivated in Java.

Botanical Source.—A climbing plant similar to pepper, but having the flowers in racemes, with short foot stalks instead of spikes.

Description.—About the size of black pepper, but readily distinguished by having foot stalks attached (*Piper caudatum*) and by its characteristic odour.

Uses.—Either powdered cubebs or the oil is a mucous stimulant in gonorrhœa, but if used in the former state it should be freshly ground to evince real activity.

Dose of the powder, $\frac{1}{2}$ to 2 drachms, stirred up in a large quantity of water.

Chemistry.—The active principle is a volatile oil, and it also contains an inert crystalline substance called cubebin, very similar to piperine.

Cubebæ Oleum, Oil of Cubebs.—The volatile oil obtained in Britain by distilling cubebs with steam, between 450° and 480° F. It is colourless or pale greenish yellow, and is polymeric with oil of turpentine, its formula being $C_{30}H_{48}$. In cold weather it deposits crystals of *cubebene-hydrate*, $C_{30}H_{48}2H_2O$, commonly called cubeb camphor. Cubebs yield from 10 to 15 per cent. of volatile oil.

Dose, 5 to 20 minims.

Tinctura Cubebæ.

Cubebs, in powder	$2\frac{1}{2}$ ounces
Rectified Spirit	20 fluid ounces

Dose— $\frac{1}{2}$ to 2 drachms (7 grains of Cubebs in 1 fluid drachm).

MATICÆ FOLIA, Matico Leaves.—The dried leaves of *Piper angustifolium* (*Artanthe elongata*, B.P.) imported from Huanuco, and other parts of Peru.

Botanical Source.—A small rough-leaved shrub, growing in the moist woods of Bolivia, Peru, Brazil, and Venezuela.

Description.—As found in commerce it is in a mass of more or less broken leaves, which are noticed to be thick, acuminate, at the apex; and cordate and unequal at the base. They are tessellated on the upper surface, and downy on the under surface.

Artanthe adunca is another species of Matico, the leaves not reticulated on the upper and downy on the under surface.

Uses.—As a powerful topical astringent, and hæmostatic (*ā'ī'μā'* blood).

Chemistry.—Matico contains a little volatile oil, a trace of tannic acid, a little resin, and a crystallisable acid called artanthic acid. Infusion of Matico becomes dark when mixed with ferric chloride.

Infusum Maticæ.

Matico Leaves, cut small ½ ounce
Boiling Distilled Water..... 10 fluid ounces

Infuse in a covered vessel for ½ hour, and strain.

Incompatibles—Salts of iron.

DOSE—1 to 4 ounces.

NATURAL ORDER.—THYMELACEÆ.

Usually shrubs with entire and exstipulate leaves and hermaphrodite flowers; calyx, inferior, imbricated; stamens, inserted on the calyx, and twice as many as its lobes; ovary, superior and 1-celled; fruit, usually drupacious.

MEZERII CORTEX, Mezereon.—

The dried bark of *Daphne Mezereum* and *Daphne Laureola* (Spurge Laurel), cultivated in England and Central Europe, wild near Andover.

Botanical Source.—The *D. Mezereum* is a small shrub with evergreen, lanceolate, smooth leaves; and rose-coloured flowers appearing in almost sessile

bunches of three, before the leaves. The calyx is monosepalous; the corolla wanting; and the stamens eight, inserted inside the calyx in two rows; the fruit is baccate, red, and fleshy. The *D. Laureola* is a small shrub, with spatulate, evergreen, smooth and entire leaves. The flowers are like those of *D. Mezereum*, monochlamydeous. They are yellowish green, and are arranged in axillary cymes. The fruit is baccate and black.

Description.—A tough and pliable, fibrous bark; olive brown externally; and white and satiny internally, with a faint odour, and acrid taste.

Uses.—Emetic and purgative in large doses. In small doses diuretic. Externally, a powerful local irritant. The ethereal extract is a constituent of *linimentum sinapis compositum*. The bark is an ingredient in compound decoction of sarsaparilla. 60 grains to the pint.

Chemistry.—The bark contains a volatile oil and a glucoside, called *Daphnin* $C_{31}H_{34}O_{19}$, which, when boiled with a dilute acid, yields Daphnetin ($C_{19}H_{14}O_9$) and sugar; but the real active principle is an acrid

resin, which has not been properly investigated. The bark will give up its acrid properties to water, vinegar, spirit and oils.

Extractum Mezerei Æthereum.

Mezereon Bark, cut small	16 ounces
Rectified Spirit	8 pints
Ether	1 pint

NATURAL ORDER.—LAURACEÆ.

CAMPHORA, Camphor.—The concrete volatile oil obtained from the wood of *Cinnamomum Camphora* (*Camphora officinarum*), imported in a crude state from China and Japan, and purified by sublimation in this country.

Botanical Source.—The *Camphora officinarum* is a handsome, evergreen tree, abounding in the Eastern provinces of China, and in the Island of Formosa. Its wood is much used in China for making cabinets, as it resists the attacks of insects.

Collection.—Camphor, which is diffused through the whole plant, is obtained by cutting the wood into

small chips and boiling with water, when the camphor volatilises with the steam, and is collected in inverted copper pots. On its arrival in this country, it is re-sublimed into thin glass vessels (*bomboloes*), which are then broken, and the pure camphor taken out, in a bell-shaped cake, about 3 inches thick.

Uses.—Diaphoretic and anti-spasmodic. Good in fever, mania, spasmodic diseases, and recently recommended in diarrhœa. In large doses it is a sedative. Dose, 2 to 10 grains.

Chemistry.—It is white, translucent and crystalline, having the formula $C_{10}H_{16}O$, and its sp. gr. is $\cdot 996$. It volatilises slowly at ordinary temperatures, the sp. gr. of its vapour being $5\cdot 3$. It is soluble in 1,300 parts of water; but freely soluble in rectified spirit, chloroform, ether, acetic acid, and the volatile and fixed oils. It melts at $288^{\circ} F.$; boils at $400^{\circ} F.$, and burns in the air with a clear, bright flame. By distillation with phosphoric anhydride, it loses water and becomes converted into cymol $C_{10}H_{14}$; and when heated with oxidising agents, such as nitric acid, it is slowly oxidised, first into *camphoric acid* $C_{10}H_{16}O_4$, and afterwards into *camphoretic acid* $C_{10}H_{14}O_7$.

Aqua Camphoræ.Camphor, in small pieces $\frac{1}{2}$ ounce

Distilled Water 1 gallon

Dose—1 to 2 ounces. (About $\frac{1}{2}$ grain of Camphor in 1 fluid ounce.)**Linimentum Camphoræ.**

Camphor 1 ounce

Olive Oil 4 fluid ounces

Linimentum Camphoræ Compositum.Camphor $2\frac{1}{2}$ ounces

Oil of Lavender 1 fluid drachm

Strong Solution of Ammonia 5 fluid ounces

Rectified Spirit 15 fluid ounces

Spiritus Camphoræ.

Camphor 1 ounce

Rectified Spirit 9 fluid ounces

Dose—10 to 30 minims. (1 in 10.)

Tinctura Camphoræ Composita. (*Vide* page 54.)

Camphor is also contained in Unguentum Plumbi Subacetatis Compositum, Unguentum Hydrargyri Compositum, and all liniments except Ammoniæ, Calcis, Crotonis, Opii and Potassii Iodidi.

Other varieties of camphor are :—1. Borneo camphor, obtained from *Dryobalanops aromatica*. It consists of Borneol, and is harder than common camphor. It is

not so volatile, and has more the odour of patchouli. Its formula is $C_{10}H_{18}O$, and it is converted into common camphor by the action of nitric acid.

2. Ngai camphor is the produce of *Blumea balsamifera*, which is known to the Chinese, and stands in value between ordinary and Borneo camphor. It has been found to be similar in composition to *Borneo camphor*.

CINNAMOMI CORTEX, Cinnamon Bark.—The inner bark of shoots from the truncated stocks of *Cinnamomum Zeylanicum*, imported from Ceylon.

Botanical Source.—The *C. Zeylanicum* is a small evergreen tree, with shining leaves and panicles of greenish flowers.

Collection.—The stems of the trees are cut down so as to form “stools,” from which young shoots spring. They are cut when about two years old, in May and December, and the bark is then cut longitudinally and peeled off, and placed on a convex piece of wood so that it contracts into quills, which are dried first in the shade and afterwards in the sun.

Description.—In close double-rolled quills of a light yellowish brown, which break readily into fragments,

having a characteristic odour and an aromatic taste. A cold decoction should give no distinct blue with tincture of iodine.

Uses.—Aromatic, stimulant, and slightly astringent. Useful in nausea and flatulence, and to conceal the taste of other remedies.

Dose, in powder, 5 to 20 grains.

Chemistry.—The active principle is a volatile oil, of which good cinnamon yields nearly 1 per cent. It also contains sugar, starch, mucilage, and tannic acid; and it is the presence of the latter in considerable proportion which, taken together with a small quantity of starch, causes the imperfection of the iodine reaction as already noticed.

Cinnamomi Oleum B.P., Oil of Cinnamon.—The oil distilled from the cinnamon bark, after a preliminary maceration in a saturated solution of common salt, about 8 ounces being usually obtained from 80 lbs. of cinnamon.

Description.—Yellowish when fresh, becoming red by keeping; sp. gr. 1.035; boiling point from 428° to 446° F.

Chemistry.—Oil of cinnamon is chiefly composed of cinnamic aldehyd (*cinnamylhydride*, C_9H_7OH), which may be precipitated by shaking the oil with potassium bisulphite. This substance is a colourless oil, rapidly oxidising in the air, and becoming coloured owing to the solid resin and *cinnamic acid*, $HC_9H_7O_2$.

Cinnamic Acid resembles benzoic acid in its crystals, is slightly soluble in cold water, freely soluble in alcohol, fusible and volatile by heat. When distilled with barium hydrate, carbonic anhydride is given off, and the hydrocarbon of the series *styrol* or *cinnamol*, C_8H_8 , is produced.

By distilling cinnamic acid with oxidising agents, such as nitric acid or potassium dichromate and sulphuric acid, it is converted into benzoic aldehyd (volatile oil of bitter almonds) and benzoic acid.

Aqua Cinnamomi.

Cinnamon Bark, bruised	20 ounces
Water	2 gallons
Distil 1 gallon. (1 of Bark in 8.)	

Pulvis Cinnamomi Compositus.

Cinnamon, in fine powder	1 ounce
Ginger ,, 	1 ounce
Cardamoms ,, 	1 ounce

Dose—3 to 10 grains.

Tinctura Cinnamomi.

Cinnamon, in coarse powder..... $2\frac{1}{2}$ ounces
 Proof Spirit.....ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE— $\frac{1}{2}$ to 2 drachms. (7 grains of Cinnamon in 1 fluid drachm.)

Also an ingredient in Acidum Sulphuricum Aromaticum, Decoctum Hæmatoxyli, Infusum Catechu, Pulvis Catechu Compositus, Pulvis Cretæ Aromaticus, Pulvis Kino Compositus, Tinctura Cardamomi Composita, Tinctura Catechu, Tinctura Lavandulæ Composita, and Vinum Opii.

CASSIÆ CORTEX, Cassia Bark.—

The dried bark of *Cinnamomum Cassia*, imported from China and Java.

Botanical Source.—A tree similar to cinnamon.

Description.—Larger than cinnamon, and the quills never in double rolls. It also breaks with a shorter fracture, and a cool decoction gives a blue with tincture of iodine.

Uses.—Similar to cinnamon.

Chemistry.—Contains an essential oil exactly similar in a chemical point of view to cinnamon, but not so fine

in flavour ; and, therefore, commercially, not so valuable. The bark also contains tannic acid.

LAURI FRUCTUS, Bayberries.—The dried berries of *Laurus nobilis*, cultivated in England, and imported from the south of Europe. Not official.

Botanical Source.—It is an evergreen plant, with oblong lanceolate leaves, with a wavy margin ; inflorescence in axillary umbels. The berry egg-shaped, and bluish black, and 1-seeded.

Description.—Dried bayberries are egg-shaped ; a little larger than *Cocculus Indicus* ; dark brown in colour, with a wrinkled pericarp, to which the seed is not attached. When shaken near the ear they are heard to rattle. They contain both a volatile and a fixed oil, and have now almost entirely fallen out of use.

NECTANDRÆ CORTEX, Bebeeru Bark. — The bark of dried *Nectandra Rodiæi* (Greenheart), imported from British Guiana.

Botanical Source.—The *Bebeeru*, or greenheart, is a lofty tree, which is a native of British Guiana. In flat pieces about $\frac{1}{4}$ inch thick, externally greyish brown,

internally dark cinnamon brown. Strongly bitter and astringent to the taste.

Chemistry.—It contains resin, tannic acid, and about 2·5 per cent. of *beberia*, $C_{35}H_{40}N_2O_6$, which is a colourless amorphous alkaloid, very slightly soluble in water, and more so in ether, but most of all in absolute alcohol. It is soluble in dilute acids, forming yellow uncrystallisable salts.

Beberia Sulphas, sulphate of *beberia*,



is prepared, according to the B.P., by the following process:—Acidulate 1 gallon of water with $\frac{1}{2}$ fluid ounce of sulphuric acid, and with this exhaust 1 lb. of coarsely powdered *beberu* bark. Concentrate to 1 lb.; and, when cold, add gradually a little milk of lime, until the fluid is nearly saturated, but still retains a distinct acid re-action. After two hours filter and wash the precipitate, and add to the filtrate a slight excess of solution of ammonia. Collect the precipitate, and wash it twice with 10 ounces of cold water. Then squeeze, and dry the precipitate on a water-bath. Pulverise the dry precipitate and exhaust it with boiling rectified spirit. Add a little water, and recover the spirit by distillation. Add a

slight excess of sulphuric acid to the remainder, and evaporate to dryness in a water-bath. Pulverise the residue, and treat it with 1 lb. of cold water; filter; evaporate to the consistence of syrup, and separate it on plates and dry at a heat not exceeding 140° F.

By this process the sulphate is not produced perfectly pure. The addition of the lime neutralises the greater part of the acid first employed, and precipitates some of the natural acids of the bark. The ammonium hydrate precipitates the crude alkaloid, which is purified by solution in spirit and converted into sulphate. As thus prepared, the sulphate is in dark brown translucent scales; soluble in water and alcohol, and strongly bitter. The aqueous solution gives a white precipitate with barium chloride, proving the presence of sulphuric acid, and the addition of caustic soda to another portion, causes the precipitation of beberia as a yellowish white deposit, which is dissolved when the mixture is shaken up with twice its bulk of ether. The ethereal solution leaves, on evaporation, a yellow translucent residue, entirely soluble in diluted sulphuric acid.

SASSAFRAS RADIX, *Sassafras* Root.—The dried root of *Sassafras officinale*, from North America.

Botanical Source.—The *Sassafras officinale* is a small, dioecious tree common throughout the United States.

Description.—In light, porous, greyish yellow chips; and also in branched pieces, sometimes 8 inches in diameter at the crown. The bark is externally greyish brown; and internally, reddish brown; of a characteristic odour, and aromatic taste. The active principle is the essential oil, which resides chiefly in the bark of the root. It has a sp. gr. of 1.09; and, at a low temperature, it deposits crystals of sassafras camphor. The oil has been found to consist of an elæopten; isomeric with turpentine, $C_{10}H_{16}$, and a liquid body called safrol, $C_{10}H_{10}O_2$, forming with bromine characteristic crystals of a substitution product, having the formula $C_{10}H_5Br_5O_2$; melting at $338^{\circ} F$.

Sassafras root also contains tannic acid, and a colouring matter analagous to cinchona-red.

Uses.—Sudorific, stimulant and diaphoretic; also used as an adjunct to perfumery.

Preparations.—An ingredient in *Decoctum Sarsæ Compositum*, $\frac{1}{4}$ ounce to 1 pint.

NATURAL ORDER.—MYRISTICACEÆ.

MYRISTICA, Nutmeg.—The kernel of the seed of *Myristica officinalis*, cultivated in the Banda Islands of the Malayan Archipelago.

Botanical Source. — The *M. officinalis* is an evergreen tree, cultivated in Sumatra, Malacca and Singapore, but truly indigenous to the Malacca Islands. It bears nearly spherical drupes, about the size of a peach, the seed of which is covered with a scarlet-covered arillode, which is exposed by the spontaneous splitting open of the pericarp, when the fruit is ripe.

Collection.—When the pericarp splits, the fruit is collected, and the arillode carefully separated, dried and sold as mace. The seeds are thus isolated, dried by stove heat, until the kernels rattle in the shell, which latter is then broken, and the nutmeg rubbed over with dry lime and packed.

Description.—Oval, about an inch long ; externally reticulated, and internally greyish red, with dark brown veins.

Uses.—Stimulant and carminative.

Chemistry.—Nutmegs contain about 25 per cent. of fixed oil and 2 to 3 per cent. of volatile oil.

Oleum Myristicæ Expressum, expressed oil of nutmeg; butter or oil of mace; a concrete oil obtained from nutmegs by expression and heat.

This substance is a firm fat with an aromatic odour, and orange-red colour, imported from Singapore in blocks, enveloped in palm leaves. It is soluble in four parts of warm rectified spirit. It contains 6 per cent. of a volatile oil, and is otherwise a mixture of fats, the chief of which is *myristicin* $C_3H_5^3C_{14}H_{27}O_2(C_{45}H_{86}O_6)$, which constitutes about half the weight of the expressed oil. It is soluble in ether, but insoluble in cold alcohol, and when saponified it yields myristic acid $HC_{14}H_{27}O_2$.

Oleum Myristicæ, volatile oil of nutmeg, obtained in England by distilling nutmegs with water.

This oil is straw-yellow, sp. gr. .93; boiling point $329^{\circ} F.$, with a powerful odour of nutmeg. It chiefly consists of a hydro-carbon isomeric with turpentine, $C_{10}H_{16}$.

Spiritus Myristicæ.

Volatile Oil of Nutmeg	1 fluid ounce
Rectified Spirit.....	49 fluid ounces

Dose— $\frac{1}{2}$ drachm to 1 drachm, (1-5th the strength of 1864.)

Nutmeg is also an ingredient in Pulvis Catechu Compositus (1 in 10); Pulvis Cretæ Aromaticus (1 in 16); Spiritus Armoraciæ Compositus, and Tinctura Lavandulæ Composita.

Adulterations.—Long or wild nutmegs, the produce of *Myristica fatua*, recognised by their shape.

NATURAL ORDER.—ULMACEÆ.

ULMI CORTEX, Elm Bark.—The dried inner bark of *Ulmus campestris*, indigenous.

Botanical Source.—The well known common elm tree.

Collection.—The bark should be taken from the tree in early spring, and then deprived of its epiphloeum, and dried.

Description.—A tough brownish yellow bark, in

broad flat pieces, about 1-8th of an inch thick ; smooth on both surfaces, with broad marks where the knife has removed the outer layer.

Uses.—Demulcent and astringent. Administered in chronic skin diseases.

Chemistry.—It contains about 3 per cent. of tannic acid, with a little resin, gum and mucilage. Its decoction is precipitated by gelatin, and turned green by ferric chloride.

Decoctum Ulmi.

Elm Bark, cut in small pieces	2½ ounces
Distilled Water	ad. 20 fluid ounces

Boil 10 minutes.

Dose—2 to 4 ounces.

NATURAL ORDER.—CANNABINACEÆ.

CANNABIS INDICA, Indian Hemp.

—The dried flowering tops of the female plant of *Cannabis sativa*, from India, Herat and Persia.

Botanical Source.—The *C. sativa* is a dioecious annual, with an erect stem and digitate leaves, of

from 5 to 7 narrow lanceolate and serrated leaflets. Male flowers in drooping panicles ; female flowers in strobile-like racemes with leafy bracts, and a perianth covered with short brownish glands embracing the ovary. The fruits are small grey nuts, containing a single seed, and known in commerce as hemp seed.

Uses.—Narcotic and anti-spasmodic. It has been used with some success in neuralgia, tetanus, and even in hydrophobia, and the tincture has been exhibited with benefit in difficult menstruation.

The tincture, the colour of which is intensely green, gives a whitish precipitate of resin when mixed with water ; mucilage, therefore, should always be added in order to suspend the resin.

Chemistry.—The essential principles are a volatile oil and a resin, of which the latter is the chief active constituent. Good hemp should yield 7 per cent. of this resin, when treated with strong alcohol. It is blackish grey in colour, soluble in ether and in fixed and volatile oils. The volatile oil exists in the dried hemp to the extent of about '3 per cent. It consists of two hydro-carbons—*cannabene* $C_{18}H_{20}$. and *cannabene*

hydride $C_{18}H_{22}$. It has been asserted by some that this is almost as important a principle as the resin.

Extractum Cannabis Indicæ.

Indian Hemp, in coarse powder	16 ounces
Rectified spirit	4 pints

Dose— $\frac{1}{4}$ to 1 grain.

Tinctura Cannabis Indicæ.

Extract of Indian Hemp	1 ounce
Rectified Spirit	ad. 20 fluid ounces

Dose—5 to 20 minims. (1 grain of Extract in 22 minims.)

Antidotes—Give hot brandy and water, allow the patient to sleep, and apply a blister to the nape of the neck, if necessary, to control its violent action.

LUPULUS, Hop.—The dried strobiles of the female plant of *Humulus Lupulus*, cultivated in England and many parts of Europe and North America.

Botanical Source.—It is a twining plant, with opposite, palmate, serrated and stipulate leaves. The flowers are diœcious: the males in axillary panicles, and the females in strobiles with concave, membranous bracts, each of which has a fruit, at its base covered with yellow, aromatic glands.

Uses.—Aromatic and tonic. Slightly sedative and soporific, but uncertain in its action in the latter respect.

Chemistry.—The active principle of the hop exists in the lupulinic glands above described as covering the fruit, 6 lbs. of hops yielding about 6 ounces. These yellow grains are called lupulin, and have been found to contain 2 per cent. of volatile oil, 10 per cent. of bitter extractive, and 55 per cent. of resin. The active principle of hops may be extracted by spirit, water and petroleum ether.

Extractum Lupuli.

Hops	16 ounces
Rectified Spirit	1½ pint
Distilled Water	1 gallon

Dose—5 to 15 grains.

Infusum Lupuli.

Hops	½ ounce
Boiling Distilled Water.....	10 fluid ounces

Infuse in a covered vessel for 2 hours, and strain.

Incompatibles—Mineral acids; metallic salts.

Dose—1 to 2 ounces.

Tinctura Lupuli.

Hops	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

Incompatibles—The salts and preparations of iron.

Dose—¼ to 2 drachms.

NATURAL ORDER.—MORACEÆ.

CARICÆ, Figs.—The dried fruit of *Ficus Carica*, imported from Asiatic Turkey, Spain, and Portugal. [*Artocarpeæ*, Hanbury.]

Botanical Source.—The *F. Carica* is a tree with large rough palmate leaves. Inflorescence is a hypanthodium, which gives birth to an anthocarpous or collective fruit called a syconus.

Uses.—Slightly laxative, and an ingredient in Confection of Senna.

Chemistry.—It has not been fully investigated, but contains a large quantity of grape sugar, which constitutes nearly 70 per cent. of the dried fruit.

MORI SUCCUS, Mulberry.—The juice of the ripe fruit of *Morus nigra*.

Botanical Source.—A tree with alternate, slightly cordate, acuminate, serrated, pubescent, and stipulate leaves. The flowers are monœcious—male flowers in a spike, and female flowers in catkins. The fruit is a black sorosis.

Description.—Dark violet colour, with an acidulous taste; contains a large quantity of grape sugar, and nearly 2 per cent. of malic acid.

Syrupus Mori.—The fruit from which the juice is pressed should be perfectly ripe, and used immediately after it is gathered.

USES.—Refrigerant and slightly laxative, and administered to children; employed as a colouring agent.

NATURAL ORDER.—EUPHORBIACEÆ.

Herbs, shrubs, or trees with an acrid milky juice; flowers, unisexual, monœcious or diœcious; calyx, sometimes present, but frequently absent, both the male and female flowers being enclosed in an involucre furnished with glands; male flowers, with one or more stamens and 2-celled anthers; female flowers, with superior, usually 3-celled ovary, frequently supported on a gynophore, with 1 or 2 suspended ovules in each cell; fruit, tricoccus.

CASCARILLÆ CORTEX.—The dried bark of *Croton Eluteria*, from the Bahama Islands.

Description.—In quills, 2 or 3 inches in length and from 2 to 5 lines in diameter, dull brown, and coated with white crustaceous lichen; breaks with a short resinous fracture, and emits a fragrant odour when burnt.

Uses.—Aromatic stimulant and tonic, and to some extent febrifuge. It is usually prescribed in dyspepsia, cholic, and general debility. It is a constituent in fumigating pastilles.

Chemistry.—The bitter principle is called *cascarillin* $C_{12}H_{18}O_4$. It is a neutral substance freely soluble in ether and hot alcohol, but sparingly soluble in water and chloroform. The bark also contains 1 per cent. of volatile oil and 15 per cent. of resin, with gum and tannin. Tincture of Cascarilla becomes milky when poured into water.

Infusum Cascarillæ.

Cascarilla Bark	1 ounce
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel 1 hour, and strain.

Incompatibles—Lime water, metallic salts, mineral acids.

Dose—1 to 2 ounces.

Tinctura Cascarillæ.

Cascarilla Bark, bruised	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

Dose—½ to 2 drachms. (7 grains of Cascarilla in 1 fluid drachm.)

Adulterations.—The bark of *Croton lucidus* has lately been met with mixed with cascarilla. It is very similar in appearance, but its tincture does not precipitate in water.

TIGLII SEMINA. — The seeds of *Croton Tiglium*, imported from Ceylon and the Indian Archipelago.

Botanical Source.—The *C. Tiglium* is a small tree, indigenous to the Malabar coast, with brown, 3-celled capsules, each cell containing one seed.

Description.—Egg-shaped seeds, about ½ inch long, with a bright cinnamon-brown coat, which is easily scraped off, showing a black testa beneath. From the hilum the raphè is seen to run to the other end of the seed terminating in the chalaza, which is easily visible as a dark spot.

CROTONIS OLEUM, Croton Oil.—

The oil expressed from the seeds, which yield from 50 to 60 per cent.

Description.— It is a brownish yellow, viscid oil, with a very burning taste.

Uses.— Drastic purgative, in doses of 1-3rd to 1 minim. Applied to the skin it produces pustules. May be administered, made into a pill with bread crumb or cocoa butter.

Antidotes—Emetics, followed by mucilaginous drinks and opium.

Chemistry.—The true irritant principle has not been isolated, but in addition to the ordinary fixed oils it is found to contain acetic, butyric, valerianic, *tiglinic* ($C_5H_8O_2$), and *crotonic* acids $C_4H_6O_2$, all of which are volatile fatty acids.

Linimentum Crotonis.

Croton Oil	1 fluid ounce
Oil of Cajuput	3½ fluid ounces
Rectified Spirit	3½ fluid ounces

(1 of Croton Oil in 8.)

EUPHORBIIUM, *Euphorbium*. — The acrid resinous exudation from incisions in the stem of *Euphorbia resinifera*, imported from Mogadore, and a native of Morocco.

Description.—In regular pieces of dull yellowish brown waxy-looking matter, sometimes hollow in the centre, but generally enclosing a spine or fragment of the flower stalk. The dust, when inhaled, causes violent sneezing. It contains an amorphous resin soluble in alcohol, and with an acrid burning taste; and also *euphorbon*, which crystallises from ether, and is free from acidity. If euphorbon be moistened with a little sulphuric acid, a fine violet is produced on the careful addition of nitric acid.

Uses.—Powerful irritant; now only used as an ingredient in paint for preventing the attacks of insects.

RICINI SEMINA.—The seeds of *Ricinus communis*, from the East Indies.

Botanical Source. — The *R. communis* is a perennial plant, but when cultivated in England becomes annual, owing to its being unable to withstand the winter. The leaves are large, simple, palmatifid,

peltate, and serrated. The flowers are monoëcious in terminal panicles; the lower usually male, and the upper female. The fruit is a tricoccus capsule, covered with spines and having 3 cells, each cell being 1-seeded.

Description.—Grey shining seeds, with brownish black bands or spots, about 6-10ths of an inch in length, and 4-10ths wide. They are, as a rule, both larger and flatter than croton seeds.

Chemistry.—The active principle is a fixed oil, of which they afford about 50 per cent.

RICINI OLEUM, Castor Oil.—The oil expressed from the seeds above described, imported chiefly from Calcutta.

Description.—A viscid colourless or very pale yellow oil; sp. gr. .96. It does not entirely solidify at zero, and when exposed to the air in thin layers it dries.* It is miscible in all proportions with absolute alcohol and in glacial acetic acid, and is soluble in twice its bulk of rectified spirit.

* Although *Ol. Ricini* does not entirely solidify until cooled to -18° C. it throws down a considerable white deposit in extreme cold, which makes keeping it in large stock very unprofitable.

Uses.—A rapid but mild cathartic; very useful as a purgative for women and children, and in diseases accompanied by inflammatory action in the bowels.

Infant dose, 1 to 3 drachms; adult dose, $\frac{1}{2}$ to 1 ounce, in an emulsion with yolk of egg or mucilage, the yellow of one egg being sufficient to suspend an ounce of oil.

Chemistry.—Castor oil is a mixture of olein with a little palmatin. The olein, however, is peculiar to castor oil, yielding, when saponified, *ricinoleic acid*, $C_{18}H_{34}O_3$. By treatment with ammonia, ricinolein yields a solid crystalline substance, *ricinolamide*, $C_{18}H_{35}NO_2$.

Varieties.—The finest castor oil is that produced by expression without heat, and afterwards bleached by exposure to the sun on the roofs of houses; the second variety is known as Calcutta oil, prepared by heat; and of late years a quantity of Italian oil has come into the market, which is very tasteless. It is expressed from seeds grown in Italy, and also from those imported for the purpose from India.

Adulteration.—Castor oil is sometimes adulterated with olive or lard oils, which may be distinguished by their insolubility in rectified spirit.

Collodium Flexile.

Collodion	6 fluid ounces
Canada Balsam	120 grains
Castor Oil.....	1 fluid drachm

Castor oil also enters into the composition of Linimentum Sinapis Compositum and Pilula Hydrargyri Subchloridi Composita.

KAMALA.—A powder consisting of the minute glands which cover the capsules of *Rottlera tinctoria*, imported from India.

Botanical Source.—The *R. tinctoria* is a shrub, or small tree, with dioecious flowers in terminal panicles. The tricoccus capsule, which ripens in spring, is covered with minute, sessile, bright red glands.

Description.—Kamala is a fine granular, mobile powder, brick-red in colour, which mixes difficultly with water, but yields its colouring matter to alcohol, ether, chloroform, benzol, acetic acid, and potassium hydrate. Heated in a flame it ignites like lycopodium, and leaves only $1\frac{1}{2}$ per cent. of a grey ash.

Uses.—For the expulsion of *tænia*. The dose is 30 to 110 grains, given in a little strong brandy-and-water.

Chemistry.—Kamala contains traces of tannic acid, gum, volatile oil, and 80 per cent. of *rottlerin* $C_{22}H_5O_6$. This is a reddish yellow resin extracted by ether, which appears to be a compound of two other resins differing only in their degree of solubility and in their melting point.

NATURAL ORDER.—ARISTOLOCHIACEÆ.

ARISTOLOCHIÆ RADIX,
Birthwort.—The dried root of *Aristolochia longa*, or *A. rotunda*, found in England, and also imported from Southern Europe.

This root is not now officially prescribed in medicine, but is frequently employed by herbalists for criminal purposes, on account of its so-called emmenagogue properties. It was also an ingredient in a once famous remedy for gout, known as the Duke of Portland's powder. Like serpentary it contains a little essential oil and a bitter principle apparently identical with *aristolochin*, but which has never really been properly investigated. The dose in powder is from 20 to 60 grains.

SERPENTARIÆ RADIX,
Serpentary Root, and Virginian Snake
Root.—The dried root of *Aristolochia Serpentaria*,
from Southern parts of North America.

Botanical Source.—A herb with a perennial stalk,
growing in shady woods, and producing small dull
purple flowers.

Description.—A knotted, contorted rhizome, about
1 inch in length by 1-8th of an inch in thickness;
bearing on its under side a bunch of numerous slender
matted rootlets, about 3 inches long. It is dull yellowish
brown in colour, and has an aromatic odour and taste.

Uses.—Diaphoretic, diuretic, stimulant and tonic.
Said to be useful in chronic rheumatism, dyspepsia, low
fever, and as a promoter of eruption in *exanthemata*.
It is useful in certain stages of typhus, and its action in
many cases is similar to that of guaiacum, only not so
irritating to the bowels.

Chemistry.—Serpentary contains about $\frac{1}{2}$ per cent.
of volatile oil, and the same of resin. The outer layer
contains tannin, and also an amorphous bitter principle
precipitated by lead acetate.

Infusum Serpentariæ.

Serpentary Root	$\frac{1}{2}$ ounce
Boiling Distilled Water	10 fluid ounces

Infuse in a covered vessel for 2 hours, and strain.

DOSE—I to 2 ounces.

Tinctura Serpentariæ.

Serpentary Root, in coarse powder	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

DOSE— $\frac{1}{2}$ to 2 drachms. (7 grains of Serpentary Root in 1 fluid drachm.)

Also in Tinctura Cinchonæ Composita, $\frac{1}{2}$ ounce to the pint.

NATURAL ORDER.—SANTALACEÆ.

SANTALI LIGNUM. — The wood of *Santalum album*, from the East Indies. It is not much known in English commerce, but yields a light yellow thick essential oil of the sp. gr. .963, which is employed as a perfume, and has lately been prescribed as a substitute for copaiba in gonorrhœa.

NATURAL ORDER.—CUPULIFERÆ

(vel. CORYLACEÆ).

QUERCUS CORTEX, Oak Bark.—

The dried bark of small branches and young stems of *Quercus pedunculata*, collected in spring, from trees grown in Britain.

Description.—It is in channelled pieces, greyish and shining outside; cinnamon coloured inside; brittle, fibrous, and strongly astringent. The outer surface is dotted over with small scars, while the inner surface is longitudinally striated.

Uses.—An astringent in diarrhœa; the decoction makes a good gargle for sore throat and an efficient injection in *leucorrhœa*.

Chemistry.—Contains about 10 per cent. of a special variety of tannic acid, called *querci-tannic* acid, which precipitates gelatin, and the precipitate does not decompose so readily as that produced by ordinary tannic acid. It is distinguished from tannic acid by not being

convertible into gallic acid. A decoction of oak-bark becomes dark purple on the addition of ferric chloride.

Decoctum Quercus.

Oak Bark, bruised	1½ ounce
Distilled Water	ad. 20 fluid ounces

Boil 10 minutes in a covered vessel ; produce, 1 pint.

Incompatibles—Metallic salts ; solution of isinglass ; decoctum cinchonæ. Alkalies destroy its astringency.

GALLÆ, Nut Galls. — The excrescences caused by the punctures and the deposited ova of *Diplolepis Gallæ tinctoriæ*, found on the *Quercus infectoria*, chiefly from Asia Minor.

Botanical Source.—The female insect above named is furnished with a little, oblong organ, called an ovipositor, with which she pierces the tender shoots of the oak, and deposits an egg. The irritation thus caused in the plant produces a flow of juice to the spot and the gradual formation of an excrescence around the egg, called a gall, inside which the larva is hatched and undergoes its various transformations. At the end of about six months it becomes a winged insect, which bores its way to the surface of the gall and escapes. The best galls are those which are gathered before the

insect obtains its freedom, and which are, therefore, not perforated.

Description.—Heavy, hard, rounded bodies of a bluish green colour, yellowish white within. The upper half of the gall is covered with small pointed nobs arranged irregularly and the lower half is smooth.

Varieties.—The finest are the Mosul galls, and the next variety Aleppo galls, imported from the Levant. Of these there are two varieties, the best being the blue galls without perforation, and the inferior the white galls, which are of a light greyish or yellowish hue, and are perforated. A third variety, called large Mecca galls, are imported from Bussorah. They are also round in shape and have a circle of horned protuberances at their centre. These latter are sometimes called Dead Sea apples.

Chemistry.—Galls contain from 30 to 40 per cent. of tannic acid, and about 3 per cent. of already formed gallic acid, with some extractive matter. An infusion of galls is acid to test paper, gives a dirty white precipitate with gelatin, and an inky black with ferric chloride.

Tinctura Gallæ.

Galls, in coarse powder	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

Dose—½ to 2 drachms. (7 grains of Galls in 1 fluid drachm.)

Unguentum Gallæ.

Galls, in fine powder	80 grains
Benzoated Lard	1 ounce

Unguentum Gallæ c. Opio.

Ointment of Galls	1 ounce
Opium, in powder	32 grains

Substitutions. — Chinese galls. From the *Rhus semialata* (*Anacardiaceæ*). They are light and hollow, and distorted by numerous protuberances, and completely covered by a thick velvety grey down.

2. English galls, smooth, brown, and slightly speckled pale brown excrescences formed on the ordinary English oak.

TANNIC ACID, $C_{27}H_{22}O_{17}$, is obtained by the following process:—Expose powdered galls to a damp atmosphere for two or three days, and afterwards add sufficient ether to form a soft paste. Let this stand in a well-closed vessel for 24 hours, and then, having

quickly enveloped it in a linen cloth, submit it to strong pressure so as to separate the liquid portion. Reduce the pressed cake to powder, mix it with sufficient ether (to which 1-16th of its bulk has been added) to form a soft paste, and press this as before. Mix the expressed liquids, and expose the mixture to spontaneous evaporation until, by the aid subsequently of a little heat, it has acquired the consistency of a soft extract; then place it in earthen plates or dishes, and dry in a hot air chamber at a temperature not exceeding 212° F.

In this process the extraction of the tannic acid is accomplished by means of moist ether, because the tannic acid, although insoluble in pure ether, is very soluble in ether containing a little water, with which it forms a heavy yellow liquid, which has been supposed to be a definite chemical compound.

Tannic acid is soluble in water, and very soluble in glycerin and dilute alcohol. Its watery solution gives a precipitate with gelatin, ferric chloride, and with salts of lead and antimony. Exposed in contact with water, its solution becomes mouldy, and is converted into gallic acid and glucose. Heated to 620° F. it decomposes, forming *pyro-gallic* acid, $C_6H_6O_3$. A piece of raw hide immersed in a solution of tannic acid,

completely absorbs the whole of it, forming leather, and leaving pure water.

Glycerinum Acidi Tannici.

Tannic Acid	1 ounce
Glycerin	4 fluid ounces

Mix and dissolve with a gentle heat.

Suppositoria Acidi Tannici.

Tannic Acid	36 grains
Benzoated Lard	44 grains
White Wax	10 grains
Oil of Theobroma	90 grains

3 grains of Tannic Acid in each. Divide into 15 grain suppositories, or into 12 equal parts.

Suppositoria Acidi Tannici c. Sapone.

Tannic Acid	36 grains
Glycerin of Starch	50 grains
Curd Soap	100 grains
Starch	q.s.

Divide like Suppositoria Acidi Tannici.

Trochisci Acidi Tannici.

Tannic Acid	360 grains
Tincture of Tolu	$\frac{1}{2}$ fluid ounce
Refined Sugar, in powder	25 ounces
Gum Acacia, in powder	1 ounce
Mucilage of Gum Acacia	2 fluid ounces
Distilled Water	1 fluid ounce

Divide into 720 lozenges.

DOSE—1 to 6 (each lozenge contains $\frac{1}{2}$ grain of Tannic Acid).

GALLIC ACID, $\text{H}_3\text{C}_7\text{H}_3\text{O}_5 \cdot \text{H}_2\text{O}$, is prepared by the following process :—Place 1 lb. of galls, in coarse powder, in a porcelain dish ; pour on as much water as will convert them into a thick paste, and keep them in this moistened condition for six weeks, at a temperature of between 60° and 70° F. ; adding water from time to time to substitute what is lost by evaporation. At the end of that time boil the paste for 20 minutes with 45 fluid ounces of water ; strain through calico, and, when the fluid is cold, collect on a filter the crystalline deposit which has formed, and let it drain. Press it strongly between the folds of filtering paper ; and re-dissolve it in 10 ounces of boiling distilled water. When the fluid is cooled to 80° F., pour it off from the crystals which are formed ; wash these with 3 ounces of ice cold water ; and dry them, first by filtering paper, and finally at a temperature not exceeding 100° F.

During the six weeks fermentation, the tannic acid of the galls is converted into gallic acid which is extracted by boiling water, and deposited on cooling.

Gallic acid is very slightly soluble in cold water and in ether, but freely soluble in boiling water and alcohol. Its precipitate gives no solution with gelatin, but it gives a bluish black precipitate with ferric chloride.

Heated to 410° F. it volatilises and becomes converted into pyro-gallic acid and CO₂. Its solutions in alkalies absorb oxygen, and become brown. It is very soluble in glycerin.

The dose is 2 to 10 grains as a hæmostatic.

Glycerinum Acidi Gallici.

Gallic Acid	1 ounce
Glycerin	4 fluid ounces

Mix and dissolve with a gentle heat.

NATURAL ORDER.—ALTINGIACEÆ.

STYRAX PRÆPARATUS (B.P.),

Prepared storax.—A balsam obtained from the bark of *Liquidambar orientale*,* and purified by dissolving in rectified spirit, and straining.

Botanical Source.—The *L. orientale* is a tree growing in the south western parts of Asia Minor, and often attaining a height of 50 feet. It has palmate leaves very like those of the common plane tree.

* According to Hanbury this plant belongs to the *Hamamelideæ*, Sub-class *Calycifloræ*, Sub-division *Epigynæ*.

Collection.—The inner bark is boiled in water and submitted to pressure in horse-hair bags, when the balsam exudes. It is then sent to Smyrna and exported *via* Trieste.

Description.—A brownish-yellow, semi-fluid resinous mass, semi-transparent and similar in consistence to thick honey. Heated to 212° F. it becomes more fluid, and boiled with a solution of potassium dichromate and sulphuric acid it evolves the odour of bitter almonds (by evolution of benzoic aldehyd).

Uses.—Storax agrees in medicinal properties with the other true balsams, and is given as a stimulant expectorant in chronic bronchitis.

Dose 10 to 20 grains.

Chemistry.—Contains styrol, styracin, cinnamic acid and resin. *Styrol* is a liquid hydrocarbon C_8H_8 , having a sp. gr. of .924, and boiling at 295° F., soluble in alcohol and ether. *Styracin* or *meta-cinnamein*, $C_{18}H_{16}O_2$, is polymeric with cinnamic aldehyd C_9H_8O . (See Oil of Cinnamon.) It is a crystalline solid, soluble in alcohol and ether, and convertible by the action of potassium hydrate into potassium cinnamate and cinnamic alcohol (*cinnyl hydrate* C_9H_9HO). These

various substances may be separated by first distilling at 212° F. which separates styrol, then treating with hot water to remove the cinnamic acid, and finally dissolving out the styracin by ether.

Prepared Storax is an ingredient in Tinctura Benzoini Composita, $1\frac{1}{2}$ ounce to 1 pint.

Adulteration and Substitutions.—1. The crude storax is much adulterated by sawdust, which is left behind in the alcohol preparation ordered by the B.P.

2. The liquid storax of Rosamala is a very fine article sometimes imported from the Eastern Archipelago, and is the produce of *Liquidambar Altingiana*.

NATURAL ORDER.—SALICACEÆ.

SALICIS CORTEX, Willow Bark.—

The bark of various species of willow, from British plants.

Botanical Source.—The *Salix fragilis*, *S. alba* and *S. caprea* are all diœcious trees, with the flowers achlamydeous and in catkins.

Description.—In thin quills, usually with a greenish brown epidermis and white inner surface, but varying much in appearance according to its age and source.

Uses.—Tonic and astringent, and to some extent anti-periodic; chiefly used for the manufacture of salicin.

Chemistry.—Contains much tannin, and a bitter principle called salicin.

Salicin $C_{13}H_{18}O_7$, is a crystalline glucoside soluble in cold water and alcohol, but not in ether. It is made by boiling an infusion of willow bark with freshly precipitated plumbic hydrate, which precipitates tannin and other matters, then filtering and crystallising the salicin. Boiled with diluted sulphuric acid it yields glucose and *saligenin* $C_7H_8O_2$, and the solution neutralised gives a fine blue with ferric chloride. Acted upon by potassium dichromate and sulphuric acid it yields *salicylol*, or *salicylous acid* $C_7H_6O_2$, which is oil of meadow-sweet.

Salicylic Acid $H_2C_7H_4O_3$, is closely allied to salicylol, and was formerly obtained from it, but is now made by heating carbolic acid (*phenyl hydrate* C_6H_5HO)

with potassium hydrate, and then passing through carbonic anhydride between 212° and 482° F., which produces *inter alia* potassium salicylate. This salt is then made into a saturated solution and acidulated with hydrochloric acid, when the salicylic acid crystallises out.

It is only slightly soluble in cold but readily in boiling water. Its solution gives a deep purple with ferric salts. It has been much used lately in the treatment of rheumatism.

DIVISION II.—GYMNOSPERMIA.

Plants having their ovules naked and impregnated by the direct action of the pollen.

NATURAL ORDER.—CONIFERÆ.

LARICIS CORTEX, Larch Bark.—

The dried bark, deprived of its outer layer, of *Larix Europæa*, a native of Central Europe, cultivated in Britain.

Description.—In flattish pieces or large quills, reddish brown externally; and smooth and pinkish brown or pale yellow, internally. It breaks with a short fracture and smells distinctly of turpentine.

Uses.—Hæmostatic and astringent.

Chemistry.—Larch Bark contains a peculiar variety of tannic acid, which gives olive-green precipitates with ferric salts. It also yields a crystalline substance called *larixin* $C_{10}H_{10}O_5$, which may be obtained by evaporating an infusion of the bark to a syrupy consistence; distilling and crystallising the distillate.

It is slightly soluble in cold water, but freely in boiling water and alcohol. It is sparingly soluble in ether. Its aqueous solutions have a feeble acid re-action, and give a purple colour with ferric chloride and a gelatinous precipitate with baryta water.

Tinctura Laricis.

Larch Bark, in coarse powder $2\frac{1}{2}$ ounces
Rectified Spiritad. 20 fluid ounces

Macerate in 15 ounces of Spirit for 48 hours, then percolate with the remaining Spirit. Afterwards press the marc, mix the products, filter, and add Spirit to 1 pint.

DOSE—20 to 30 minims.

TEREBINTHINA VENETA,
Venice Turpentine. — The oleo-resin obtained by incision from *Larix Europæa*, imported from Italy, Switzerland, and the Tyrol.

Chemistry.—It yields about 15 per cent. of volatile oil and the rest resin. The true Venetian turpentine is rarely imported into this country, the article sold as Venice turpentine being a mixture of ordinary resin and oil of turpentine, which may be readily distinguished from the true article by the facility with which it dries up when spread on paper.

JUNIPERI FRUCTUS. — The fruit of *Juniperus communis*, from Northern Europe, and cultivated in England.

Botanical Source.—The *J. communis* is a dioecious shrub with dark green, subulate leaves in whorls of three. The fruit is a purple black galbulus, covered with greenish blue and consisting of three fleshy bracts enclosing as many seeds.

Chemistry.—The active principle is a volatile oil.

JUNIPERI OLEUM.—The oil distilled in Britain from the unripe fruit of the plant above described.

Description. — Juniper oil is colourless, or light greenish. Sp. gr. $\cdot 83$. Its formula is $C_{10}H_{16}$. It is, therefore, isomeric with oil of turpentine, and about equally soluble in alcohol.

Spiritus Juniperi.

Oil of Juniper	1 fluid ounce
Rectified Spirit	49 fluid ounces

($9\frac{1}{2}$ minims of Ol. Junip. in 1 fluid ounce.)

Dose— $\frac{1}{4}$ drachm to 1 drachm. (1-5th of the strength of 1864).

Uses.—Stimulant and diuretic, in doses of 4 to 6 minims, usually prescribed with sweet spirit of nitre. It is also used in the manufacture of Holland's gin.

JUNIPERI OLEUM EMPYREUMATICUM. — *Juniperus Oxycedrus* yields Huile de Cade, or Juniper tar, by destructive distillation.

SABINÆ CACUMINA, Savin. — The fresh and dried tops of *Juniperus Sabina*, collected in spring, from plants cultivated in Britain.

Botanical Source.—The *♀. Sabina* is a small, bushy shrub, with minute, ovate, convex, imbricated and appressed leaves in whorls of four. The fruit is a bluish galbulus, about the size of a currant.

SABINÆ OLEUM, Oil of Savin.—The oil distilled in Britain from the fresh plant, which yields about 3 per cent. It is colourless, or pale yellow; sp. gr. .915, moderately soluble in spirit and freely soluble in ether. It is isomeric with turpentine $C_{10}H_{16}$.

Uses.—An emmenagogue, and frequently used to procure abortion, but usually with fatal effects, being a powerful irritant poison.

The dose in powder is 4 to 10 grains, and of the oil 1 to 5 minims.

Tinctura Sabinæ.

Savin Tops, dried and coarsely powdered	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—20 minims to 1 drachm. (1 grain of Savin in 9 minims.)

Antidotes—Emetics; afterwards opiates and demulcents.

Unguentum Sabinæ.

Fresh Savin Tops, bruised	8 ounces
Yellow Wax	3 ounces
Prepared Lard.....	16 ounces

Digest 20 minutes, and strain through calico.

AMERICAN TURPENTINE, Horse Turpentine.—The produce chiefly of *Pinus palustris* and also of *P. tæda* and *P. pinaster*, imported from America.

Collection.—It is obtained by cutting a hollow in the trunk of the tree, and collecting the juice which exudes, transferring it to casks.

Description.—A yellowish white semi-solid substance resembling honey, with a strong odour of turpentine.

Chemistry.—Contains about 15 per cent. of a volatile oil and .85 per cent. of resin.

TEREBINTHINÆ OLEUM.—The oil distilled from the oleo-resin described above as American turpentine.

Preparation.—When the crude oleo-resin is distilled with water in a copper still, the volatile oil distils over, and the resin remains in the retort.

Chemistry.—The volatile oil of turpentine is a limpid, colourless, and inflammable liquid, having the formula $C_{10}H_{16}$, and a sp. gr. .864. It is very slightly soluble in water, more so in rectified spirit, and miscible in all proportions with ether, absolute alcohol, and benzol. It has the power of dissolving volatile and fixed oils, resins, caoutchouc, sulphur, and phosphorus. When exposed to the air it gradually absorbs oxygen and becomes converted into resin. When treated with hydrochloric acid it forms two crystalline compounds, $C_{10}H_{16}.HCl$ and $C_{10}H_{16}2HCl$, which are known as artificial camphors. Treated with strong nitric acid it inflames and is converted into a variety of substances.

Dose, as a diuretic and hæmostatic, 5 to 20 minims : as an anthelmintic, from 2 to 4 fluid drachms, mixed with 4 to 6 drachms of castor oil.

Confectio Terebinthinæ.

Oil of Turpentine.....	1 fluid ounce
Liquorice Powder	1 ounce
Clarified Honey	2 ounces

Dose—1 to 2 drachms. (15 minims of Turpentine in 1 drachm.)

Enema Terebinthinæ.

Oil of Turpentine	1 fluid ounce
Mucilage of Starch.....	15 fluid ounces
Mix for 1 Enema.	

Linimentum Terebinthinæ.

Oil of Turpentine	8 fluid ounces
Soft Soap	1 ounce
Camphor	$\frac{1}{2}$ ounce

Linimentum Terebinthinæ Aceticum.

Oil of Turpentine.....	1 fluid ounce
Acetic Acid	1 fluid ounce
Liniment of Camphor.....	1 fluid ounce

Unguentum Terebinthinæ.

Oil of Turpentine.....	1 fluid ounce
Resin, in coarse powder.....	60 grains
Yellow Wax.....	$\frac{1}{2}$ ounce
Prepared Lard	$\frac{1}{2}$ ounce

CREASOTUM, Creasote ($\kappa\epsilon\acute{\iota}\alpha\varsigma$ flesh, $\sigma\acute{\omega}\zeta\omega$ I preserve).—A product of the distillation of wood tar.

Description.—A colourless or slightly yellow liquid, with a strong empyreumatic odour. Its name is derived from its antiseptic power of preserving flesh. Unlike phenyl hydrate, it is insoluble in commercial glycerin, nor does it cause any effect when added to the solution known as collodium B.P.

Uses.—Creasote is a popular remedy for toothache, and employed by dentists in dressing carious teeth. It is frequently prescribed in order to allay sickness in doses of $\frac{1}{2}$ to 1 minim; also in phthisis and chronic bronchitis, especially where there is a tendency to hæmorrhage.

Chemistry.—Consists chiefly of *creasol*, $C_8H_{10}O_2$, which forms an almost insoluble crystalline salt with ammonium hydrate. Like carbolic acid, it coagulates albumen. Sparingly soluble in water, freely soluble in alcohol, ether, and glacial acetic acid. Sp. gr. 1.071. It is dextro-rotatory, and is not solidified by the cold produced by a mixture of hydrochloric acid and sodium sulphate.

Mistura Creasoti.

Creasote	16 minims
Glacial Acetic Acid.....	16 minims
Spirit of Juniper	$\frac{1}{2}$ fluid drachm
Syrup	1 fluid ounce
Distilled Water	15 fluid ounces
DOSE— $\frac{1}{2}$ to 1 ounce.	

Unguentum Creasoti.

Creasote	1 fluid drachm
Simple Ointment.....	1 ounce

Vapor Creasoti.

Creasote	12 minims
Boiling Water	8 fluid ounces

SANDARACH.—A white brittle resin, the produce of *Callitris quadrivalvis*, imported from Mogadore. It is not used in medicine, but its powder is known under the name of **POUNCE**.

RESINA (B.P.), Resin, or Rosin.—*Rosin colophony*, the residue left after the distillation of the oil of turpentine.

Description.—Translucent, amber coloured, and brittle, with a shining fracture, easily reduced to a white powder, fusible by heat, and burning with a dense, yellow, smoky flame.

Uses.—Slightly stimulant, but chiefly employed in pharmacy for its adhesive qualities.

Chemistry.—Pure resin has a sp. gr. of 1·07, and consists chiefly of a crystallisable acid called abietic or sylvic acid, and a non-crystallisable resin known as pinic acid, both isomeric, and having the formula $C_{20}H_{30}O_2$. These acids may be separated by fractional solution in cold spirit of wine, the latter being the more soluble. When acted upon by alkalies, resin combines with them, forming resin soap, which is a constituent of the ordinary yellow kitchen soap. In this decomposition

the acids of resin give up one atom of hydrogen in exchange for the alkaline base.

Specially Distinctive Tests.—Common resin and all coniferous products containing it, except Canada balsam, are—1. Completely soluble in chloroform, soluble in ether, and the latter solution is not rendered turbid by alcohol.

2. Soluble in alcohol, and the solution is coloured greenish brown by alcoholic ferric chloride (1 in 10), and precipitated by saturated alcoholic plumbic acetate, not soluble on boiling.

3. Acted upon by a saturated solution of sodium carbonate.

4. Extracted by petroleum spirit and the solution evaporated, a residue is obtained which is coloured reddish violet with blue streaks by impure chloral hydrate.

Emplastrum Resinæ.

Resin	4 ounces
Lead Plaster.....	32 ounces
Hard Soap	2 ounces

Unguentum Resinæ.

Resin, in coarse powder.....	8 ounces
Yellow Wax.....	4 ounces
Simple Ointment.....	16 ounces

Resin is also contained in Charta Epispastica, Emplastrum Cantharidis, Emplastrum Hydrargyri, Emplastrum Picis, Emplastrum Saponis and Unguentum Terebinthinæ.

Resin plaster is an ingredient in Emplastrum Belladonnæ, Emplastrum Opii, Emplastrum Calefaciens, and Emplastrum Plumbi Iodidi.

THUS AMERICANUM, Common Frankincense.—A concrete acrid turpentine obtained by spontaneous exudation from the *Pinus tæda* and *P. palustris*, from the Southern States of North America.

Description.—It is a softish, bright yellow, opaque solid, having the odour of turpentine.

Chemistry.—Thus Americanum is a mixture of essential oil of turpentine and resin, and enters into the composition of Emplastrum Picis.

TEREBINTHINA CANADENSIS, Canada Balsam.—An oleo-resin, obtained by incision in the stem of *Abies balsamea*, from Canada.

Description.—A pale yellow oleo-resin, perfectly transparent and of the consistence of very thin honey.

When mixed with 1-6th of its weight of magnesia it solidifies, and it also dries up spontaneously on exposure to the air. It contains about 18 per cent. of volatile oil and 82 per cent. of resin.

Uses.—The medicinal properties of Canada Balsam are similar to turpentine, but more mild. Dose 20 to 30 grains. It is chiefly used for mounting microscopic objects.

A constituent of *Charta Epispastica* and *Collodium Flexile*.

Specially Distinctive Tests.—This substance is—
1. Completely soluble in chloroform, and gives a perfect solution in ether, which becomes turbid on the addition of alcohol.

2. An alcoholic solution, becomes turbid with an alcoholic solution of ferric chloride (1 in 10), but clears up on boiling.

3. The extract from the alcoholic solution is coloured violet by impure chloral hydrate containing alcoholate.

PIX BURGUNDICA, Burgundy Pitch.—Stated in the B.P. to be the resinous exudation from the stem of *Abies excelsa*, imported from Switzerland.

Collection.—The crude oleo-resin, which flows from the tree, is melted in hot water ; strained through a cloth and cast into moulds. Although officially stated to be imported from Switzerland, it is really principally produced in the Black Forest, Austria and Finland.

Description.—Burgundy pitch should be of a dull reddish brown colour, hard and brittle, free from air vesicles, and giving off no water when heated. It should be soluble in rectified spirit and glacial acetic acid.

Chemistry.—Burgundy Pitch consists chiefly of resin, with a little volatile oil of turpentine, which escapes dissipation during the melting.

Emplastrum Picis.

Burgundy Pitch	26 ounces
Frankincense	13 ounces
Resin	4½ ounces
Expressed Oil of Nutmeg	1 ounce
Yellow Wax.....	4½ ounces
Olive Oil	2 fluid ounces
Water	2 fluid ounces

Emplastrum Ferri.

Hydrated Peroxide of Iron, in fine powder	1 ounce
Burgundy Pitch	2 ounces
Lead Plaster	8 ounces

Substitutions.—False Burgundy pitch is made by mixing American turpentine with palm oil and resin.

PIX LIQUIDA, Wood Tar.—Obtained by the destructive distillation of the wood of *Pinus sylvestris*, from Finland, Northern Russia and Sweden.

Uses.—A common remedy in cutaneous diseases.

Chemistry.—The distillate from wood consists of two portions: the lighter aqueous portion which contains impure acetic acid, wood spirit, acetone and creasote, and is separated for the manufacture of these substances; and the heavier portion, which is the true official tar, and is a mixture, affording, amongst other products, *toluene* C_7H_8 ; *naphthalene* $C_{10}H_8$; and *anthracene* $C_{14}H_{10}$.

Unguentum Picis Liquidæ.

Tar.....	5 ounces
Yellow Wax.....	2 ounces

PIX NIGRA.—The residuum in the still after the distillation of tar. Not official.

CLASS II.—MONOCOTYLEDONES.

Having the seed monocotyledonous; germination endorhizal; root fibrous; stem endogenous; leaves usually with parallel venation, and parts of the flowers in multiples of 3.

SUB-CLASS I.—DICTYOGENÆ.

Having the leaves with reticulated venation; the underground stem and root with concentric rings; and the floral envelopes verticillate.

NATURAL ORDER.—SMILACEÆ.

SARSÆ RADIX, Jamaica Sarsaparilla.

—The dried root of *Smilax officinalis*, a native of Central America, and imported from Jamaica.

Botanical Source.—The various species of *Smilax* which yield sarsaparilla are evergreen twining shrubs, with unisexual and diœcious flowers.

Description.—Roots about as thick as a goose quill, reddish brown externally, folded in bunches about 18 inches long, and covered with small rootlets, which are known technically as “the beard.” A section of good sarsaparilla should show a thick medullium and only a very slight deposit of starch in the inner cortical layer. A drop of sulphuric acid should colour the whole section dark purple, and a cool decoction should not become blue with tincture of iodine.

Uses.—Alterative. Used in secondary syphilis, and all diseases dependent on a depraved state of the system.

Chemistry.—The active principles are a very minute quantity of volatile oil and a white crystallisable body called *smilacin* or *parillin*. Parillin is prepared by exhausting the root with heated rectified spirit, and then evaporating until the remaining fluid weighs only 1-6th of the amount of the root started with. This liquid is then gradually diluted with $1\frac{1}{2}$ times its volume of water, which causes the evaporation of crude parillin.

This deposit, after settling, is then treated with $\frac{1}{2}$ its volume of alcohol, filtered, and again precipitated with water, and the precipitate washed with dilute spirit not containing more than 25 per cent. of alcohol, in which it is insoluble. It may lastly be again dissolved in rectified spirit, agitated with animal charcoal and crystallised. Thus prepared, parillin is obtained in pure white thin scales or prisms, only soluble in 10,000 parts of cold water, but more freely in boiling water. Its best solvent is rectified spirit, but it also forms a solution with chloroform which cannot be filtered.

Specially Characteristic Tests.—1. With strong sulphuric acid parillin gives a pure yellow solution becoming cherry red at the edges.

2. Heated on a water bath with diluted sulphuric acid (10 per cent.), it becomes first greenish, then red, and finally brown.

3. A solution of parillin in chloroform treated with dry hydrochloric acid gas suddenly becomes brown by transmitted light, and full green by reflected light.

4. A similar green fluorescence is observed when it is dissolved in any dilute mineral acid, but too much water causes it to disappear.

Decoctum Sarsæ.

Jamaica Sarsaparilla, cut transversely .. 2½ ounces
 Boiling Distilled Water..... 1½ pint

Digest 1 hour, boil 10 minutes in a covered vessel ; product
 1 pint.

Incompatibles—Alkalies ; lime water ; plumbi acetas ; some
 salts of mercury, and other metals.

Dose—2 to 10 ounces.

Decoctum Sarsæ Compositum.

Jamaica Sarsaparilla, cut transversely .. 2½ ounces
 Sassafras, in chips ¼ ounce
 Guaiacum Wood Turnings ¼ ounce
 Fresh Liquorice Root, bruised ¼ ounce
 Mezereon Bark 60 grains
 Boiling Distilled Water..... 1½ pint

Digest 1 hour, boil 10 minutes in a covered vessel,
 product, 1 pint.

Dose—2 to 10 ounces.

Extractum Sarsæ Liquidum.

Jamaica Sarsaparilla, cut transversely .. 16 ounces
 Distilled Water at 160°..... 14 pints
 Rectified Spirit 1 fluid ounce

Product 8 ounces ; sp. gr. 1.095. 100 lbs. of Root yield
 from 25 to 30 lbs. of solid extract. The cortical part
 will yield one half.

Dose—2 to 4 drachms. 1 fluid ounce is equal to 2 ounces
 of Root.

Other Varieties of Sarsaparilla.—The various

sarsaparillas of commerce are divided into two classes—

1. "Mealy," or "gouty," which contain much starch.
2. "Lean," or non-mealy, having very little starch, and consequently more valuable for extract.

I. *Mealy Sarsaparillas.*

1. *Honduras*, in rolls 30 inches long, and closely wound round with a long root so as to form a neat bundle.

2. *Guatemala*, in similar bundles to the Honduras, but of a more decidedly orange tint. The rootlets become thicker as they descend, and the bark inclines to peel off. It is supposed to be the produce of *S. papyracea*.

3. *Brazilian*. Not much imported, but when seen it is in bundles over 3 feet in length and 6 inches in diameter, tightly compressed, and neatly cut at the ends. These bundles are wound round by the stem of a plant not a smilax. Brazilian is supposed also to be chiefly *S. papyracea*.

II. *Non-Mealy Sarsaparillas.*

1. The official sarsa, *S. officinalis*, already described.
2. *Mexican*. Supposed to be produced by *S. medica*,

and not in bundles, with portions of the rhizome and stem attached. The stem portions are angular and thorny.

3. *Guayaquil*. Similar to Mexican, but the portions of the stem are round and not thorny, and the root has many rootlets attached.

SUB-CLASS II.—PETALOIDEÆ.

Endogenous herbs with parallel veined leaves, and flowers in white or coloured perianths.

SUB-DIVISION I.—EPIGYNÆ.

With superior perianths and inferior ovaries.

NATURAL ORDER.—ORCHIDACEÆ.

Usually herbs terrestrial or epiphytcal; leaves entire and sheathing; perianth irregular and superior, with the large outer leaf (*labellum*) often assuming curious shapes; stamens gynandrous, with the pollen grains collected into masses (*pollinia*); ovary inferior 1-celled. with 3 parietal placentæ.

VANILLA.—The dried pods of *Vanilla planifolia* indigenous to Eastern Mexico.

Also the dried pods of *V. aromatica*.

Botanical Source.—A tree growing in damp situations, and producing a long fleshy fruit.

Description.—In long dry pods, 3-8ths of an inch long and 3-10ths to 4-10ths of an inch wide, growing narrower and hooked at the stalk end. The surface is striated longitudinally, and is shining, unctuous, and often covered with an efflorescence of minute crystals.

Chemistry.—It owes its flavour to a substance called **Vanillin**, which crystallises in prisms, and is soluble in alcohol, ether, and oils, but less so in water. The flavour of vanillin is diffusible to a remarkable extent in sugar. It is slightly acid to test paper, and its formula is $C_{16}H_8O_6$. Vanillin has been lately obtained artificially by oxidising *coniferin*, $C_{16}H_{22}O_8 \cdot 2H_2O$, a body existing in the sap of pines, with potassium dichromate and sulphuric acid.

Salep.—The root of several indigenous and other species, chiefly *Orchis mascula*. When dried it forms an amylaceous product which was formerly esteemed.

NATURAL ORDER.—ZINGIBERACEÆ.

GRANA PARADISI, Grains of Paradise.—The seeds of *Amomum Melegueta*, from the West Coast of Africa.

Description.—Hard seeds, about 1-10th of an inch in diameter, with a shining reddish-brown surface and variable angular form.

Uses.—In veterinary medicines.

Chemistry.—The chief principle is an essential oil.

CURCUMÆ RHIZOMA, Turmeric.—The dried rhizome of *Curcuma longa*. A native of Southern Asia, and imported from India and the Eastern Archipelago.

Description.—The most common is the long turmeric, about the size of the finger; pointed; and externally of a yellow colour, and marked with transverse annular wrinkles, but it is often found as short turmeric, which is ovate or pear-shaped, much thicker and shorter.

Uses.—As an ingredient in condiments, specially in curry ; in dyeing ; and in testing for alkalies and boracic acid.

Chemistry.—Turmeric contains about 1 per cent. of volatile oil and a colouring matter called *curcumin*, obtained by exhausting the root with benzol, after the essential oil has been distilled off. Its formula is supposed to be $C_{10}H_{10}O_{13}$. Its yellow colour becomes red in the presence of alkalies, and orange, after drying, in the presence of boracic acid.

Tincture of Turmeric.

Turmeric, bruised	1 ounce
Rectified Spirit	6 fluid ounces

Macerate 7 days.

Turmeric Paper (B.P.)

Unsize paper dipped in the above tincture and dried.

CARDAMOMUM, Cardamom.—The dried capsules containing the seeds of *Elettaria Cardamomum*, cultivated in Malabar.

Description.—The capsules are imported entire, as the seeds, which are the real part required, would lose much of their activity if separated before being required

for use. Good cardamomum fruits should be plump and heavy, and should yield seeds to the extent of 3-4ths of their weight. The separated seeds are reddish brown, wrinkled and angular externally, and whitish internally.

Uses.—Similar to ginger, but more aromatic.

Chemistry. — The seeds contain 4 per cent. of volatile oil, 10 per cent. of fixed oil and acrid resin. The taste and odour are due to the volatile oil, which has a sp. gr. of .93 and contains the usual elæopten and stearopten.

Tinctura Cardamomi Composita.

Cardamom Seeds, freed from the Pericarps, and bruised	¼ ounce
Carraway Fruit, bruised.....	¼ ounce
Raisins, freed from their seeds	2 ounces
Cinnamon Bark, bruised	½ ounce
Cochineal, in powder	60 grains
Proof Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

DOSE— $\frac{1}{2}$ to 2 drachms.

Also an ingredient in Extractum Colocynthis Compositum; Pulvis Cinnamomi Compositus (1 in 3):

Pulvis Cretæ Aromaticus (1 in 44); Tinctura Gentianæ Composita; Tinctura Rhei; and Vinum Aloes.

Varieties.—There are three varieties of cardamoms: the best are Malabar shorts. The second are the Madras shortlongs, lighter in colour, and shipped at Madras. The third variety are Aleppy shorts, with a peculiar greenish tint, imported from Calicut. As a rule, shorts are always to be preferred.

Further information respecting cardamoms must be sought for in treatises devoted to the subject. The natural history of this class of fruits has always proved a difficulty to pharmacologists.

ZINGIBER, Ginger.—The scraped and dried rhizome of *Zingiber officinale*, from plants cultivated in the East and West Indies and tropical regions of Asia, Africa, and America.

Description.—In more or less brownish zigzag pieces called “hands” or “races.” Good official races should be scraped, and have a yellowish white but not a chalky appearance on the surface, with a short mealy fracture.

Uses.—Aromatic stimulant and carminative; good in flatulence and to allay the griping of purgatives; when chewed, ginger behaves as a sialagogue.

Dose, 5 to 30 grains.

Chemistry.—According to the older writers, the active principle of ginger is the volatile oil, C_5H_8 ; but Flückiger states that he obtained, by distilling ginger with water, a yellow oil having the sp. gr. .878, and possessing the odour but *not* the pungent taste of ginger. He is of opinion that the acidity of ginger is due to the resin it contains, not yet, however, properly investigated.

Syrupus Zingiberis.

Strong Tincture of Ginger.....	6 fluid drachms
Syrup.....	19 fluid ounces

Dose—1 drachm.

Tinctura Zingiberis.

Ginger Root, in coarse powder.....	2½ ounces
Rectified Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—15 minims to 1 drachm. (7 grains of Ginger in 1 fluid drachm.)

Tinctura Zingiberis Fortior.

Ginger, in fine powder	10 ounces
Rectified Spirit	ad. 20 fluid ounces

Pack the Ginger tightly in a percolator, and pour carefully over it $\frac{1}{2}$ pint of spirit. In 2 hours add more, until 1 pint has passed through.

Dose—5 to 20 minims. (28 grains of Ginger in 1 fluid drachm.)

Also contained in Confectio Opii, Confectio Scammonii, Infusum Sennæ, Pilula Scillæ Composita, Pulvis Cinnamomi Compositus, Pulvis Jalapæ Compositus, Pulvis Opii Compositus, Pulvis Rhei Compositus, Pulvis Scammonii Compositus, Syrupus Rhamni, and Vinum Aloes.

Varieties.—There are four chief varieties of ginger at present found in the market. These are Jamaica, Cochin, Bengal, and African; the Jamaica being the best, Cochin second, and Bengal third of the scraped gingers, while the African is unscraped.

GALANGAL (*Radix Galangæ minoris*) is derived from *Alpinia officinarum*. The drug is an aromatic stimulant, and might take the place of ginger. Its chief consumption is in Russia.

NATURAL ORDER.—MARANTACEÆ.

MARANTÆ AMYLUM, Arrowroot.
—The starch from the tubers of *Maranta arundinacea*, from the West Indies.

Collection.—The tubers (rhizomes) are beaten into a pulp, when one year old, and thrown into water and agitated, and rapidly strained. The starch passes through with the water, and is allowed to dry in the sun. Besides the West Indian arrowroot, East Indian arrowroot is known in commerce, and is usually yielded by *Curcuma augustifolia*.

Maranta arrowroot is of three qualities: namely, Bermuda, St. Vincent and Natal. The first importation of Natal arrowroot was of unexceptionable quality, but less care was bestowed on its subsequent preparation.

Tous les Mois. — The starch of *Canna edulis*, produced in Barbadoes in a similar manner to arrowroot. It is an exceedingly pure starch, and an article of great consumption.

Canna speciosa is said to produce African turmeric.

NATURAL ORDER.—IRIDACEÆ.

Herbs with bulbs, corms or rhizomes ; leaves equitant ; perianth 3 partite and superior ; stamens 3 on the outer segments of the perianth, and with extrorse anthers ; ovary inferior 3-celled, with axile placentation, and a 3-lobed and often petaloid stigma.

CROCUS, Saffron.—The dried stigma and part of the style of *Crocus sativus*, imported from Spain, France and Italy.

Botanical Source. — An annual, very like the ordinary crocus of our gardens ; with a pale, purple perianth, and a specially large, somewhat petaloid stigma.

Description.—A mass of deep red, shrub-like styles, each terminated by 3 long orange brown stigmas, expanded towards their ends. Rubbed on damp paper it leaves an orange colour, but pressed between folds of filtering paper it leaves no oily stain.

Chemistry.—The colouring matter of saffron is

supposed to be a substance called polychroit, which is a viscid, deliquescent body ; readily soluble in water, but sparingly soluble in alcohol, and insoluble in ether. It is a glucoside, and by the action of a dilute acid is decomposed into *crocin*, an essential oil and glucose. The red colour of saffron is changed to blue, violet, and brown by strong sulphuric, and to green, yellow and brown by strong nitric acid.*

Tinctura Croci.

Saffron	1 ounce
Proof Spirit	ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Saffron is also contained in Decoctum Aloes Compositum ; Pilula Aloes et Myrrhæ ; Pulvis Cretæ Aromaticus ; Tinctura Cinchonæ Composita ; Tinctura Opii Ammoniata ; and Tinctura Rhei.

Varieties.—The best is the hay saffron, and the inferior is the cake saffron, which is much adulterated by the florets of safflower, *Carthamus tinctorius*, which are easily recognisable after steeping in warm water.

* A very dilute infusion of saffron is decolorised by diluted hydrochloric acid. Subsequently boiled in the presence of sugar, the yellow tint will be restored and a beautiful red fluorescence make its appearance on the surface of the liquid.—(Stoddart.)

The expanded stigma of the saffron-crocus exhibits a characteristic and unmistakable form.

Adulterations.—The hay saffron is frequently made heavy by dipping it in coloured chalk, which may be detected by stirring with a very small quantity of water.* It has also been said to be mixed with dried shreds of beef and the dried stamens of the plant. Saffron is also found which has been exhausted of a portion of its colouring matter. This can be discovered by steeping in water and observing the amount of colour.

SUB-DIVISION II.—HYPOGYNÆ.

Having the perianth inferior and the ovary superior.

NATURAL ORDER.—LILIACEÆ.

Herbs, shrubs or trees with sessile or sheathing leaves ; perianth 6 partite and inferior ; stamens 6 ; anthers introrse ; ovary, superior 3-celled with axile placentæ ; fruit with loculicidal dehiscence or succulent.

* A white powder will instantly separate, causing the water to appear turbid ; and if a drop of hydrochloric acid be now added, a brisk effervescence will take place.—(*Hanbury.*)

ALOE BARBADENSIS, Barbadoes Aloes. — The inspissated juice of the leaves of *Aloe vulgaris*.

Botanical Source.—The *A. vulgaris* is a succulent plant, with lanceolate and curved leaves, having large reddish spines and bright yellow flowers. Aloes are cultivated in Barbadoes, being planted in rows, carefully attended to, and manured.

Collection.—The leaves are cut off close to the plant in March or April, during the heat of the day, and instantly thrown into a trough, where the juice is allowed to exude spontaneously. The juice is then exposed to the sun until sufficiently inspissated, or it may be rapidly boiled down in copper vessels.

Description.—Good samples of Barbadoes aloes should be hard, dry, and have a deep chocolate-brown colour, with a clean, dull, conchoidal fracture. When breathed upon, it exhales a characteristic odour. It is usually imported in gourds.

Uses.—In full doses cathartic, especially on the large intestines. It is slow in operation and useful as a

purgative for women, being to some extent a stimulant of the action of the uterus, but it must not be given when the patient suffers from hæmorrhoids. Used in the form of enema it is very efficient as an anthelmintic in thread worms (*ascaris vermicularis*). A point in the administration of aloes, is, that if 1 to 2 grains does not act, a better effect cannot be produced by increasing the dose.

Chemistry.—The purgative constituent is supposed to be *aloin* $C_{34}H_{36}O_{14}.H_2O$, which is accompanied by a little amorphous aloin, volatile oil and resin. Aloin is prepared by boiling 1 lb. of aloes with 1 gallon of water slightly acidulated with sulphuric acid. The solution thus obtained, having been allowed to stand for 24 hours, is poured off from the fixed resinous matter, and quickly concentrated by evaporation to about 30 ounces, and set to crystallise. The crystals of aloin thus obtained are purified by re-crystallisation from very dilute spirit.

Each of the commercial varieties of aloes seems to contain a peculiar aloin of its own. Thus barbaloin is in prismatic crystals, soluble in water, alcohol and ether, and turns red when treated with nitric acid, the colour rapidly fading. It melts at 300° F., and when subjected

to oxidation with nitric acid it yields both picric and chrysamic acids. When mixed with a drop of sulphuric acid, and then a rod moistened with nitric acid placed in contact, no blue colour is developed.

Socaloin, obtained from Socotrine aloes, $C_{30}H_{32}O_{14}$, is much less soluble in ether than *barbaloin*, and does not give a brilliant red with nitric acid.

Nataloin, $C_{25}H_{28}O_{11}$, crystallises in plates, but is only slightly soluble in water and alcohol. With nitric acid it produces a strong red colour which does not fade like that from barbaloin, and when treated with a drop of sulphuric and then nitric acid it yields picric and oxalic acids, but not chrysamic acid.

Enema Aloes.

Aloes	40 grains
Carbonate of Potash	15 grains
Mucilage of Starch	10 fluid ounces

Mix and rub together.

Extractum Aloes Barbadosis.

Barbadoes aloes purified by solution in water, straining, and subsequent evaporation.

(100 parts yield 75 parts of extract.)

DOSE—2 to 6 grains.

Pilula Aloes Barbadensis.

Barbadoes Aloes, in powder	2 ounces
Hard Soap, in powder	1 ounce
Oil of Carraway	1 fluid drachm
Confection of Roses	1 ounce

Dose—5 to 10 grains. (50 per cent. stronger than P.L.)

Pilula Aloes et Ferri.

Sulphate of Iron	1½ ounce
Barbadoes Aloes, in powder	2 ounces
Compound Powder of Cinnamon.....	3 ounces
Confection of Roses	4 ounces

Dose—5 to 10 grains. (1½ grains of Iron and 4 grains of Aloes in 10½ grains.)

Also contained in *Pilula Cambogiæ Composita* ;
Pilula Colocynthis Composita ; *Pilula Colocynthis*
et Hyoscyami.

ALOE SOCOTRINA, Socotrine Aloes.

—The inspissated juice of the leaf of *Aloe Socotrina* and allied species imported from Bombay, to which it has been carried from Socotra to Zanzibar.

Description.—It is imported in kegs in tin-lined boxes. When of a fine quality it is a dark reddish brown, and when breathed upon it emits a peculiar

odour. Very frequently it is opaque and liver-brown in colour, with a dull waxy fracture and not very fragrant, when it receives the name of "hepatic aloes."

Uses.—Similar to Barbadoes aloes.

Decoctum Aloes Compositum. (*Baume de Vie.*)

Extract of Socotrine Aloes.....	120 grains
Myrrh, bruised.....	90 grains
Saffron, chopped fine	90 grains
Carbonate of Potash	60 grains
Extract of Liquorice	1 ounce
Compound Tincture of Cardamoms	8 fluid ounces
Distilled Water	22 fluid ounces

Boil 5 minutes, product 30 ounces.

Contains 4 grains Extract of Aloes in 1 fluid ounce.

Incompatibles—Acids; acidulous, earthy, and metallic salts; and all substances which are decomposed by potassa carbonas.

Dose— $\frac{1}{2}$ to 2 ounces.

Extractum Aloes Socotrinæ. (Ext. Aloes P.L.)

Socotrine Aloes purified.

(100 parts yield 50 parts of extract.)

Dose—2 to 6 grains.

Pilula Aloes et Asafœtidæ.

Socotrine Aloes, in powder	1 ounce
Asafœtida	1 ounce
Hard Soap, in powder	1 ounce
Confection of Roses	1 ounce

Dose—5 to 10 grains.

Pilula Aloes et Myrrhæ.

Socotrine Aloes	2 ounces
Myrrh	1 ounce
Saffron, dried	$\frac{1}{2}$ ounce
Confection of Roses	$2\frac{1}{2}$ ounces

Dose—5 to 10 grains. (2 grains of Aloes and 1 grain of Myrrh in 6 grains.)

Pilula Aloes Socotrinæ.

Socotrine Aloes, in powder	2 ounces
Hard Soap, in powder	1 ounce
Volatile Oil of Nutmeg	1 fluid drachm
Confection of Roses	1 ounce

Dose—5 to 10 grains. (5 grains of Aloes in 10 grains.)

Tinctura Aloes.

Socotrine Aloes, in coarse powder	$\frac{1}{2}$ ounce
Extract of Liquorice	$1\frac{1}{2}$ ounce
Proof Spirit	20 fluid ounces

Macerate 7 days, and then filter.

Dose—1 to 2 drachms. (11 grains Aloes in 1 fluid ounce.)

Vinum Aloes.

Socotrine Aloes	$1\frac{1}{2}$ ounce
Cardamom Seeds, bruised.....	80 grains
Ginger, in coarse powder	80 grains
Sherry	40 fluid ounces

Digest 7 days, and filter.

Dose—1 to 2 drachms. (2 grains of Aloes in 1 fluid drachm.)

Socotrine Aloes is also contained in Extractum Colocynthidis Compositum, Pilula Rhei Composita (1 in 6 nearly), Tinctura Benzoini Composita (160 grains to 1 pint).

ALOE CAPENSIS, Cape Aloes.—The inspissated juice of the leaf of *Aloe spicata* and other species, imported from the Cape of Good Hope.

Description.—This aloes is distinguished by its brilliant glossy fracture and characteristic odour. When viewed by reflected light it has a greenish appearance, and when splinters are examined by transmitted light they are found to be amber-coloured.

Natal Aloes is a variety of aloes imported from Natal since 1870 to a considerable extent. It is more like hepatic aloes in appearance, and contains *Nataloin*, already described.

SCILLA, Squill. — The sliced and dried bulb of *Urginea Scilla* (*Scilla maritima*), from the Mediterranean coasts.

Botanical Source.—The squill is endogenous, being a pear-shaped bulb, usually weighing about 4 lbs. It is

tunicated, and has reddish brown outer scales and pale rose fleshy inner scales.

Collection.—The bulbs are dug up in August, freed from their dry outer scales, cut across into slices, and dried in the sun.

Description.—In dried slices, white or yellowish white, and slightly translucent, something like tragacanth. When exposed to damp the slices become flexible, but when dried they are brittle and readily powdered. The powder is apt to absorb moisture and concrete into an intractable hard mass.

Uses.—Expectorant, emetic, and diuretic; used in chronic pulmonary complaints, whooping cough, and in dropsies, when not accompanied by inflammatory symptoms.

Chemistry.—Squills contain a large quantity of mucilage, which may be precipitated by the addition of alcohol to a watery infusion. If, after thus separating the mucilage, the alcohol be evaporated and tannic acid added, a bitter body will be precipitated, which has been called scillitin or skulëin, but which has not been properly investigated.

Acetum Scillæ.

Squill, bruised	2½ ounces
Dilute Acetic Acid	20 fluid ounces
Proof Spirit	1½ fluid ounce

Macerate 7 days; add Spirit, and filter.

Dose—15 to 40 minims.

Oxymel Scillæ.

Vinegar of Squill.....	20 fluid ounces
Clarified Honey	32 ounces

Evaporate in water-bath to sp. gr. 1.32.

Dose—½ to 1 drachm.

Pilula Ipecacuanhæ c. Scillâ. (Vide p. 53.)

Pilula Scillæ Composita.

Squill, in fine powder.....	1¼ ounce
Ginger, in fine powder	1 ounce
Ammoniacum, in fine powder	1 ounce
Hard Soap, in powder	1 ounce
Treacle, by weight	2 ounces or q.s.

Dose—5 to 10 grains. (1 grain of Squill in 5 grains.)

Syrupus Scillæ.

Vinegar of Squill.....	20 fluid ounces
Refined Sugar	40 ounces

Product, 60 ounces by weight.

Dose—½ to 1 drachm.

Tinctura Scillæ.

Squill, bruised	2½ ounces
Proof Spirit	20 fluid ounces

Macerate 48 hours, then percolate.

Dose—10 to 30 minims. (1 grain of Squill in 9 minims.)

NATURAL ORDER.—MELANTHACEÆ.

Similar to Liliaceæ, but having *extrorse* anthers, and the fruit commonly dehiscing septicidally.

SABADILLA, Cevadilla.—The dried fruit of *Asagrea officinalis*, imported from Mexico.

Description.—Each fruit consists of three brown papery follicles, about $\frac{1}{2}$ inch long, containing each from 1 to 3 seeds.

Chemistry.—It is used for extracting the alkaloid veratria.

Veratria, $C_{32}H_{52}N_2O_8$, is prepared as follows:—Macerate 2 pounds of Cevadilla with half its weight of boiling distilled water in a covered vessel for 24 hours; remove the cevadilla; squeeze it and dry it thoroughly with a gentle heat; beat it now in a mortar, and separate the seeds from the capsules by brisk agitation in a deep narrow vessel, or by winnowing it gently on the table with a sheet of paper; grind the seeds in a

coffee-mill, and form them into a thick paste with rectified spirit. Pack this firmly in a percolator, and pass rectified spirit through it until the fluid ceases to be coloured; concentrate the spirituous solution by distillation so long as no deposit forms, and pour the residue while hot into 12 times its volume of cold water. Filter through calico, and wash the residue on the filter with distilled water until the fluid ceases to precipitate with ammonia. To the united filtered liquids add solution of ammonia in slight excess; collect the precipitate on a filter, and wash it with water until the fluid passes colourless. Diffuse the moist precipitate through 12 ounces of water, and add gradually, with diligent stirring, sufficient hydrochloric acid to make the fluid feebly but persistently acid. Then add 60 grains of purified animal charcoal; digest at a gentle heat for 20 minutes; filter, and allow the liquid to cool. Add ammonia in slight excess; collect the precipitate on a filter, and wash it with cold water till the washings cease to be affected by nitrate of silver acidulated with nitric acid. Lastly, dry the precipitate—first by imbibition with filtering paper, and then by the application of a gentle heat.

The *Veratria* is, in this process, extracted in its

natural state as a gallate of veratria, together with resin and alkaloid matter. When the concentrated tincture is poured into water, the resin is precipitated. The addition of ammonia then causes precipitation of the crude veratria, which is further purified by solution in acid and decolorisation by animal charcoal, and reprecipitation by ammonia.

As thus prepared veratria is pale grey and amorphous, the minutest quantity producing instant irritation if breathed into the nostrils. It is insoluble in water, but soluble in spirit, ether, and dilute acids. With concentrated sulphuric acid it become first yellow, then red, and finally violet.

Veratria is a violent irritant poison, which has been used as a substitute for colchicum in gout. Externally, in the form of ointment, it is rubefacient and errhine; and sometimes used in neuralgia and rheumatism.

Unguentum Veratriæ.

Veratria.....	8 grains
Prepared Lard.....	1 ounce
Olive Oil	$\frac{1}{2}$ fluid drachm

COLCHICI CORMUS.—The fresh corm, and the same stripped of its coats, sliced transversely

and dried, at a temperature not exceeding 150° F. of *Colchicum autumnale*, indigenous.

Botanical Source.—The colchicum is a herb with leaves broadly lanceolate, and about 1 foot long, appearing in spring. In early summer the leaves fall off, and in autumn the flowers appear in the form of pale lilac perianths, which die away and form so-called capsules, ripening in the following spring.

Collection.—The corm should be dug up and brought to market in July, during the period of the decay of the leaf and the production of the flower.

Description.—When dried in flat pieces about 1-10th of an inch thick, white, starchy and more or less reniform in shape.

Uses.—Colchicum is a sedative and irritant; in small doses nauseant, cathartic, diuretic, frequently given in gout and acute rheumatism; in large doses an irritant poison.

Chemistry.—The active constituent is an alkaloid called *colchicia*, $C_{17}H_{19}NO_5$. It is soluble in water and alcohol, and is closely allied to veratria, but does not cause sneezing. It turns deep violet with nitric acid.

Extractum Colchici.

The juice of the fresh *Colchicum* corms, evaporated to an extract, after the feculence has deposited.

(100 lbs. of corms yield 4 lbs. of extract.)

Dose— $\frac{1}{2}$ to 2 grains.

Extractum Colchici Aceticum.

Fresh *Colchicum* Corms, deprived of their

coats 7 lbs.

Acetic Acid 6 fluid ounces

(100 lbs. of Corms yield 5½ lbs. of extract.)

Dose— $\frac{1}{2}$ to 2 grains.

Vinum Colchici.

Colchicum Corm, dried, sliced and bruised 4 ounces

Sherryad. 20 fluid ounces

(Macerate 7 days, press, and strain through calico.)

Dose—10 to 30 minims. (1 grain of *Colchicum* Corm in 9 minims.)

COLCHICI SEMINA, *Colchicum* Seeds.—The ripe seeds of the plant already described. They are small and spherical, about 1-10th of an inch in diameter, hard and difficult to powder, reddish brown in colour, but dull and wrinkled.

Chemistry.—Like the *Colchicum* corm.

Antidotes.—Similar to *veratria*.

Tinctura Colchici Seminum.

Colchicum Seeds, bruised 2½ ounces
 Proof Spirit.....ad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—10 to 30 minims. (1 grain of Colchicum Seeds in 9 minims.)

VERATRI ALBI RADIX, White Hellebore Root. — The dried acrid rhizome of *Veratrum album*, from the Alps, Pyrenees, and other mountainous districts of Central and Southern Europe.

Botanical Source. — The *V. album* grows from 1½ to 3 feet high, has a præmorse root stalk, and elliptic and plicate leaves, pubescent beneath. The flowers are in paniculate racemes of a yellowish white colour, greenish at the base.

Description.—A cylindrical root of a dull earthy black, the top being crowned with the bases of leaves, and the lower part much beset by the scars of rootlets. Internally it is nearly colourless, and a section shows a bright white ring surrounding a spongy buff coloured medullium.

Uses.—A hydragogue, cathartic, and in over doses

produces vomiting and purging. It is now chiefly used to destroy pediculi.

Chemistry.—Contains the alkaloid veratria, and has also been said to contain traces of three other allied alkaloids, viz., sabadilline, colchicine and jervine.

VERATRI VIRIDIS RADIX,
Green Hellebore Root. — The dried rhizome of *Veratrum viride*, collected in autumn in the United States and Canada.

Botanical Source.—The green hellebore is an annual herb with bright green leaves, oval-acuminate in shape, pubescent, plated and amplexicaul. The inflorescence is a panicle of greenish yellow flowers with a six-parted perianth.

Description.—It is very similar to white hellebore, but easily distinguishable by being either sliced or cut lengthwise before drying.

Uses.—Similar to white hellebore.

Chemistry. — Contains the alkaloids veratria and jervia, and is in most respects similar to the white hellebore.

Tinctura Veratri Viridis.

Green Hellebore Root, in coarse powder 4 ounces
 Rectified Spiritad. 20 fluid ounces

Macerate 48 hours, then percolate.

Dose—5 to 20 minims. (1 grain of Hellebore Root in
 5½ minims.)

NATURAL ORDER.—PALMACEÆ.

ARECA, Areca or Betel Nut.—The seeds of *Areca Catechu*, from Ceylon and the Malayan Archipelago.

Botanical Source.—The areca palm is a tall tree, frequently 50 feet in height.

Description.—Similar to a nutmeg, but rather larger, and often enclosed in a peculiar fibrous pericarp.

Uses. — An astringent and anthelmintic for the expulsion of *tænia*. Charcoal prepared from the nuts is esteemed as a dentifrice.

Dose, 10 to 15 grains for diarrhœa, and ½ to ¾ ounce for tapeworm.

Chemistry.—Its active principle is tannic acid.

SANGUIS DRACONIS, 'Dragon's Blood.—A resinous substance exuding from the fruit of *Calamus Draco*, chiefly from Singapore and Batavia.

Description.—It is a hard deep red substance, almost entirely soluble in spirit of wine. The finest quality is in sticks neatly wrapped in palm leaf, and tied with transverse bands of dried grass. The interior is in lumps containing remains of the fruit, of which only about two-thirds are soluble in spirit of wine.

Uses.—As a colouring matter for plasters, tooth powders, and varnishes.

Chemistry.—The resin of dragon's blood has been stated to have the formula of $C_{20}H_{20}O_4$.

Specially Distinctive Tests. — 1. Completely soluble in chloroform and ether, the latter solution giving no turbidity with alcohol.

2. Soluble in alcohol, and the addition of alcoholic ferric chloride (1 in 10) gives a greenish brown; but alcoholic plumbic acetate gives no precipitate, while ammonium hydrate gives a turbid mixture.

The dragon's blood from *Pterocarpus Draco*

(Leguminosæ) is not so soluble in chloroform, but readily dissolves in ether; the addition of ammonium hydrate to the red alcoholic solution gives no turbidity.

SAGO.—The starch obtained from the interior of the stem of various species of *Sagus*, growing in swampy districts of the Indian Archipelago.

NATURAL ORDER.—ACORACEÆ.

[AROIDEÆ.]

CALAMI AROMATICI RHIZOMA,
Sweet Flag.—The rhizome of *Acorus Calamus*, indigenous.

Description.—As usually found it is rough and shrunken, varying in colour from orange-brown to dark brown externally, with a dull brown spongy interior. On the upper surface it is marked with scars of leaves, and on the end with zigzag lines of little elevated dot-like rings.

Uses.—An aromatic stimulant, and believed by some to be anti-spasmodic.

Chemistry.—It yields an essential oil, isomeric with turpentine and a bitter principle called *acorin*.

SUB-CLASS III.—GLUMACEÆ.

Plants having the flowers enclosed in glumes and paleæ.

NATURAL ORDER.—GRAMINACEÆ.

Herbs, shrubs, or arborescent plants with hollow, round and jointed stem; leaves alternate with ligules and split sheaths; inflorescence in locustæ; flowers achlamydeous, furnished with paleæ and glumes; stamens 1—6, with versatile anthers, and ovary superior 1-celled, with a feathery stigma; fruit, a caryopsis.

HORDEUM DECORTICATUM,
Pearl Barley. — The husked seeds of *Hordeum distichon*, cultivated in Britain.

Uses.—Emollient, demulcent and nutritive.

Chemistry.—Similar to wheat flour, but not quite so rich in gluten.

Decoctum Hordei.

Pearl Barley.....	2 ounces
Distilled Water	1½ pint

Wash in cold water, boil 20 minutes, and strain.

TRITICI FARINA, Wheat Flour.—

The flour ground from the fruits of *Triticum vulgare*, which have been separated from the perianth by thrashing.

Chemistry.— It contains 68 per cent. of starch, 10 per cent. of gluten, 5 per cent. of sugar, 4 per cent. of gum, and the remainder water and mineral matters, consisting chiefly of phosphates. When some flour is placed on a cloth and washed with water, the starch passes through, the gluten remaining on the cloth. Gluten differs from starch in containing nitrogen, and it therefore is considered to be a flesh-forming article of food, in contradistinction to starch, sugar or fat, which are only heat-producing bodies.

AMYLUM, Starch, $C_6H_{10}O_5$.—Procured from the above. In white columnar masses. Triturated with a little cold distilled water it is neither acid nor alkaline to test paper; the filtered liquid does not become blue

on the addition of iodine in solution; mixed with boiling water and cooled it gives a deep blue colour with iodine, of amidin iodide. (See Author's Pharmaceutical Chemistry.)

Glycerinum Amyli. (1 in 11, by weight.)

Starch	1 ounce
Glycerin	8 fluid ounces

Rub them together until they are intimately mixed, then transfer the mixture to a porcelain dish, and apply a heat gradually raised to 240° F. It should be constantly stirred until the starch particles are completely broken, and a translucent jelly is formed. The original compound was called Plasma, before it became an official preparation.

Mucilago Amyli. (12 grains to 1 fluid ounce.)

Triturate the starch with water, gradually added, and boil for a few minutes, constantly stirring.

Starch is also contained in Pulvis Tragacanthæ Compositus (1 in 6).

SACCHARUM PURIFICATUM. —

(See Author's Pharmaceutical Chemistry.)

SUB-KINGDOM II.—CRYPTOGAMIA.

FLOWERLESS PLANTS PROPAGATED BY SPORES.

CLASS III.—ACOTYLEDONES.

Plants having no seeds but propagated by spores ; germination heterorhizal ; stem (when present) acrogenous ; leaves (when present) furcate in venation ; flowers none.

SUB-CLASS I.—ACROGENÆ.

Plants with stems and leaves distinguishable and possessing stomata.

NATURAL ORDER.—FILICES.

Herbs with rhizomes or arborescent plants with acrogenous stems ; leaves (fronds) with furcate venation : fructification in *sori*, usually on the under surface or margins of the frond ; spores enclosed in *sporangia*.

FILIX MAS, Male Fern. — The dried rhizome, with the bases of the foot stalks and portions of the root fibres attached of *Aspidium Filix Mas*, collected in summer from indigenous plants.

Uses.—For exciting muscular action in the bowels, and therefore useful for expelling worms. The dose in powder is from 50 to 100 grains.

Chemistry.—It contains a little volatile oil, 6·9 per cent. of a green fixed oil, 4·1 per cent. of resin, together with tannin, starch, sugar and gum. When treated with ether it yields 8 per cent. of an extract containing a crystalline substance called *flicic* acid, $C_{14}H_{18}O_5$, which, when fused with potash, is resolved into butyric acid and another substance. This is probably the active constituent.

Extractum Filicis Liquidum.

Fern root, in coarse powder	32 ounces
Ether	4 pints, or q.s.

Exhaust by percolation, and evaporate to an oily extract.

Dose—15 to 30 minims.

NATURAL ORDER.—LYCOPODIACEÆ.

LYCOPODIUM.—The spores of *Lycopodium clavatum*, or club-moss, from indigenous plants, but chiefly imported from Central and Northern Europe.

Botanical Source.—The *L. clavatum* is a moss with a long creeping stem; leaves linear-cylindrical and imbricate, terminating in hair-like processes; fructification sessile, in the axils of the leaves; spores tetrahedral, with 4 facets, and covered with hexagonal network.

Description.—A straw-yellow mobile and inodorous powder, which, although specifically heavier than water, floats on its surface, owing to occlusion of air on the netted surface of the grains. Very inflammable.

Uses.—For dusting scalded surfaces, and for preventing the adhesion of soft pills. Reputed among herbalists as violently purgative and emmenagogue.

Chemistry.—Contains, when pure, under 5 per cent. of water, and yields an ash of 4 per cent., the chief constituent of which is aluminium phosphate. Ether extracts 48 per cent. of a fixed oil. It is adulterated sometimes by starch and magnesia; test, the microscope.

SUB-CLASS II.—THALLOGENÆ.

Plants without distinction of stem and leaf and destitute of stomata.

NATURAL ORDER.—LICHENES.

CETRARIA (B.P.), Iceland Moss.—

The entire plant of *Cetraria islandica*, from the North of Europe.

Botanical Source.—The *C. islandica* is an epiphyte with an erect, smooth, leathery and dry thallus, from 2 to 4 inches high: fructification in rusty brown rounded apothecia on the apices of the thallus.

Description.—Has the characters named above, and when dry is brownish white above but lighter in colour beneath.

Uses. — Demulcent, nutritive and slightly tonic; useful in advanced phthisis.

Chemistry. — By first extracting with boiling rectified spirit, and then boiling the residue in water, concentrating and precipitating by spirit of wine, about

70 per cent. of a peculiar body called *lichenin*, or lichen-starch is obtained. Its formula is $C_{12}H_{20}O_{10}$,* and it may be regarded as a polymer of starch and cellulose, partaking of certain qualities of both, and being soluble in ammoniacal oxide of copper as well as tinged blue by iodine. By boiling with dilute acid it is converted into dextrin and sugar. The bitter principle is *Cetraric acid*, $C_{18}H_{16}O_8$, of which it contains 3 per cent., and there is also a little fat, fumaric acid and sugar. The cetraric acid may be extracted by a weak solution of potassium carbonate, and this yields, on evaporation, a yellow bitter salt.

Decoctum Cetrariæ.

Iceland Moss	1 ounce
Distilled Water	1 pint

Boil 10 minutes in a covered vessel, and strain while hot, product, 20 fluid ounces.

Incompatibles—The alkalies and their carbonates; most metallic salts, and iodine.

LITMUS (B.P.), Lacmus.—A blue pigment prepared from various species of *Rocella*, and imported from Holland.

* This formula is wrong in the books, it should be C_{12} , as here written.

Botanical Sources.—The two chief species used are the *R. tinctoria* and *R. fuciformis*, the former having a thallus branching and cylindrical in form, and the latter possessing a flat one.

Manufacture.—The lichens are exposed to the air after grinding, in a solution of potassium and ammonium carbonates. They are then treated with ammonium hydrate, and the whole digested for some hours more, and decanted. After exposure to the air for several weeks, this solution is evaporated and made into cakes with plaster of Paris and chalk.

Description.—In small light cubical masses about $\frac{1}{4}$ inch square, having an indigo-blue colour, and an odour something like that of violets. It yields its blue colour both to water and spirit.

Uses.—Acids and acid salts turn blue litmus paper red, while alkalies or any alkaline salt, such as sodæ phosphas, or plumbi subacetat, turn red litmus paper blue.

Chemistry.—The essential constituent of the blue colour is *orcein*, $C_7H_7NO_3$, which is produced by the joint action of oxygen and ammonia upon *orcin*, $C_7H_8O_2$,

The latter substance, even, is not contained in the litmus, but is the product of the action of potassium carbonate upon certain acids existing in the litmus, the chief of which are *orsellic* acid, $C_8H_8O_4$, *erythric* acid, $C_{20}H_{22}O_{10}$, and *evernic* acid, $C_{17}H_{16}O_7$.

Tincture of Litmus.

1 in 10 of proof spirit. Macerate for 2 days, and filter.

Blue Litmus Paper.

By steeping slips of unsized paper in the tincture, and drying by exposure to the air.

Red Litmus Paper.

By reddening the tincture with a minute quantity of diluted sulphuric acid, and steeping paper into the solution.

Instead of using two kinds of paper, it is preferable to prepare what is called "neutral" litmus paper of a pale purple colour, which will show both the changes to red and to blue. This is effected by boiling $\frac{1}{2}$ ounce of litmus in 4 ounces of water for 20 minutes, filtering, making up with water to exactly 4 ounces, and dividing into two equal parts. To one part *exceedingly* diluted sulphuric acid is added, drop by drop, till it turns red; and the other portion is then poured into it, and slips of unsized paper passed slowly through the liquid and hung up to dry.

NATURAL ORDER.—FUNGI.

Cellular plants growing in or upon living or decaying organic substances, and nourished through their vegetative structure called the mycelium (spawn); fructification various and produced in the air.

ERGOTA (B.P.), Ergot of Rye. — The compact mycelium or spawn (sclerotium) of *Claviceps purpurea*, produced within the paleæ of *Secale cereale*, in Southern Russia and Spain, but chiefly in Galicia.

Botanical Source.—The first sign of the grain of rye being attacked by ergot is, according to Tulasne, the appearance of a yellow mucus called honeydew. This sweet exudation attracts many insects which thus help to propagate the infection to other grains. The dew soon disappears and leaves the young ovaries covered by a white spongy mycelium of the fungus which penetrates the entire grain and develops *hyphæ*, or thread-like cells, covered externally by *basidia*. The

epicarp and embryo of the caryopsis are especially attacked and its growth is stopped, while at its base is formed by tumefaction and compaction of the mycelium, a mass gradually increasing in size, which is the *sclerotium*. This in time separates from the first mycelium by the drying up of the latter, and pushes the surviving portion of the apex of the caryopsis up so that it is visible above the paleæ. In collection this portion usually drops off, and is not found in commercial ergot.

If a portion of the ergot be placed on damp earth it develops a little *stipe* and *pileus* and all the other characteristics of a minute mushroom; but this development does not take place on the grain, and unless the *sclerotium* falls on suitable places its vitality for propagation only appears to last one season.

Description. — In violet-brown elongated masses about 1 inch long, obtuse at the ends, curved, and sub-triangular, with a longitudinal furrow on the concave side. It is pinkish within and breaks with a short fracture, and gives a faint but peculiar odour, which is much increased by rubbing with potassium hydrate.

Uses.—To assist parturition by causing, usually within 20 minutes after its administration, rapid and violent uterine contractions which continue without intermission until the birth is completed and the placenta expelled. It is also used in menorrhagia in doses up to 15 grains. The dose in parturition is up to 30 grains. It is an aid to midwifery which should only be given under certain circumstances and with extreme discretion, and under the advice of an experienced practitioner.

Chemistry.—Ergot contains 30 per cent. of a non-drying oil which is a mixture of glycerides of oleic and palmitic acids, with several glycerides of soluble and volatile fatty acids, such as butyric, lactic, acetic and formic. It also yields 7 per cent. of resin, with traces of cholesterin, mycose (a peculiar sugar), and some soluble albumen. The colouring matter is red and insoluble in ether and alcohol, but soluble in alcoholic ammonium hydrate.

The real active principle appears to be a base called *ecbolia*, which exists to the extent of .16 per cent. combined with ergotic acid. Another, but apparently inert base called *ergotine*, $C_{50}H_{52}N_2O_3$, is

also present to the amount of .12 per cent. These bases, after mutual extraction in the ordinary manner suitable for alkaloids, may be separated by mercuric chloride, which precipitates only *ecbolia*.

Extractum Ergotæ Liquidum.

Ergot, in coarse powder	16 ounces
Ether..... (should be 40 fluid ounces)	20 fluid ounces
Rectified Spirit	8 fluid ounces
Distilled Water	3½ pints

Product, 16 ounces.

The Ether is merely employed to separate the inert oil.

(1 drachm is equivalent to 1 drachm of Ergot.)

DOSE—10 to 30 minims.

Infusum Ergotæ.

Ergot, in coarse powder.....	¼ ounce
Boiling Distilled Water.....	10 fluid ounces

Infuse in a covered vessel for ½ hour, and strain.

Incompatibles—Astringents, metallic salts.

DOSE—1 to 2 ounces. (About 11 grains of Ergot in 1 fluid ounce.)

Tinctura Ergotæ.

Ergot, in coarse powder	5 ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 48 hours, then percolate.

DOSE—10 minims to 1 drachm. (1 grain of Ergot in 4½ minims.)

NATURAL ORDER.—ALGÆ.

Cellular plants, growing in fresh or salt water, or in moist situations; thallus branched or filamentous; fructification various or wanting; reproduction by spores, tetraspores, and antheridia, or by simple conjugation and cell division.

CHONDRUS CRISPUS, Carragheen, *Irish Moss*.—The entire alga bleached and dried by exposure to the sun, chiefly on the West Coast of Ireland.

Description.—A whitish dichotomously-cleft thallus, with cuneate segments; having here and there elliptical *sori* embedded in the thallus (distinction from Iceland moss).

Uses.—Demulcent, and to a small extent nutritive and alterative. Half an ounce, steeped in water and boiled in a pint of milk, makes a very firm jelly for invalids.

Chemistry.—Irish Moss consists chiefly of mucilage, which, although it contains no starch, is yet slightly coloured blue by iodine. Ash 15 per cent., with traces of iodides and bromides.

GRACILARIA LICHENOIDES,
Ceylon Moss.—A decoction is used as a demulcent.

MATERIA MEDICA
OF THE
ANIMAL KINGDOM.





SUB-KINGDOM I.—PROTOZOA.

(πρῶτος, first; ζῶον, animal.)

Size minute, composed of sarcode; the body not showing any differentiation of parts, and the nervous system absent.

CLASS.—RHIZOPODA.

(ῥίζα, a root; ποὺς, ποδός, a foot; so-called because the pseudopodia spread out like roots.)

Simple or compound animals, possessing the power of emitting pseudopodia.

ORDER.—SPONGIDA.

SPONGIA OFFICINALIS, Sponge.—

The skeleton of *Spongia officinalis*, imported from the Mediterranean and Red Seas, and from the West Indies.

The sponges were placed, by Grant, in a class named **Porifera**.

Varieties.—The finest are Turkey sponges, and the inferior the West Indian; all contain sand and minute shells, which have to be removed by washing before use.

Uses. — Domestic. Medically, calcined sponge, *Spongia usta*, was used for the cure of glandular swellings, probably on account of the sodium iodide and magnesium bromide it contained.

Sponge tents, being compressed sponge saturated with wax, were formerly employed to dilate contracted canals, chiefly the cervix uteri.

Chemistry.—Sponge tissue is chiefly gelatin and coagulated albumen, with a good deal of mineral matter, especially silicates, carbonates, phosphates and iodides.

SUB-KINGDOM II. — CÆLENTERATA

AND

SUB-KINGDOM III.—ANNULOIDA,

Not important to Materia Medica.

SUB-KINGDOM IV.—ANNULOSA.

[RINGED ANIMALS.]

Animals possessing a segmented body arranged along a longitudinal axis. The nervous system consists of a double chain of ganglia situated on the neutral surface of the body, and transversed anteriorly by the œsophagus.

DIVISION I.—ANARTHROPODA.

(Without joints—*α*, without; *ἄρθρον*, a joint; *πούς*, a foot.)

Locomotive appendages not distinctly jointed or articulated to the body.

CLASS.—ANNELIDA.

External surface of the body clearly segmented. The ventral nerve cord ganglionated.

ORDER.—HIRUDINEA.

The sexes are united, and the young do not pass through any metamorphosis. The posterior, or both ends of the body, possess a locomotive and adhesive sucker. Foot tubercles and bristles absent.

HIRUDO (B.P.), the Leech.—Official leeches are either the speckled variety, *Sanguisuga medicinalis*, or the green leech, *S. officinalis*, collected in Spain, France, Italy and Hungary. Leeches are androgynous, each possessing male and female organs. They propagate by eggs.

Description. — *S. medicinalis*. Back with six longitudinal reddish bands, on a greenish brown ground; belly yellowish green with many black spots and yellowish intervening spaces; rings varying from 93 to 108; teeth 80 to 90.

S. officinalis. Back dark green, with six longitudinal but interrupted reddish stripes; belly olive-green and without spots; teeth not exceeding 75, usually 70.

Preservation.—In water kept from putrefaction by the addition of a few fragments of iron. Unglazed

brown pans may be advantageously used tied over with coarse cloth so as not to exclude air. The aquarium principle has been adopted with success. A bed composed of clean pebbles of various sizes mixed with sand and a little moss has been found to answer well. The containing jar should be kept at a moderate temperature, and preferably in an underground situation.

Uses.—For extracting blood, each leech taking from 1 to 2 drachms, capable of being increased to 4 drachms by subsequent fomentation.

DIVISION II.—ARTHROPODA, Or ARTICULATA.

(ἀρθρον, a joint; πούς, a foot.)

CLASS.—INSECTA.

Head, thorax and abdomen distinct. Six legs borne on the thorax, and two pairs of wings on the same portion of the body; antennæ 2 in number; respiration effected by tracheæ.

ORDER.—HEMIPTERA.

(ἥμι half; πτεξὸν a wing; *i.e.*, having half of the elytra membranous.)

SUB-ORDER.—HOMOPTERA.

(ὁμός, similar; πτεξὸν, a wing.)

COCCUS (B.P.), Cochineal. — The dried, matured and fecundated female insect of *Coccus cacti*, reared in Mexico and Teneriffe.

Description. — *Females.* — Body broadly elliptical. bluish red, about 1-6th of an inch long, and usually covered with a silvery white powder which is of a fatty nature melting by heat, when the insect becomes reddish black; *apterous* with a *rostrum* and antennæ short.

Males. — Deep red, only half the size of the female. with two long divergent setæ at the posterior end: *rostrum* absent; antennæ nearly as long as the body.

Uses. — As a colouring matter, and reputed (but without foundation) to be anodyne and anti-spasmodic.

Chemistry.—The colouring matter of cochineal may be isolated by first percolating with ether to remove fat, and then macerating in alcohol. After distilling off the spirit it remains as a rose coloured powder, soluble in water and alkalies, forming deep purple solutions. It is not changed by dilute sulphuric or hydrochloric acids: it is turned yellow by nitric acid, chlorine, iodine and bromine. It may be precipitated as an insoluble *lake* by plumbic acetate, and forms with tin salts a bright crimson solution. Its formula is $C_{14}H_{14}O_8$, and it has been variously called *carmine*, *carminic acid* and *cochinellin*.

Tinctura Cocci.

Cochineal, in powder	2½ ounces
Proof Spirit.....ad.	20 fluid ounces

Macerate 7 days, strain, press, and filter.

(7 grains of Cochineal in 1 drachm.)

Cochineal is an ingredient in Tinctura Cardamomi Composita and Tinctura Cinchonæ Composita.

Commercial Varieties.—Three kinds are imported under the names of Honduras, Mexican or Vera Cruz, and Teneriffe. The two first are distinguished into the

silver and black varieties. An inferior kind is known under the name of *Granilla*.

It has been usually stated that the silver-grey cochineal are most esteemed, but at present it is not really so, the finest being the large, heavy, and perfect blacks, the seconds being the silver-grey, and the lowest the light and broken blacks. It has also been stated that the low quality of blacks are mixtures of unfecundated females and males, but the author's researches conducted on a great number of samples. have proved that all the cochineal of commerce is practically composed of fecundated females. The inferior qualities, however, consist chiefly of females which have been attacked by a species of ichneumon during life, and the interior organs either partially or wholly eaten away.

Adulterations.—Powdered barium sulphate used to be dusted over genuine cochineal previously moistened with gum water. A much heavier sp. gr. was thus obtained. The fraud was concealed by means of ivory black. Lead carbonate has been employed to give a silvery appearance.

ORDER.—HYMENOPTERA.

(ὑμῆν, a membrane; πτερόν, a wing.)

SECTION.—ACULEATA.

The ovipositor is in connection with a gland which secretes a poisonous fluid, and is used as a weapon of defence (*aculeus*, a sting).

MEL (B.P.), Honey.—A saccharine secretion deposited in the honeycomb by the *Apis mellifica*, or hive bee.

Collection.—The bees are killed by sulphurous vapours, and, the honeycombs having been removed, the honey is permitted to drain spontaneously. An inferior quality is obtained by pressure in bags.

Other modes of collection have been introduced with success. In Denmark an ingenious plan is adopted of replacing the honeycomb three separate times by means of moveable frames. A much larger yield of honey is thus secured, and its production has become a national industry.

The mechanical arrangement of the frames is an improvement over the ordinary construction of the hive. Experiment has shown that the bee gathers honey three times in excess of the wax necessary to form the honeycomb.

Uses.—Demulcent, nutritive, and slightly laxative.

Chemistry.—Honey consists essentially of sugar and water, with flavouring matters from the flowers from which it has been collected by the bees. It contains three distinct sugars, viz.:—

1. **Sucrose**, or *cane sugar*, $C_{12}H_{22}O_{11}$, which is crystallisable and immediately blackened by heating with dilute sulphuric acid, but not by potassium hydrate.

2. **Glucose**, *dextrose* or *grape sugar*, $C_6H_{12}O_6 \cdot H_2O$, which is also crystalline, but does not blacken on heating with dilute sulphuric acid, but at once turns dark with potassium hydrate.

3. **Lævulose**, or *fruit sugar*, $C_6H_{12}O_6$, which is uncrystallisable, behaves with re-agents like glucose, but turns the plane of polarisation of light to the left.

Both glucose and lævulose precipitate red cuprous oxide from **Fehling's Solution of Copper**, while sucrose does not.

This solution is made by dissolving 350 grains of cupric sulphate in water, adding 1730 grains of Rochelle salt, and 600 grains of potassium hydrate, and making the whole up to 10,000 grains with water. It is of a clear deep blue, and should not alter when boiled with ten times its bulk of distilled water. 100 grains of this solution will be exactly decolorised by .5 grain of pure grape or fruit sugar.

Mel Boracis.

Borax, in fine powder	64 grains
Clarified Honey	1 ounce

The solution of the borax in 1 drachm of glycerin, added to 6 drachms of Honey, is an improvement. (*Squire.*)

Mel Depuratum.

Honey melted in a water-bath, and strained, while hot, through flannel moistened with warm water.

Oxymel.

Clarified Honey	40 ounces
Acetic Acid	5 fluid ounces
Distilled Water	5 fluid ounces

Sp. gr. 1.320.

Dose—1 to 2 drachms.

Honey is also contained in *Confectio Piperis*, *Confectio Scammonii*, *Confectio Terebinthinæ*, and *Oxymel Scillæ*.

Adulterations.—1. *Starch*, detected by boiling with water, and then testing with iodine solution.

2. *Cane sugar*. Owing to the fact that honey really contains a small quantity of sucrose, this adulteration can only be detected by a quantitative estimation of the cane sugar, by first estimating the glucose and lævulose by means of Fehling's solution, and then boiling another portion with a few drops of dilute sulphuric acid to convert the sucrose into lævulose. and once more estimating with Fehling's solution. Any excess of sugar found in the second estimation over 5 per cent. may be taken as deliberate adulteration. The older the honey the less is the natural amount of sucrose, and in most commercial samples it does not exceed 1 per cent.

3. *Glucose*. This sugar is now much more used than sucrose for adulterating honey, as it is made from starch in great quantities and at very small cost by the action of dilute sulphuric acid and subsequent neutralisation with lime. Its detection is a matter of experience in the analysis of sugars, and would be too lengthy for detail, suffice it to say that the process depends:—(1). On a total estimation of the glucoses (both dextrose and lævulose) by means of Fehling's

solution. (2). The subsequent use of the polarimeter to determine the respective amounts of right and left handed sugar present. Those who have no polarimeter may separate the glucose and lævulose by taking 100 grains of the honey dried at 212° F. and triturating well with 60 grains of calcium hydrate (pure) and 1,000 grain measures of water. The insoluble calcium salt of lævulose thus produced, having been separated by filtration and pressure, is suspended in water, carbonic anhydride is passed to precipitate the calcium, and the whole is then filtered and evaporated on a water-bath and dried at 212° F., or still better *in vacuo*.

CERA FLAVA, Yellow Wax. — The prepared honeycomb of *Apis mellifica*.

CERA ALBA, White Wax.—Yellow wax bleached by exposure to moisture, air and light.

Collection.—Yellow wax is left in the bags after the expression of the honey, and is melted in hot water and then cooled. White wax is prepared by allowing melted wax to run in thin streams upon revolving cylinders kept wet. It thus forms very thin ribbons which are

exposed moist to air and light until bleached. The same action is more rapidly obtained by the action of oxidants, such as dilute nitric acid and potassium permanganate. Chlorine is sometimes employed, but it acts chemically on the wax.

Description.—*Yellow wax* has an odour of honey, is not greasy to the touch, and does not melt under 140° F. It is quite insoluble in cold alcohol, but soluble in turpentine.

White wax is nearly white and translucent, not greasy and does not melt under 150° F.

Chemistry.—When wax is boiled with alcohol it is divided into 2 parts : one totally insoluble (73 per cent.), and the other soluble (22 per cent.) but crystallising out on cooling. The former is *Melissyl palmitate* $C_{30}H_{61}.C_{16}H_{31}O_2$, which can be obtained by dissolving the insoluble portion in benzol and crystallising. It melts at 147.2° F., and is soluble in ether and turpentine. The latter is *Ceryl Cerotate*, $C_{27}H_{55}.C_{27}H_{53}O_2$, and is also soluble in ether. The colour and odour of yellow wax is due to about 5 per cent. of an acid fat called *cerolein*, soluble in cold alcohol.

Yellow wax is contained in Emplastrum Calefaciens, Emplastrum Cantharidis, Emplastrum Cerati Saponis, Emplastrum Galbani, Emplastrum Picis, Unguentum Cantharidis, Unguentum Hydrargyri Compositum, Unguentum Hydrargyri Oxidi Rubri, Unguentum Picis Liquidæ, Unguentum Resinæ, Unguentum Sabinæ, and Unguentum Terebinthinæ.

White wax is contained in Charta Epispastica, Suppositoria Acidi Tannici, Suppositoria Hydrargyri, Suppositoria Morphię, Suppositoria Plumbi Composita, Unguentum Cetacei, Unguentum Plumbi Subacetatis Compositum, and Unguentum Simplex.

Unguentum Simplex.

White Wax	2 ounces
Prepared Lard.....	3 ounces
Almond Oil	3 ounces

Unguentum Simplex is contained in Unguentum Antimonii Tartarati, Unguentum Cadmii Iodidi, Unguentum Creasoti, Unguentum Elemi, Unguentum Hydrargyri Ammoniati, Unguentum Hydrargyri Iodidi Rubri, Unguentum Plumbi Carbonatis, and Unguentum Plumbi Iodidi.

ORDER.—COLEOPTERA.

(κολεός, a sheath or case ; πτερόν, a wing.)

CANTHARIS (B.P.), Spanish Fly.—The dried beetle *Cantharis vesicatoria*, collected chiefly in Hungary.

Description.—About $\frac{1}{2}$ to 5-6ths of an inch long and 1-6th of an inch broad. At once recognised by their long shining and golden-green wing cases (*elytra*) covering two thin membranous and brownish wings.

Collection.—At a certain time in the hot season they suddenly make their appearance, and are in such quantity that their offensive odour causes the inhabitants to desert the public streets in the villages during the period of ten days, which is their time of existence. They are found chiefly on the ash, privet, and lilac, and are shaken off the trees into vinegar, and then dried in the sun or by artificial heat.

Uses.—Externally as a vesicant, and internally as a violent irritant poison. It is reputed to be a powerful aphrodisiac, or stimulant to the sexual organs. The

tincture is given in leucorrhœa, gleet, and incontinence of urine, but ought only to be taken under the advice of an experienced physician.

The best antidotes are oil, Pulvis Cretæ c. Opio, both by the mouth and by injection.

Chemistry.—The active principle is *cantharidin*, $C_5H_6O_2$. It is a solid crystalline body, fusing to a yellow oil at 405° F., and capable of sublimation by heat. It is soluble in ether, acids, and alkalies; and although, *after isolation*, it is insoluble in water and spirit, it can yet be extracted by these menstrua from the flies themselves, owing, doubtless, to its being in some state of combination at present unknown. Its vapour is powerfully irritant, and 1-100th of a grain will distinctly blister the skin.

Acetum Cantharidis.

Cantharides, in powder	2 ounces
Glacial Acetic Acid.....	2 fluid ounces
Acetic Acid	18 fluid ounces

Mix 13 ounces of the acetic acid with the glacial acid and Cantharides; digest at 200° F. for two hours, then percolate; lastly, pass the remaining acetic acid through it, press the contents, and filter.

Product, 1 pint.

(1 part of Cantharides in 10 parts.)

Charta Epispastica. (Blistering Paper.)

White Wax	4 ounces
Spermaceti	1½ ounce
Olive Oil	2 fluid ounces
Resin	$\frac{3}{4}$ ounce
Canada Balsam	$\frac{1}{4}$ ounce
Cantharides, in powder	1 ounce
Distilled Water	6 fluid ounces

Digest all the ingredients, excepting the Canada balsam in a water-bath for 2 hours, stirring constantly, then strain. Mix the balsam with the plaster in a shallow vessel and pass strips of paper over the surface of the hot liquid, so as to receive a thin coating of the plaster.

Emplastrum Calefaciens.

Cantharides, in coarse powder.....	4 ounces
Boiling Water.....	20 fluid ounces
Expressed Oil of Nutmeg	4 ounces
Yellow Wax	4 ounces
Resin	4 ounces
Soap Plaster.....	52 ounces
Resin Plaster	32 ounces

Emplastrum Cantharidis.

Cantharides, in fine powder	12 ounces
Yellow Wax.....	7½ ounces
Prepared Suet	7½ ounces
Resin	3 ounces
Lard	6 ounces

Liquor Epispasticus.

Cantharides, in powder	8 ounces
Acetic Acid	4 fluid ounces
Ether	ad. 20 fluid ounces

(1 part of Cantharides in $2\frac{1}{2}$ parts.)

Macerate the Cantharides in the acid 24 hours, then percolate with the ether.

Tinctura Cantharidis.

Cantharides, in coarse powder.....	$\frac{1}{4}$ ounce
Proof Spirit.....ad.	20 fluid ounces

Macerate 7 days, strain, press, and filter.

DOSE—5 to 20 minims.

Antidotes—Emetics, demulcent drinks, and opiates.

Unguentum Cantharidis.

Cantharides	1 ounce
Yellow Wax.....	1 ounce
Olive Oil	6 fluid ounces

Infuse the Cantharides in the oil, in a covered vessel for 12 hours, then place the vessel in boiling water for 15 minutes, strain through muslin with strong pressure, add the product to the wax previously melted, and stir constantly while it cools.

SUB-KINGDOM V.—MOLLUSCA.

Not of importance to Materia Medica.

SUB-KINGDOM VI.—VERTEBRATA.

The principal portion of the nervous system is dorsal, and is completely shut off from the hæmal system ; all possess a vertebral column. The limbs are never more than four in number.

Vertebrata are divided by Professor Huxley into three primary sections.

DIVISION I.—ICTHYOPSIDA.

(ἰχθύς a fish ; ὄψις appearance.)

CLASS.—PISCES.

Provided with gills throughout the whole of their life. The heart consists of a single auricle and a single ventricle. The limbs are expansions of the integument or fins.

ORDER.—TELEOSTEI.

(τέλειος perfect ; ὀστέον a bone.)

SUB-ORDER.—ANACANTHINI.

(a privative ; ἀκάνθινος spiny.)

OLEUM MORRHUÆ, Oleum Jecoris aselli, *Cod-Liver Oil*. — The oil extracted from the fresh liver of the cod *Gadus Morrhua*, by the application of a heat not exceeding 180° F.

Description.—Pale yellow, with a slight fishy odour and taste and a slightly acid reaction. Sp. gr. .928. A few drops of the oil treated on a slab with a drop of strong sulphuric acid, become first violet and then reddish brown. This test is due to cholic acid, one of the constituents of bile, and is common to all liver oils.

Uses.—Nutritive, being a fat which is more easily digested and assimilated than usual. Given in phthisis and wasting diseases generally.

Chemistry.—The analysis of cod liver oil, according to De Jongh, is, olein, 80·00; palmitin, 15·00; acetin, traces; butyric acid, traces; iodine, ·37; bromine and chlorine, ·15; phosphorus, ·02; cholic acid, traces; salts, 4·46; total, 100·00.

Winkler states that the fats of this oil are not true glycerides, but are salts of *propyl* C_3H_7 , and yield *propyl hydrate* C_3H_7HO , instead of glycerin.

It is certain that by distilling the oil with potassium hydrate and ammonium chloride, we obtain *propylamine* $NH_2C_3H_7$, which is a colourless oily liquid, having a strong odour of herring.

Varieties.—The commercial oils are known as pale, light brown, and dark brown. The pale only is official.

ICHTHYOCOLLA, Isinglass.—The swimming-bladder or sound of various species of sturgeon, *Acipenser*, prepared, cut into shreds, and dried.

Chemistry.—The best Russian isinglass contains 98 per cent. of *gelatin*, which is an albuminous body containing carbon, 50·1; hydrogen, 6·6; nitrogen, 18·3; sulphur, ·14; and the rest oxygen. It is insoluble in

alcohol and ether and when steeped in cold water, swells and dissolves on the application of heat, but sets to a jelly on cooling. It is not affected by dilute acids except tannic acid, which precipitates it from its aqueous solution in the form of finely-divided leather. (It is due to this action that tannic acid turns hides into leather by the process called tanning.) By long boiling with water the solutions of gelatin lose their property of congealing when cold.

Gelatin, Solution of

Isinglass, in shreds	50 grains
Warm Distilled Water	5 fluid ounces

Use.—To distinguish gallic from tannic acid, the latter only giving a yellowish white precipitate.

DIVISION II.—SAUROPSIDA.

(σαῦρα, lizard; ὄψις, appearance.)

CLASS.—AVES.

Circulation double, blood warm. The tarsal and metatarsal bones anchylosed. Covered externally with feathers.

ORDER.—CARINATÆ.

SUB-ORDER.—RASORES, OR SCRATCHERS.

ALBUMEN OVI, White of Egg.—The white of the egg of the common fowl, *Gallus Bankiva* var. *domestica*.

Uses.—Demulcent, and a good antidote for poisoning by any corrosive metallic salt.

Chemistry.—It consists of water, 85; albumen, 12; mucus, 2·7; salts, ·3. The *albumen* of egg differs from that of the serum of blood by being coagulable by ether. Its formula is $C_{72}H_{110}N_{18}SO_{22} \cdot H_2O$. It is soluble in cold water, but coagulable on boiling or on the addition of a few drops of nitric acid, but with excess of this acid it dissolves, forming a pinkish solution. Solution of white of egg may be precipitated by salts of lead, mercury, silver, and copper; the precipitate with the latter being soluble in excess. It is not precipitated by acetic acid (distinction from *casein* of milk), but is thrown down by potassium ferrocyanide in the presence of acetic acid (distinction from gelatin).

Albumen, Solution of

The white of one Egg.

Distilled Water 4 fluid ounces

(This must be recently prepared.)

Use.—To distinguish the glacial from the B.P. phosphoric acids: giving a precipitate with the former, but not with the latter.

VITELLUS OVI, Yolk of Egg.—The yolk contained in a delicate sac called the vitelline membrane of the egg of *Gallus Bankiva* var. *domestica*.

Uses.—Highly nutritive, and of great use in pharmacy for making emulsions.

Chemistry.—A natural emulsion containing 28 per cent. of a yellow oil suspended in water by the aid of 17.5 per cent. of albuminous matter containing phosphorus.

Mistura Spiritus Vini Gallici.

Spirit of French Wine 4 fluid ounces

Cinnamon Water..... 4 fluid ounces

Yolks of Eggs 2

Refined Sugar..... ½ ounce

Dose—1 to 2 ounces.

DIVISION III.—MAMMALIA.

The skull articulates to the vertebral column by two occipital condyles. The blood corpuscles are not nucleated. The young are nourished during a portion of their existence by milk, a secretion of the mammary glands.

ORDER.—CETACEA.

CETACEUM (B.P.), SPERMACETI.—

Nearly pure *cetin* obtained, mixed with oil, from the head of the sperm whale *Physeter macrocephalus*, inhabiting the Pacific and Indian Oceans.

Collection.—After removal from the head it is freed from oil by filtration and pressure, and afterwards purified by melting.

Description.—A white glistening crystalline fat, sp. gr. .940, reducible to powder by rubbing with a little rectified spirit. It melts at 120° F., and is insoluble in cold, but entirely dissolved by boiling alcohol.

Chemistry.—The cetin of spermaceti is not a true glyceride like most other saponifiable fats, but is *Cetyl palmitate* $C_{16}H_{33}.C_{16}H_{31}O_2$, and yields when boiled with alkalies a palmitate of the alkali used, and *Cetyl hydrate*, $C_{16}H_{33}.HO$, commonly called *ethal*, instead of glycerin. Ethal is an alcohol homologous with ordinary alcohol. It is volatile by heat and by oxidation, and yields palmitic acid just as common alcohol yields acetic acid.

Unguentum Cetacei.

Spermaceti	5 ounces
White Wax	2 ounces
Almond Oil	20 fluid ounces

Spermaceti is an ingredient in Charta Epispastica.

ORDER.—UNGULATA.

(*Or* HOOFED QUADRUPEDS.)

This important division of Mammalia includes three entire old orders, Pachydermata, Solidungula, and Ruminantia.

SUB-ORDER.—ARTIODACTYLA.

(ἄρτιος, even; δάκτυλος, a toe.)

The toes even in number, two or four.

ADEPS PRÆPARATUS, Prepared Lard.—The purified internal abdominal fat of the pig *Sus scrofa*, belonging to the old order **Pachydermata** (*παχὺς*, thick; *δέρμα*, the skin.)

Purification.—Remove as much of the membranes as possible from the internal fat of the abdomen of the hog perfectly fresh. Cut the fat into small pieces, put it into a suitable vessel with about four gallons of *cold water*, and, while a current of water is running through the vessel, break up the masses of the fat with the hands, exposing every part to the water, so that whatever is soluble may be thus dissolved and carried away. Afterwards collect the washed fat on a sieve or in a cloth, drain away the water, liquefy the fat at a heat not exceeding 212° , and strain through flannel, pressing the residue while hot; then put it into a pan heated by steam, and keep it a temperature a little, but not much, above 212° , stirring it constantly until it becomes clear and entirely free from water. Finally, strain it through flannel.

Chemistry.—A pure white fat, without odour, melting about 100° F., and entirely soluble in ether. Stated in the books to be a mixture of 62 per cent.

glyceryl oleate (olein), $C_3H_5\beta C_{18}H_{33}O_2$, with palmitin and stearin; but the author, in his researches into lard, has found that these proportions are neither accurate nor constant, and that, in fact, they vary with the age and sex of the animal. Water which has been boiled with lard should not precipitate argentic nitrate (absence of salt), nor turn blue with solution of iodine (absence of starch).

Prepared lard is contained in the following ointments:—*Aconitiæ*, *Atropiæ*, *Belladonnæ*, *Hydrargyri*, *Hydrargyri Nitratis*, *Hydrargyri Subchloridi*, *Iodidi*, *Potassæ Sulphuratæ*, *Potassii Iodidi*, *Sabinæ*, *Simplex*, *Sulphuris Iodidi*, *Terebinthinæ*, and *Veratriæ*.

Adeps Benzoatus. (*Vide* BENZOIN, page 249.)

Prepared Lard	16 ounces
Benzoin, in coarse powder	160 grains

Heat in a water bath for two hours, stir frequently, and strain. (About 10 grains to 1 ounce.)

Benzoated lard is contained in *Suppositoria Acidi Tannici*, *Suppositoria Hydrargyri*, *Suppositoria Morphiæ*, *Suppositoria Plumbi Composita*, *Unguentum Gallæ*, *Unguentum Gallæ c. Opio*, *Unguentum Plumbi Acetatis*, *Unguentum Sulphuris*, and *Unguentum Zinci*.

PEPSIN (B.P.).—A preparation of the mucous lining of the fresh and healthy stomach of the pig, sheep, or calf.

Preparation.—The stomach of the recently killed animal having been cut open, and laid on a board with the inner surface upwards, any adhering portions of food, or other undissolved matter, are to be removed, and the exposed surface slightly washed with cold water; the cleansed mucous membrane is then scraped with a blunt knife or other suitable instrument, and the viscid pulp thus obtained is to be immediately spread over the surface of glass or glazed earthenware, and quickly dried at a temperature not exceeding 100°. The dried residue is to be reduced to powder, and preserved in a stoppered bottle.

Characters and Tests.—A light yellowish brown powder, having a faint but not disagreeable odour, and a slightly saline taste, without any indication of putrescence. Very little soluble in water or spirit. Two grains with 1 ounce of water, to which 5 minims of hydrochloric acid have been added, form a mixture in which 100 grains of hard boiled white of egg in thin

shavings will dissolve on their being digested together for about 4 hours at 98° F.

This is caused by the power which the pepsin has of converting albuminous bodies into *peptone* in presence of a dilute acid.

LAC, Milk.—A natural emulsion of fat (butter) in a very weak solution of sugar by the aid of albuminous matter (casein), and containing a small proportion of mineral matter, chiefly chlorides and phosphates. Average cow's milk may be thus represented:—
Water, 87.5; fat, 3.2; sugar, 5.0; casein, 3.6; ash, .7; total, 100.00.

Owing, however, to the fact that cows yield a greater or less quantity of fat according to the nature of their food, analysts generally have agreed that the lowest limit of any sample of mixed milk should be fat, 2.5; solids not fat, 9.0; total solids, 11.5; ash, .65.

To ascertain these points, about 50 grains of the milk should be evaporated to dryness in a weighed platinum dish by means of a water-bath, and the perfectly dry residue weighed. Ether should then be poured on the residue, boiled, and poured off, and this having been repeated until all the fat is extracted, the ether should

be evaporated to dryness in a small weighed beaker, and the residual fat weighed. By deducting this from the total solids, the difference gives "solids not fat," or, in other words, sugar, casein, and mineral matter. The portion remaining in the platinum dish insoluble in ether should then be ignited at a gentle heat, and the ash weighed. The calculation to obtain the amount of pure milk in the sample is

$$\frac{100 \times \text{solids not fat}}{9} = \text{per-centage of pure milk.}$$

The butter of cow's milk is peculiar in containing besides the ordinary glycerides, such as olein, stearin, &c., a considerable per-centage of the glycerides of butyric, caproic, and caprylic acids. When, therefore, 50 grains of pure butter fat, separated from casein, &c., by melting till it becomes clear, and filtering, are saponified by alcoholic solution of potassium hydrate. the solution diluted, and the fatty acids thrown up by adding excess of hydrochloric acid, only the oleic, stearic, and palmitic acids float to the surface, while the butyric, caproic, and caprylic acids remain in solution. The insoluble acids, when washed with boiling water. dried, and weighed, should not exceed 89 per cent. as the utmost limit. (For further information as to the

taking of the "actual density" and soluble acids, see the author's paper in the "*Analyst*," No. 13, for April, 1877.)

SACCHARUM LACTIS, *Sugar of Milk*, $C_{12}H_{22}O_{11} \cdot H_2O$. A crystallised sugar, obtained from the whey of milk by evaporation. Usually in cylindrical masses, 2 inches in diameter with a cord in the axis. Greyish white and crystalline on the surface and in its texture; gritty when chewed and slightly sweet. Less soluble in water than common sugar; slightly soluble in alcohol, and yields mucic acid by the action of nitric acid. It forms a sort of connecting link between gums and sugars.

Uses.—To dilute stronger medicines, and as a means of giving them in the state of powder.

FEL BOVINUM PURIFICATUM
B.P., *Purified Ox Bile*.—The purified gall of the ox, *Bos Taurus*, belonging to the old order **Ruminantia**.

Uses.—In habitual constipation and in morbid states of the liver involving defective secretion.

Chemistry.—Bile is the fluid secreted by the liver of all mammalia. It is practically a solution of a natural soap formed by glycocholic and taurocholic acids in

combination with sodium. Both these acids are conjugate bodies, the glycocholic having a formula of $C_{26}H_{43}NO_6$, and being built up of cholic acid, $HC_{24}H_{39}O_5.H_2O$, and glycocin, $C_2H_5NO_2$. It is soluble in alcohol and ether, but only slightly in water. The taurocholic acid has the composition $C_{26}H_{45}NO_7S$, and is produced by the union of cholic acid with *taurin* $C_2H_7NO_3S$.

The official extract is yellowish green, soluble in water and spirit, but insoluble in ether. Its watery solution, mixed with a drop of syrup and then sulphuric acid added until the precipitate first formed is redissolved, becomes first cherry-red, then carmine, purple, and finally violet. This is a test for the presence of cholic acid, and works with any solution of bile.

Fel Bovinum Purificatum.

Fresh Ox Bile	20 fluid ounces
Rectified Spirit	40 fluid ounces

Mix and set aside for 12 hours, decant and evaporate to an extract.

DOSE—5 to 10 grains, in pills, coated with an ethereal tincture of tolu.

MOSCHUS (B.P.), Musk.—The inspissated and dried secretions from the preputial follicles of the *Moschus moschiferus*, or Musk Deer, belonging to the

old order **Ruminantia**. Imported from China and Japan.

Description. — In small reddish black grains, unctuous in appearance, and contained in a round or slightly oval membranous sac, about 2 inches in diameter, curved on the outer side with stiff greyish hairs surrounding its central orifice. Each sac weighs about $\frac{3}{4}$ of an ounce, and contains about 160 grains weight of musk.

Varieties.—1. Chinese, which is the official musk.
2. Russian or Siberian, in longer grains, but inferior.

Uses. — Stimulant, anti-spasmodic, and reputed aphrodisaic.

Chemistry.—The odorous principle of musk is volatile, but has not been yet satisfactorily investigated.

Substitute.—Chinese musk, made up of dried blood with a little ammonia and a few true musk grains, sewed up in a piece of the skin of the musk ox.

SEVUM PRÆPARATUM, Prepared Suet. — The internal abdominal fat of the sheep, *Ovis aries*, purified by melting and straining.

Description.—White, smooth, greasy, and fusible at 103° F. Entirely soluble in ether, but crystallising out on the addition of a little rectified spirit.

Chemistry.—Is nearly all glyceryl stearate (stearin), $C_3H_5.3C_{18}H_{35}O_2$, with a little olein and palmitin. Saponified by alcoholic potash, and the solution diluted and acidulated with hydrochloric acid, 95 per cent. of fatty acids separate, which solidify on cooling, and consist chiefly of stearic acid.

Prepared suet is used in making *Sapo animalis*, which is chiefly sodium stearate and is a constituent of Emplastrum Cantharidis and Unguentum Hydrargyri.

ORDER.—RODENTIA.

CASTOREUM, Castor.—The dried preputial follicles of *Castor fiber* (the Beaver), separated from the somewhat shorter and smaller oil sacs which are frequently attached to them. From Hudson's Bay Territory.

Source.—The beaver is peculiar in possessing as an appendage to its prepuce a pair of sacs of a pear-like shape, and about 3 inches long. They are found both in the male and female, and are situated under the skin

in a hollow space between the cloaca and the pubic arch, into which both the organs of generation and the rectum open. They must not be confused with the testicles, having no connection therewith.

Description.—In pairs about 3 inches long, and in appearance something like a large and very dark-coloured fig, but much heavier.

Chemistry.—The secretion contained in the sacs is really orange-yellow during life, but in the dried article it is brown and almost entirely soluble both in alcohol and ether. It contains a volatile oil, an amorphous, dark brown, bitter and acrid resin, and a crystalline fat called *castorin*, soluble in ether. This fatty body resembles cholesterin, and is not susceptible of saponification. Wöhler has found that the volatile oil contains both carbolic and salicylic acids.

Uses. — Reputed anti-spasmodic, and used like asafoetida in hysteria. It has been also stated to possess emmenagogue properties.

Tinctura Castorei.

Castor, in coarse powder	1 ounce
Rectified Spirit	ad. 20 fluid ounces

Macerate and strain.

Dose— $\frac{1}{2}$ to 1 fluid drachm.



APPENDICES.



APPENDIX I.

A Classification and Description of the Chief Starches existing in the Commoner Articles of Materia Medica and Food.

The following Classification, first made public by me in the "ANALYST" for December, 1876, is the one which I have for some years adopted in teaching Microscopic Analysis. The whole of the matter contained in these explanatory Notes depends for its correct application on the use of the 4-10th-inch power and "B" eye-piece.

SAMPLING, MOUNTING, AND OBSERVING.

The best way of obtaining a fair sample of the starch is to mix it thoroughly in a mortar, and then take a clean and dry glass rod and with it stir the whole well up.

Having now got the slide ready, the rod is withdrawn from the starch, and touched sharply on the glass. By this means a sufficient number of granules will detach themselves and fall upon the slide. A drop of distilled water having been added, the covering-glass put on, and superfluous moisture removed with a slip of blotting paper, all is ready. It should be noted that the less the number of granules transferred to the slide the better; indeed, after mounting, the film of starch should just be visible to the eye as a thin milky cloud.

The next point to be considered is the power suitable for use; and here I must say that many of the common errors are induced by the tendency to employ too high an objective at the first observation. The most satisfactory power for general use by commercial or food analysts is a 4-10th-inch objective, and a "B" micrometer eye-piece. This will show any starch with sufficient distinctness, and will not unduly magnify. Anyone not having this power should certainly provide himself with one if he wants to work accurately and comfortably.

A very important matter in the examination of starches is the arrangement of light, which should *never*

be direct, but always somewhat oblique. Great advantages will be apparent by changing the direction of the light now and then during the examination of a field; indeed, I am of opinion that the first thing which distinguishes the true microscopist is the judicious handling of his mirror, by which almost any effect can be attained. When polarising effects are wanted I always mount in water, using a blue selenite plate beneath the object; and the descriptions given hereafter of the behaviour of the starches in polarised light are dependent on these conditions. As a rule, the flat starches polarise better than the convex ones, and angular granules do not show any marked effects at all.

The points to be observed in looking at a starch are—

1. The shape.
2. The position and nature of the *hilum*.
3. The concentric rings.

The two first observations are simple, but the examination for rings requires care. The usual idea produced in the mind of anyone from observation of the ordinary woodcut drawings is that nearly all

starches are covered with distinct rings. In practice with the 4-10th power this delusion is soon dissipated, as, even in the larger starches, the rings can only be caught by altering the focus continually with the fine adjustment, and keeping the eye, while doing so, on certain of the bolder granules. There are only really two starches which can be honestly presented as showing highly-marked rings on nearly every granule with an ordinary microscope, namely, turmeric and Natal arrowroot. It is very rarely that we find more than three or four granules, in a field of potato, for example, so equal in size as to show rings at the same focus. The oblique arrangement of the light is an absolute necessity in examining for rings; and it is here that a slight alteration of light during observation, as already recommended, tells wonderfully. There are several starches (wheat, for example) which show no rings, even in the best light. When the hilum is anywhere near the centre of the granules, the rings are seen all round it; but when it is near one end, only a segment of each ring is seen. For the purpose of brevity, I will call the former *complete*, and the latter *incomplete*, rings.

MOUNTING STARCHES FOR PRESERVATION.

After trying numerous systems, I have finally arrived at the conclusion that starch retains its characters longest by being simply mounted in glycerin one part, and camphor water two parts. After removing superfluous moisture carefully, I surround the covering-glass edge with a good layer of blue mounting varnish by means of a turn-table. The varnish can be obtained from the usual dealers in microscopic apparatus.

MEASUREMENTS OF STARCHES.

The *measurements* of starches are most important, and are performed in the usual manner by first comparing a stage micrometer, graduated in $\cdot 01$ of an inch, with the divisions on a "B" micrometer eye-piece. By a careful study of the relation of the lines on the stage and in the eye-piece the exact value of each division of the latter with the 4-10th power is ascertained, and it is not then necessary to calculate each starch to fractions of an inch, but simply to note the number of divisions of the eye-piece which the image of the granule covers. A very little practice with the stage and eye micrometer will enable the

drawing up of a special table for the eye-piece of one's own microscope, which is a most essential point in calculating the per-centages of mixtures of starches otherwise alike in general appearances. Every food analyst should have such a table of values of his eye-piece divisions always ready for reference.

In measuring I have used the English inch, as a more familiar standard to most persons, than the French millimetre. For example, in measuring with my eye-piece, I find that $\cdot 01$ of an inch equals 27 main divisions. Each main division of the eye-piece, therefore, represents $\cdot 00037$ of an inch. Suppose now we find that the largest granule of arrowroot in a field covers 6 main divisions, we have $\cdot 00037 \times 6 = \cdot 00222$ of an inch for the largest size of arrowroot. In measuring any starches, such as rice and pepper, which run less in size than one division with the 4-10th power, it is necessary to use a higher power, say 1-5th or 1-8th. The divisions will now probably run 50 to the $\cdot 01$, and each division with this power will show only $\cdot 00002$ of an inch, which will enable the measurement to be made more accurately, a great point in distinguishing rice from pepper.

CLASSIFICATION OF THE STARCHES.

There is no possibility of making any classification of starches based upon the natural orders, and therefore I have adopted a purely empirical one, based on the three points of observation already mentioned. It will be found of the greatest use in practice as at once reducing the starch under observation to being one of a certain few, and this narrowing of the ground being gained, then the more minute points have to be considered for final identification. It is to be distinctly understood that the characters are those of the *majority* of the granules in any one field.

The whole of the common starches may be classified into five characteristic groups as follows :—

GROUP I.

The starches of this group are characterised by their more or less oval or ovate forms, and by their always having the *hilum*, and usually the *concentric rings*, clearly *visible*. The group includes tous les mois, potato, arrowroot, calumba, orris root, ginger, galangal, and turmeric.

GROUP II.

Starches having the *concentric rings* all but *invisible*, and a very strongly marked *stellate hilum*. The group includes bean, pea, maize, lentil, dari, and nutmeg.

GROUP III.

Starches having both the *concentric rings* and *hilum* invisible in the majority of the granules. The group includes wheat, barley, rye, jalap, rhubarb, senega, sumbul, bayberry, chestnut, acorn, calabar bean, liquorice root, white hellebore, green hellebore, and black hellebore.

GROUP IV.

All the granules being more or less truncated at one end. The group includes sago, tapioca, belladonna, colchicum, scammony, podophyllum, canella, aconite, arum, cassia, and cinnamon.

GROUP V.

Having all the granules angular in form. The group includes oats, tacca, rice, pepper, and ipecacuanha.

Having decided to which group the starch belongs, the following table will aid in its individual identification.

Table for the Detection of Starches when magnified to about 225 diameters.
All measurements are given in decimals of an inch.

GROUP I.—All more or less oval in shape, and having both *hilum* and *rings* visible.

NAME.	SHAPE.	NORMAL MEASUREMENT.	REMARKS.
Tous les Mois	Oval, with flat ends	·00370 to ·00185	Hilum annular near one end, and incomplete rings.
Potato	Oval	·00270 to ·00148	Hilum annular, rings incomplete, shape and size very variable.
Bermuda Arrowroot	Sack-shaped	·00148 to ·00129	Hilum distinct annular, shape variable, rings faint.
St. Vincent Arrowroot	Oval-oblong	·00148 to ·00129	Hilum semilunar, rings faint, shape not very variable.
Natal Arrowroot	Broadly ovate	·00148 to ·00129	Hilum annular in centre, and well marked complete rings.
Galangal	Skittle-shaped	About ·00135	Hilum elongated, very faint incomplete rings.
Calumba	Broadly pear-shaped	·00185	Hilum semilunar, faint but complete rings, shape variable.
Orris Root	Elongated oblong	·00092	Hilum faint, shape characteristic.
Turmeric	Oval-oblong and conical	·00148	Very strongly marked incomplete rings.
Ginger	Shortly conical, with [rounded angles	·00148	Hilum and rings scarcely visible, shape variable but characteristic.

GROUP II.—With strongly developed *hilum*, more or less stellate.

NAME.	SHAPE.	NORMAL MEASUREMENT.	REMARKS.
Bean	Oval-oblong	About '00135	Fairly uniform.
Pea	Like bean	'00111 to '00074	Very variable in size, with granules under '00111 preponderating.
Lentil	Like bean	About '00111	Hilum a long depression, seldom radiate.
Nutmeg	Rounded	" '00055	The small size and rounded form distinctive.
Dari.....	Elongated hexagon	" '00074	Irregular appearance and great convexity distinctive.
Maize	Round and polygonal ..	" '00074	The rounded angles of the polygonal granules distinctive.

GROUP III.—*Hilum* and *rings* practically invisible.

NAME.	SHAPE.	NORMAL MEASUREMENT.	REMARKS.
Wheat	Circular and flat	'00185 to '00009	<i>Very variable</i> in size, and <i>very dull</i> polarisation in water.
Barley	<i>Slightly</i> angular circles.	'00073, a few 4 times [the size	The majority measuring about '00073 distinct.
Rye	Like barley	'00148 to '00009	Small granules quite round, and here and there cracked.
Jalap	Like wheat	Like wheat	Polarises <i>brightly</i> in water.
Rhubarb	"	'00055 to '00033 for [small	Polarises between jalap and wheat, and runs smaller and more convex.
Senega	"	'00148 to '00009	Measurements the only guide.
Bayberry	"	'00074 to '00011	
Sumbul	"	'00066 to '00009	
Chestnut	Very variable	'00090 to '00009	Variable form, and small but regular size distinctive.
Acorn	Round-oval	About '00074	Small and uniform size distinctive.
Calabar Bean	Oval-oblong	'00296 to '00180	Large size and shape characteristic.
Liquorice	Elongated oval	About '00018	Small size and shape distinctive.
Hellebore, green or [black	Perfectly rotund	'00037 to '00009	Small regular size and rotundity distinctive.
" white ..	Irregular	'00055 to '00009	Irregular shape and faint central depression distinctive.

GROUP IV.—More or less truncated at one end.

NAME.	SHAPE.	NORMAL MEASUREMENT.	REMARKS.
Cassia	Round	'00111 to '00018	Round and muller-shaped granules, and faint circular hilum.
Cinnamon	Like cassia.....	'00074 to '00009	More frequently truncated than cassia, and smaller.
Sago, raw	Oval-ovate	'00260 to '00111	Has circular hilum at convex end, and rings faintly visible.
" prepared	"	"	Has a large oval or circular depression, covering $\frac{1}{3}$ nearly of each granule.
Tapioca	Roundish	'00074 to '00055	A little over 50 per cent. truncated by <i>one</i> facet and a pearly hilum.
Arum	"	About '00056.....	Smaller than tapioca, and truncated by <i>two</i> facets.
Belladonna	"	Not distinguishable from tapioca.
Colchicum	"	About '00074.....	Larger than tapioca, and contains many more truncated granules.
Scammony.....	"	" '00045.....	Smaller than tapioca, more irregular, and hilum not visible.
Canella	Very variable.....	'00033 to '00022	Very variable form and small size the only points.
Podophyllin	Like tapioca	About '00040.....	Like scammony, but has visible hilum in most of the granules.
Aconite	"	" '00037.....	Like tapioca, but half the size.

GROUP V.—All granules more or less polygonal.

NAME.	SHAPE.	NORMAL MEASUREMENT.	REMARKS.
Tacca	Polygonal or hexagonal	'00075 to '00037	Distinguished from maize by its <i>sharp</i> angles.
Oat	Polygonal	About '00037	Larger than rice, and hilum visible in some granules.
Rice.....	"	'00030 to '00020	Measurement using $\frac{1}{8}$ or $\frac{1}{16}$ -inch power, and then hilum visible.
Pepper	"	'00020 to '00002	" " "
Ipecacuanha	"	About '00018	Some round and truncated granules adhering in groups of three.



APPENDIX II.—MUSEUM CABINET.

DRAWERS FOR THE RECOGNITION
OF
MATERIA MEDICA SPECIMENS,
ARRANGED FOR THE USE OF STUDENTS
AT
THE SOUTH-LONDON SCHOOL OF PHARMACY,
KENNINGTON.

STUDENTS HAVE ALSO FREE ACCESS TO THE GENERAL MUSEUMS
OF CHEMISTRY, BOTANY, AND MATERIA MEDICA.

1878.

DRAWER A.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Nux Vomica.	Seeds.	Strychnos Nux vomica.
1 A	St. Ignatius' Beans.	Seeds.	Strychnos Ignatia.
2	Croton Seeds.	Oil.	Croton Tiglium.
3	Cocculus Indicus.	Fruit.	Anamirta Cocculus.
4	Bay Berries.	Fruit.	Laurus nobilis.
5	Coriander.	Fruit.	Coriandrum sativum.
6	Fœnugreek.	Seeds.	Trigonella Fœnum-græc.
7	Colchicum.	Seeds.	Colchicum autumnale.
8	Stramonium.	Seeds.	Datura Stramonium.
9	Nutmegs.	Kernels.	Myristica officinalis.
10	Cubebs.	Fruit.	Cubeba officinalis.
11	White Pepper.	Ripe fruit.	Piper nigrum.
12	Black Pepper.	Unripe fruit.	Piper nigrum.
13	Dill.	Fruit.	Anethum graveolens.
14	Anise.	Fruit.	Pimpinella anisum.
15	Hemlock.	Fruit.	Conium maculatum.
16	Carrot.	Fruit.	Daucus Carota.
17	Fennel.	Fruit.	Fœniculum dulce.
18	Cumin.	Fruit.	Cuminum Cyminum.

DRAWER A.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Loganiaceæ.	East Indies, [Coromandel.	Ext. Nucis vom. 1 from 16. Liq. Strychniæ, 1 in 120. Strychnia. Tinct. Nucis vom. 1 in 10.
Loganiaceæ.	Philippine Islands.	Non-official.
Euphorbiaceæ.	Ceylon.	Lin. Crotonis, 1 in 8.
Coniispermaceæ.	Malabar.	Non-official.
Scrophulariaceæ.	Southern Europe.	Non-official.
Campanulaceæ.	Cultivated in Britain.	Conf. Sennæ. Mist. Gentianæ. Syr. Rhei. Tinct. Rhei. Tinct. Sennæ. Ol. Coriandri (Syr. Sennæ).
Leguminosæ.	Southern Europe,	Non-official.
Colchicaceæ.	Cultivated in Britain.	Tinct. Sem. Colchici, 1 in 8.
Scrophulariaceæ.	Indigenous.	Ext. Stramon. 1 from 8. Tinct. „ 1 in 8.
Ericaceæ.	Banda Islands.	Pulv. Catechu Co. Spirit, 1 volat. oil. in 50.
Myricaceæ.	Java.	Oleum Cubebæ. Tinct. „ 1 in 8.
Myricaceæ.	Telicherry.	Non-official.
Myricaceæ.	East Indies.	Conf. Piper., 1 in 10.
Campanulaceæ.	Southern Europe.	Aqua Anethi, 1 in 10. Oleum Anethi.
Campanulaceæ.	Asia Minor, cultivated in Southern Europe.	Ess. Anisi, 1 in 4. Sp. Rect. Oleum Anisi.
Campanulaceæ.	Indigenous.	Tinct. Conii, 1 in 8.
Campanulaceæ.	Indigenous.	Non-official.
Campanulaceæ.	Malta.	Aqua Fœniculi, 1 in 10.
Campanulaceæ.	Malta and Sicily.	Non-official.

DRAWER B.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Gamboge.	Gum-resin.	(Garcinia Morella var. Pedicellata.
2	Guaiacum.	Resin.	Guaiacum officinale.
3	Camphor.	Concrete Volatile Oil.	Camphora officinarum.
4	Mastich.	Resin.	Pistacia Lentiscus.
5	Sandarach.	Resin.	Callitris Quadri-valvis.
6	Opium.	Inspissated Juice.	Papaver somniferum.
7	Burgundy Pitch.	Resinous exudation.	Abies excelsa.
8	Resin.	Resin.	Var. Spec. Pinus and Abies
9	Thus.	Crude Turpentine,	Pinus Tæda and P. palustris
10	Olibanum.	Gum-resin.	Var. Spec. Boswellia.
11	Benzoin.	Balsamic Resin.	Styrax Benzoin.
12	"	"	"
13	Barbadoes Aloes.	Inspissated Juice.	Aloe vulgaris.
14	Socotrine Aloes.	Inspissated Juice.	Aloe Socotrina.
15	Cape Aloes.	Inspissated Juice.	Aloe spicata.
16	Hepatic Aloes.	Inspissated Juice.	Aloe Socotrina.
17	Natal Aloes.	Inspissated Juice.	
18	Zanzibar Aloes.		

DRAWER B.

RURAL ORDER.	HABITAT.	PREPARATIONS.
Rutiferae.	Siam.	Pil. Cambogiæ Comp. 1 in 6.
Gophyllaceæ.	San Domingo and [Jamaica.	Mist. Guaiaci, 1 in 40. Pil. Hydrarg. Subchlor. Comp. 1 in 2½ Tinct. Guaiaci Ammoniat. 1 in 5.
Graceæ.	China and Japan.	Aqua Camph. ½ oz. to 1 gall. Liniment, 1 in 5 nearly. " Comp. 1 in 8 nearly. Spirit, 1 in 10 nearly. Tinct. Camph. Comp.
Incardiaceæ.	Scio.	No B.P. prep.
Miferae.	Northern Europe.	Non-official. Pounce.
Baveraceæ.	Smyrna, Persia, Con- stantinople, Egypt, and India.	Opium Preparations. <i>Vide</i> final page.
Miferae.	Black Forest.	Emp. Picis, 1 in 2. " Ferri, 1 in 5½.
Miferae.	America.	Emp. Resinæ, 1 in 9½. Ung. " 1 in 3½.
Miferae.	North America.	Emp. Picis.
Mridaceæ.	Africa and Southern Arabia.	Non-official.
Graceæ.	Siam and Sumatra.	Adeps Benzoatus, 10 grains to 1 oz.
"	" "	Acid. Benzoicum. Tinct. Benzoin Co. 1 in 10.
Graceæ.	Barbadoes.	Enema Aloes, 4 grains to 1 ounce. Ext. Aloes Barb. 8 from 10. Pil. Aloes " 1 in 2. " " et Ferri, 1 in 5½. " Cambogiæ Co. 1 in 6. " Colocynth Co. 1 in 3. " " et Hyos. 1 in 4½.
Graceæ.	Socotra. Imported [from Bombay.	Dec. Aloes Co. 4 grains (ext.) to 1 oz. Enema Aloes, 4 grains to 1 ounce. Ext. Aloes Socot. 1 from 2. " Coloc. Co. 1 (ext.) in 2½. Pil. Aloes et Asafœtidæ, 1 in 4. " " et Myrrhæ, 1 in 3. " " Socot. 1 in 3. " Rhei Co. 1 in 6. Tinct. Aloes, 1 in 40. " Benzoin Co. 8 grains to 1 oz. Vin. Aloes, 16½ grains to 1 ounce.
Graceæ.	Cape of Good Hope.	
Graceæ.	Imported from Bombay.	<i>Vide</i> Supra.
Graceæ.	Natal.	Nataloin.

DRAWER C.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Yellow Wax.	Prepared Honeycomb.	Apis mellifica.
2	White Wax.	Bleached.	"
3	Spermaceti.	Concrete Oil.	Physeter macrocephalus.
4	Cochineal.	Dried insect.	Coccus Cacti.
5	Sugar of Milk.		
6	Cantharides.	Beetle.	Cantharis vesicatoria.
7	Asafoetida.	Gum-resin.	Narthex Asafoetida.
8	Kino.	Inspissated Juice.	Pterocarpus Marsupium.
9	Ammoniacum.	Gum-resin.	Dorema Ammoniacum.
10	Pale Catechu.	Extract.	Uncaria Gambier.
11	Black Catechu.	Extract.	Acacia Catechu.
12	Scammonium.	Gum-resin.	Convolvulus Scammonia
13	Scammony Resin.	Resin.	Convolvulus Scammonia
14	Jalap Resin.	Resin.	Exogonium Purga.
15	Bdellium.	Gum-resin.	Balsamodendron Mukul. pubescens.
16	Myrrh.	Gum-resin.	Balsamodendron Myrrha
17	Acacia.	Gum.	Acacia vera.
18	Tragacanth.	Gum.	Astragalus verus.

DRAWER C.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Hymenoptera.	Everywhere.	Ung. Resinæ, &c.
" "	Pacific and Indian Oceans.	Ung. Simplex, &c.
Diptera.	Mexico and Teneriffe.	Ung. Cetacei, &c.
" "	Switzerland.	Tinct. Cocci, 1 in 8, &c.
Hymenoptera.	Hungary and Russia.	Acet. Canthar. 1 in 10.
" "	Affghanistan and Punjaub.	Charta Epispastica.
" "	Malabar.	Emplast. 1 in 3.
" "	Persia and Punjaub.	Emp. Calefaciens, 1 in 25.
" "	Singapore.	Liquor Epispasticus, 1 in 2½.
" "	Pegu.	Tinct. Canthar. 1 in 80.
" "	Syria and Asia Minor.	Ung. " 1 in 8.
" "	Prepared in Britain.	Enema Asafœtidæ, 30 grs. to 4 oz.
" "	Prepared in Britain.	Pil. " Co. 1 in 3½.
" "	Scinde.	" , Aloes et Asafœtidæ, 1 in 4.
" "	Arabia Felix and Abyssinia.	Spt. Ammon. foetid, 1 in 13½.
" "	Eastern Africa.	Tinct. Asafœtidæ, 1 in 8.
" "	Asia Minor.	Pulv. Catechu Co. 1 in 5.
		" Kino Co. 3¼ in 5.
		Tinct. Kino, 1 in 10.
		Emp. Ammon. c. Hyd. 12 in 15.
		" Galbani, 1 in 11.
		Mist. Ammon. 1 in 32.
		Pil. Ipecac. c. Scillâ, 1 in 7.
		" Scillæ Co. 1 in 6.
		Inf. Catechu, 1 in 30.
		Pulv. " Co. 1 in 2½.
		Tinct. " 1 in 8.
		Troch. " 1 gr. in each.
		Non-official.
		Conf. Scam. 1 in 3.
		Pil. Coloc. Co. 1 in 3.
		" , et Hyos. 1 in 4½.
		Pulv. Scam. Co. 1 in 2.
		Ext. Coloc. Co. 1 in 7.
		Mist. Scam. 4 grs. to 2 oz.
		Pil. " Co. 1 in 3.
		Pil. Scam. Co. 1 in 3.
		Non-official.
		Dec. Aloes Comp. 3 grs. to 1 oz.
		Mist. Ferri Comp. 6 grs. to 1 oz.
		Pil. Aloes et Myrrh, 1 in 6.
		Pil. Asafœtid. Comp. 1 in 3½.
		Pil. Rhei Comp. 1 in 8.
		Tinct. Myrrh, 1 in 8.
		Mucil. Acaciæ, 4 and 6, &c.
		Conf. Opii, 1 in 120.
		Mucilag. Tragac. 1 in 80.
		Pulv. Opii, Comp. 1 in 30.
		" Tragac. Comp. 1 in 6.

DRAWER D.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Lobelia.	Flowering Herb.	Lobelia inflata.
2	Matico.	Leaves.	Artanthe elongata.
3	Senna (inferior).	Leaflets.	Cassia elongata.
4	" (very inferior).	Leaflets.	C. angustifolia.
5	" (very good).	Leaflets.	C. elongata.
6	"	Leaflets.	C. lanceolata and obo-
7	Cusso.	Flowering Tops.	Brayera anthelmintica.
8	Indian Hemp.	Female flowering tops.	Cannabis sativa (Indica.)
9	Aconite.	Leaves.	Aconitum napellus.
10	Henbane.	Leaves.	Hyoscyamus niger.
11	Belladonna.	Leaves.	Atropa Belladonna.
12	Digitalis.	Leaves.	Digitalis purpurea.
13	Hemlock.	Leaves.	Conium maculatum.
14	Chamomile.	Flower heads.	Anthemis nobilis.
15	Bearberry.	Leaves.	Arctostaphylos Uva
16	Buchu.	Leaves.	Barosma betulina.
17	"	Leaves.	B. crenulata.
18	"	Leaves.	B. serratifolia.

DRAWER D.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Solanaceæ.	North America.	Tinct. Lobeliæ, 1 in 8. " " Æther, 1 in 8.
Geraceæ.	Peru.	Inf. Maticæ, 1 in 20.
Rumicinosæ.	East Indies.	Non-official.
"	"	"
"	Southern India.	May be used in place of Alex.
"	Alexandria.	Conf. Sennæ, 1 in 11. Inf. " 1 in 10. Mist. " Co. Pulv. Glycyrr. Co., 1 in 5. Syr. Sennæ, 1 in 2. Tinct. " 1 in 8.
Solanaceæ.	N.E. Abyssinia.	Inf. Cusso, 1 in 16.
Cannabinaceæ.	India.	Ext. Cannabis Ind. Tinct. " " 1 Ext. in 20.
Aconitaceæ.	Cultivated in Britain.	Ext. Aconiti, from fresh leaves and flowers.
Solanaceæ.	Indigenous.	Ext. from fresh leaves and young branches. Succus. Tinct. Hyos. 1 in 8.
Solanaceæ.	Indigenous.	Ext. from fresh leaves and young branches. Emplast. Succus. Tinct. 1 in 20. Unguent.
Scrophulariaceæ.	Indigenous.	Infus. Digital. 30 grains to 10 ounces. Tinct. " 1 in 8.
Belliferaæ.	Indigenous.	Cataplasm Conii, 1 in 14. Ext. " fresh leaves and young branches. Succus " " Pil. Conii Comp. 2½ in 3. Vapor Coniæ.
Compositæ.	Indigenous.	Ext. Anthemidis. Inf. " 1 in 20. Oleum "
Solanaceæ.	Indigenous.	Inf. Uvæ Ursi, 1 in 20.
Solanaceæ.	Cape of Good Hope.	} Inf. Buchu, 1 in 20. Tinct. " 1 in 8.
"	"	
"	"	

DRAWER E.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Scammony.	Root.	Convolvulus Scammonia.
2	Rhubarb.	Root.	Rheum officinale.
3	"	Root.	"
4	"	Root.	"
5	"	Root.	Rheum Rhaponticum.
6	Rhatany.	Root.	Krameria triandra.
7	Belladonna.	Root.	Atropa Belladonna.
8	Gentian.	Root.	Gentiana lutea.
9	Dandelion.	Root.	Taraxacum Dens Leonis.
10	Pellitory.	Root.	Anacyclus Pyrethrum.
11	Aconite.	Root.	Aconitum Napellus.
12	Podophyllum.	Rhizome.	Podophyllum peltatum.
13	Pareira.	Root.	Chondodendron tomentosum.
14	Pareira.	Stem.	Cissampelos Pareira.
15	Ipecacuanha.	Root.	Cephaëlis Ipecacuanha.
16	Hemidesmus.	Root.	Hemidesmus Indicus.
17	Jalap (true).	Root.	Exogonium purga.
18	Jalap (false).	Root.	Ipomœa simulans.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Evolvulaceæ.	Syria and Asia Minor.	Resin.
Legonaceæ.	Chinese Tartary and Thibet.	Ext. Rhei.
"	Imported from E.I.	Inf. " 1 in 40.
"	" Batavian."	Pil. " Co. 1 in 4.
"		Pulv. " Co. 1 in 4½.
"		Syrup. Rhei.
"		Tinct. " 1 in 10.
"		Vinum " 1 in 13½.
Legalaceæ.	Oxfordshire.	Non-official.
"	Peru.	Ext. Krameriaæ.
"		Inf. " 1 in 20.
"		Pulv. Catechu Co. 1 in 5.
"		Tinct. Krameriaæ, 1 in 8.
Opopaceæ.	Indigenous.	Atropia.
"		Liquor Unguentum.
"		Atropia Sulph. Liquor.
"		Lin. Bellad. 1 in 1.
Rutianaceæ.	Southern Europe.	Ext. Gent.
"		Inf. " Co. 1 in 80.
"		Mist. " 1 in 40.
"		Tinct. " Co. 1 in 13½.
Staphositaæ.	Indigenous.	Decoct. Tarax. 1 in 20.
"		Ext. " (fresh root).
"		" " Liq. " "
"		Succus " " 3 in 4, &c.
Staphositaæ.	Spain and Levant.	Tinct. Pyrethri, 1 in 5.
Thunbulaceæ.	Southern Europe.	Aconitia.
"		Unguentum Aconitiaæ, 1 gr. to 3j.
"		Tincture, 1 in 8.
"		Liniment, 1 in 1.
Thunbulaceæ.	North America.	Podophylli Resina.
Uispermaceæ.	Brazil.	Decoct. Pareiraæ, 1 in 13½.
"		Ext. " "
"		" " Liq. 1 in 1.
Uispermaceæ.	Brazil.	Non-official.
Uichonaceæ.	Brazil.	Pil. Conii Co. 1 in 6.
"		Pil. Ipecac. c. Scilla, 1 in 23.
"		Pulv. Ipecac. Co. 1 in 10.
"		Troch. Ipecac. ¼ gr. in each.
"		Troch. Morph. c. Ipecac. 1½ gr. in each.
"		Vin. Ipecac. 1 in 20.
Uolepiadaceæ.	India.	Syrup.
Uolvulaceæ.	Vera Cruz.	Ext. Jalapæ.
"		Pulv. " Co. 1 in 3.
"		" Scam. Co. 1 in 3 nearly.
"		Resina Jalapæ.
"		Tinct. " 1 in 8.
Uolvulaceæ.	Tampico.	Non-official.

DRAWER F.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Oak Bark.	Bark of young branches	<i>Quercus pedunculata.</i>
2	Winter's Bark.	Bark.	<i>Drimys Winteri.</i>
3	Bebeeru.	Bark.	<i>Nectandra Rodiæi.</i>
4	Elm.	Bark.	<i>Ulmus campestris.</i>
5	Cusparia.	Bark.	<i>Galipea Cusparia.</i>
6	Canella.	Bark.	<i>Canella alba.</i>
7	Yellow Cinchona.	Bark.	<i>Cinchona Calisaya.</i>
8	Pale Cinchona.	Bark.	<i>Cinchona officinalis, var. Condaminea.</i>
9	Red Cinchona.	Bark.	<i>Cinchona succiruba.</i>
10	Lancifolia.	Bark.	<i>Cinchona lancifolia.</i>
11	Cascarilla.	Bark.	<i>Croton Eluteria.</i>
12	Simaruba.	Root bark.	<i>Simaruba amara.</i>
13	Cassia Pods.	Ripe fruit.	<i>Cassia Fistula.</i>
14	Horse Cassia.	Ripe fruit.	<i>Cassia Brasiliana.</i>
15	Nux Vomica.	Bark.	<i>Strychnos Nux vomica.</i>
16	Wild Cherry.	Bark.	<i>Prunus serotina virginiana.</i>
17	Pomegranate.	Root bark.	<i>Punica Granatum.</i>
18	Pomegranate.	Fruit bark.	<i>Punica Granatum.</i>

DRAWER F.

GENERAL ORDER.	HABITAT.	PREPARATIONS.
Guliferæ.	Indigenous.	Dec. Quercus, 1 in 16.
Euphoniaceæ.	Straits of Magellan, South America.	Non-official.
Euphorbiaceæ.	British Guiana.	Beberia Sulphas.
Euphorbiaceæ.	Indigenous.	Dec. Ulmi, 1 in 8.
Euphorbiaceæ.	South America.	Inf. Cusparia, 1 in 20.
Euphorbiaceæ.	West Indies.	Vin. Rhei, 60 grs. to 1 pint.
Euphorbiaceæ.	Bolivia and Southern Peru.	Dec. Cinch. 1 in 16. Ext. „ Liq. 1 from 4. Inf. „ 1 in 20. Quina Sulph. and various alkaloids. Tinct. Cinch. 1 in 5.
Euphorbiaceæ.	Loxa in Ecuador.	Mist. Ferri Arom. 1 in 16. Tinct. Cinch. Co. 1 in 10.
Euphorbiaceæ.	Western Slopes of Chimborazo.	No B.P. prep.
Euphorbiaceæ.	New Granada.	Quina Sulph.
Euphorbiaceæ.	Bahama Islands.	Inf. Cascarilla, 1 in 10. Tinct. „ 1 in 8.
Euphorbiaceæ.	Jamaica.	Non-official.
Euphorbiaceæ.	East and West Indies.	Conf. Senna, 1 in 8.
Euphorbiaceæ.	Central America.	Non-official.
Euphorbiaceæ.	Coromandel.	Non-official.
Euphorbiaceæ.	North America.	Official in United States Phar.
Euphorbiaceæ.	Southern Europe.	Dec. Granati Rad. 1 in 10.
Euphorbiaceæ.	Southern Europe.	Non-official.

DRAWER G.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Poppy Heads.	Capsules.	<i>Papaver somniferum.</i>
2 & 3	Galls.	Excrescences.	<i>Quercus infectoria</i> formed by punctures and deposits of ova of <i>Diplolepis</i> <i>G.</i> <i>tinctoria</i> .
4	Capsicum.	Fruit.	<i>Capsicum fastigiatum.</i>
5	White Mustard.	Seed.	<i>Sinapsis alba.</i>
6	Black Mustard.	Seed.	„ <i>nigra.</i>
7	Grains of Paradise.	Seed.	<i>Amomum Melegueta.</i>
8	Linseed.	Seed.	<i>Linum usitatissimum.</i>
9	Star Anise.	Fruit.	<i>Illicium anisatum.</i>
10	Pimento.	Berries.	<i>Eugenia pimenta.</i>
11	Ergot.	Sclerotium.	<i>Claviceps purpurea.</i>
12	Calabar Bean.	Seed.	<i>Physostigma venenosum.</i>
13	Stavesacre.	Seed.	<i>Delphinium staphisagria</i>
14	Cevadilla.	Fruit.	<i>Asagraea officinalis.</i>
15	Areca Nat.	Seed.	<i>Areca Catechu.</i>
16	Cardamoms.	Fruit.	<i>Elettaria Cardamomum.</i>
17	Colocynth.	Fruit freed from seeds.	<i>Citrullus Colocynthis.</i>
18	Bael.	Fruit.	<i>Ægle Marmelos.</i>

DRAWER G.

JRAL ORDER.	HABITAT.	PREPARATIONS.
averaceæ.	Cultivated in Britain.	Decoct. Papav. 1 in 10. Ext. " " Syr. " 1 in 3 nearly.
uliferæ.	Mosul and Aleppo.	Tinct. Gallæ, 1 in 8. Ung. " 80 grs. to 1 oz. " " c. Opio. Acid Gallici. " Tannici.
anaceæ.	Zanzibar.	Tinct. Capsici, 1 in 26.
ciferæ.	Indigenous.	{ Cataplasma Sinapis. Charta Sinapis. Liniment Sinapis Comp. Ol. Sinapis.
ciferæ.	Indigenous.	
giberaceæ.	West Coast of Africa.	Non-official.
aceæ.	Indigenous.	Cataplasma Lini. Farina Lini. Inf. Lini. 1 in 30. Ol. Lini.
gnoliaceæ.	China and Japan.	Ol. Anisi.
rtaceæ.	Jamaica.	Aq. Pimentæ, 1 in 11½.
ngi.	Gallicia (chiefly).	Ext. Ergotæ Liq. 1 in 1. Inf. " 1 in 40. Tinct. " 1 in 4.
guminosæ.	Western Africa.	Ext. Physostigmatis.
nunculaceæ.	Southern Europe.	Non-official.
lanthaceæ.	Mexican Andes.	Veratria. Ung. Veratriæ, 8 grs. to 1 oz.
maceæ.	Southern India.	No B.P. prep.
giberaceæ.	Malabar.	Pulv. Cinnam. Co. 1 in 3. Tinct. Cardam. Co. 1 in 80, &c.
curbitaceæ.	Smyrna, Trieste, France and Spain.	Ext. Coloc. Co. 1 in 4½. Pil. " " 1 in 6. " " " et Hyos. 1 in 9.
rantiaceæ.	Malabar and Coro- mandel.	Ext. Bælæ Liq. 1 in 1.

DRAWER H.

NO.	NAME.	PART USED.	WHENCE OBTAINED
1	Cinnamon.	Inner bark.	<i>Cinnamomum zeylanic</i>
2	Cassia. .	Inner bark.	<i>Cinnamomum Cassia.</i>
3A	Orange Peel.	Rind of fruit.	<i>Citrus vulgaris</i> , v. <i>Bigar</i>
3B	Lemon.	Rind of fruit.	<i>Citrus limonum.</i>
4	Liquorice.	Root.	<i>Glycyrrhiza glabra.</i>
5	Irish Moss.	Entire Alga.	<i>Chondrus Crispus.</i>
6	Iceland Moss.	Lichen.	<i>Cetraria islandica.</i>
7	Senega.	Root.	<i>Polygala Senega.</i>
8	Elder Flowers.	Flowers.	<i>Sambucus nigra.</i>
9	Serpentary.	Rhizome.	<i>Aristolochia Serpentari</i>
10	Red Rose.	Unexpanded petals.	<i>Rosa Gallica.</i>
11	Hops.	Strobiles.	<i>Humulus Lupulus.</i>
12	Worm Seed.	Florets.	<i>Artemisia Cina.</i>
13	Quassia.	Wood in chips and raspings.	<i>Picræna excelsa.</i>
14	Sassafras.	Root in chips.	<i>Sassafras officinale.</i>
15	Chiretta.	Entire plant.	<i>Ophelia Chirata.</i>
16	Dulcamara.	Young branches.	<i>Solanum Dulcamara.</i>
17	Savin.	Dried tops.	<i>Juniperus Sabina.</i>
18	Mezereon.	Dried bark.	<i>Daphne Mezereum</i> , <i>Laureola.</i>

DRAWER H.

FAMILIAL ORDER.	HABITAT.	PREPARATIONS.
Rutaceæ.	Ceylon.	Aqua Cinnam. 1 in 8. Ol. " Pulv. C. Comp. 1 in 3. Tinct. Cinnam. 1 in 8, &c.
Rutaceæ.	China and Java.	Non-official.
Gentianaceæ.	Southern Europe.	Inf. Aurant. 1 in 20. " " Comp. 1 in 40. " Gentian Com. Mist. Gentian. Syr. Aurant. 1 Tinct. in 8. Tinct. " 1 in 10. " Recent. 3 in 10. " Gentian Comp. Vinum Aurant.
"	" "	Ol. Limonis. Succus " Syrup " Tinct. " 1 in 8, &c.
Leguminosæ.	Cultivated in Britain.	Ext. Glycyrrh. " " Liq. Pulv. Glycyrrh. Comp. 1 in 5, &c.
Fagaceæ.	West Coast of Ireland.	Non-official.
Urticaceæ.	Northern Europe.	Dec. Cetrariæ, 1 in 20.
Euphyllaceæ.	North America.	Inf. 1 in 20. Tinct. Senegæ, 1 in 8.
Sambucifoliaceæ.	Indigenous.	Aqua Sambuci, 1 in 1.
Asarachaceæ.	North America.	Inf. Serpentariæ, 1 in 40. Tinct. " 1 in 8.
Rosaceæ.	Cultivated in Britain.	Conf. Rosæ Gal. 1 in 4. Inf. " 1 in 40. Syr. " 1 in 17½.
Papaveraceæ.	England.	Ext. Lupuli. Inf. " 1 in 20. Tinct. " 1 in 8.
Compositæ.	Russia.	Santonin.
Ranunculaceæ.	Jamaica.	Ext. Quassiæ. Inf. " 1 in 80. Tinct. " 1 in 27.
Rutaceæ.	North America.	Dec. Sarsæ Co.
Gentianaceæ.	Northern India.	Inf. Chirata, 1 in 40. Tinct. " 1 in 8.
Ranunculaceæ.	Indigenous.	Inf. Dulcamaræ, 1 in 10.
Rubiaceæ.	Cultivated in Britain.	Tinct. Sabinæ, 1 in 8.
Rubiaceæ.	Cultivated in Britain.	Ung. Sabinæ (fresh tops) 1 in 2½.
Simulaceæ.	" "	Dec. Sarzæ Co. Ext. Mezerei Æth.

DRAWER I.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Balsam of Peru.	Balsam.	Myroxylon Pereira
2	Balsam of Tolu.	Balsam.	Myroxylon Toluife
3	Storax.	Balsam.	Liquidambar orient
4	Elemi.	Resin.	Canarium commun
5	Kamala.	Glands of Capsules.	Rottlera tinctoria.
6	Elaterium.	Sediment of juice of fruit.	Ecballium officinar
7	Indigo.	Blue pigment.	Indigofera tinctoria
8	Litmus.	Blue pigment.	Roccella tinctoria.
9	Dragon's Blood.	Resin.	Calamis Draco.
10	Red Sandal Wood.	Wood.	Pterocarpus santal
11	Wheat Starch.	Starch.	Triticum vulgare.
12	Manna.	Exudation.	Fraxinus' Ornus. F. rotundifolia.
13	Castor.	Preputial follicles.	Castor fiber.
14	Spigelia.	Entire herb.	Spigelia Marilandic
15	Saffron.	Stigma and part of style.	Crocus sativus.
16	Cloves.	Unexpanded flowers.	Caryophyllus arom ticus.
17	Bitter Almonds.	Seeds.	Amygdalus commun v. amara.
18	Sweet Almonds.	Seeds.	Amygdalus commun v. dulcis.

DRAWER I.

FAMILIAL ORDER.	HABITAT.	PREPARATIONS.
Euminosæ.	San Salvador.	No B.P. prep.
Euminosæ.	New Granada.	Syr. Tolu. Tinct. „ 1 in 8. Tinct. Benzoin Co.
Angiaceæ.	Smyrna.	Tinct. Benzoin Co.
Abridaceæ.	Manilla.	Ung. Elemi 1 and 4.
Ehorbiaceæ.	India.	No. B.P. Prep.
Curbitaceæ.	Cultivated in Britain.	Pulv. Elaterii Co. 1 in 10.
Euminosæ.	India, chiefly Bengal.	Indigo Sulphate Solution.
Eenes.	Imported from Holland.	Tinct. Litmus 1 in 10.
Enaceæ.	Singapore, Batavia.	Non-official.
Euminosæ.	Ceylon.	Tinct. Lavand Co.
Eninaceæ.	Indigenous.	Glycerinum Amyli. Mucilago „ 12 grs. to fl. oz.
Eceæ.	Sicily.	No B.P. prep.
Eentia.	Hudson's Bay.	Tinct. Castori 1 in 20.
Eaniaceæ.	North America.	Non-official.
Euceæ.	Spain and France.	Tinct. Croci, 1 in 20.
Enaceæ.	Penang, Bencoolen, and Amboyna.	Inf. Caryophyl. 1 in 40.
Euceæ.	Southern Europe, Mogadore.	Ol. Amygdal. Ol. Amygd. Essent.
Euceæ.	Malaga.	Mist. Amygdal. Ol. „ Pulv. „ 8 in 13.

DRAWER J.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Galangal.	Root.	<i>Alpinia officinarum.</i>
2	Ginger.	Scraped rhizome.	<i>Zingiber officinale.</i>
3	Orris.	Rhizome.	<i>Iris florentina.</i>
4	Birthwort.	Root.	<i>Aristolochia longa.</i>
5	Long Turmeric.	Rhizome.	<i>Cucuma longa.</i>
6	Round Turmeric.	Rhizome.	„ <i>rotunda.</i>
7	Alkanet.	Root.	<i>Anchusa tinctoria.</i>
8	Logwood.	Sliced duramen.	<i>Hæmatoxylon campechianum.</i>
9	Squill.	Sliced bulb.	<i>Urginea Scilla.</i>
10	Colchicum.	Sliced corm.	<i>Colchicum autumnale.</i>
11	Calumba.	Root.	<i>Jateorhiza palmata.</i>
12	Sumbul.	Root.	<i>Euryangium sumbul.</i>
12	Male Fern.	Rhizome.	<i>Aspidium Filix Mas.</i>
14	White Hellebore.	Rhizome.	<i>Veratrum album.</i>
15	Green Hellebore.	Rhizome.	<i>V. viride.</i>
16	Black Hellebore.	Rhizome.	<i>Helleborus niger.</i>
17	Arnica.	Rhizome.	<i>Arnica montana.</i>
18	Valerian.	Root.	<i>Valeriana officinalis.</i>

DRAWER J.

FAMILIAL ORDER.	HABITAT.	PREPARATIONS.
Gygiferaceæ.	China and America.	Non-official.
Gygiferaceæ.	West Indies, Africa, &c.	Syrup. Tinct. Zingib. Fort. 1 in 2. " " 1 in 8. Pulv. Cinnam. Co. 1 in 3.
Racææ.	Leghorn and Trieste.	Non-official.
Stolochiaceæ.	Indigenous.	Non-official.
Gygiferaceæ.	Bengal and China.	Paper. Tinct.
Gygiferaceæ.	"	Paper. Tinct.
Diaginacææ.	Southern Europe, Asia Minor.	Non-official.
Huminosæ.	Campeachy, Honduras and Jamaica.	Dec. Hæmatoxyli, 1 in 20. Ext. "
Racææ.	Mediterranean Coast.	Acetum Scillæ, 1 in 8. Oxymel " Pil. Ipec. c. Scillâ. Pil. Scillæ Co. 1 in 5. Syrup. Tinct. Scillæ, 1 in 8.
Lanthaceæ.	Indigenous.	Ext. Colch. (fresh corm). " " Acet. " Vin. " 1 in 5 (dried).
Mispermaceæ.	Eastern Africa.	Ext. Calumbæ, 1 from 8. Inf. " 1 in 20. Mist Ferri Aromat. Tinct. Calumbæ, 1 in 8.
Belliferæ.	Russia and India.	Tinct. Sumbul, 1 in 8.
Ices.	Indigenous.	Ext. Filicis Liq.
Lanthaceæ.	Alps, Pyrenees, &c.	Non-official.
Lanthaceæ.	Canada and United States.	Tinct. Verat. Vir. 1 in 5.
Junculaceæ.	Southern and Eastern Europe.	Non-official.
Compositæ.	Middle and Southern Europe.	Tinct. Arnicæ, 1 in 20.
Merianaceæ.	Indigenous.	Inf. Valer. 1 in 40. Tinct. " 1 in 8. " " Ammon. 1 in 8.

DRAWER K.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Broom.	Tops.	Sarothamnus Scopariae.
2	Castor Oil.	Seeds.	Ricinus communis.
3	Caraway.	Fruits.	Carum Carui.
4	Copaiba.	Oleo-resin.	Copaifera multijuga.
5	Mulberry.	Ripe Fruit.	Morus nigra.
6	Tobacco.	Leaves.	Nicotiana Tabacum.
7	Raisins.	Dried Fruit.	Vitis vinifera.
8	Cacao Butter.	Concrete oil.	Theobroma Cacao.
9	Prepared Lard.	Abdominal fat.	Sus scrofa.
10	Honey.	Saccharine secretion.	Apis mellifica.
11	Musk.	Dried secretion.	Moschus moschiferus.
12	Carded Cotton.	Hairs of seeds.	V. S. of Gossypium.
13	Treacle.	Residue of refining sugar.	Saccharum officinarum.
14	Tamarinds.	Pulp of fruit.	Tamarindus Indica.
15	Canada Balsam.	Oleo-resin.	Abies balsamea.
16	Pearl Barley.	Husked seeds.	Hordeum distichon.
17	Black Snake Root.	Rhizome.	Actæa racemosa.
18	Indian Aconite.	Root.	Aconitum ferox.

DRAWER K.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Euphorbiaceæ.	Indigenous.	Dec. Scoparii (dried), 1 in 20. Succus „ (fresh), 3 in 4.
Euphorbiaceæ.	Calcutta.	Ol. Ricini. Collodium flexile.
Euphorbiaceæ.	Southern Europe.	Aqua Carui, 1 in 10. Ol. Carui.
Euphorbiaceæ.	Valley of Amazon.	Ol. Copaibæ.
Euphorbiaceæ.	Cultivated in Britain.	Syrup Mori (juice), 1 in 2.
Euphorbiaceæ.	Virginia.	Enema Tabaci, 20 grs. to 8 ozs.
Euphorbiaceæ.	Southern Europe.	Tinct. Card. Comp. Tinct. Sennæ.
Euphorbiaceæ, Byttneriaceæ	Mexico and West Indies.	Various suppositories.
Euphorbiaceæ, Mammalia.		Adeps Benzoat. Ung. Simplex, 3 in 8.
Euphorbiaceæ, Insecta.		Mel. Boracis. Mel. Depurat. Oxymel.
Euphorbiaceæ, Mammalia.	India and China.	No B.P. prep.
Euphorbiaceæ.	North America.	Pyroxylin.
Euphorbiaceæ.	West Indies.	Various pills.
Euphorbiaceæ.	West and East Indies.	Conf. Sennæ.
Euphorbiaceæ.	Canada.	Charta Epispastica. Collodium flexile.
Euphorbiaceæ.	Cultivated in Britain.	Dec. Hordei.
Euphorbiaceæ.	North America.	Official in United States' Pharma- copœia.
Euphorbiaceæ.	India (Nepal).	Non-official. Aconitia.

DRAWER L.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Yellow Root.	Rhizome.	Hydrastis Canadensis.
2	Gulancha.	Stems and roots.	Tinospora cordifolia.
3	Maw Seed.	Seed.	Papaver somniferum.
4	Blood Root.	Rhizome.	Sanguinaria Canadensis.
5	Kokum Butter.	Concrete Oil.	Garcinia Indica.
6	Gurjun Balsam, Wood Oil.	Oleo-resin.	Dipterocarpus turbinatus
7	Marshmallow.	Root.	Althæa officinalis.
8	Cocoa Nibs.	Seed.	Theobroma Cacao.
9	Guarana.	Seed.	Paullinia sorbilis.
10	Guaiacum.	Wood.	Guaiacum officinale.
11	Jaborandi.	Leaves and young branches.	Pilocarpus pennatifolius.
12	Orange Flowers.	Flowers.	Citrus Bigaradia and Aurantium.
13	Buckthorn.	Dried Fruit.	Rhamnus cathartica.
14	Alder.	Bark.	Rhamnus frangula.
15	Arnica.	Flowers.	Arnica montana.
16	Cherry Laurel.	Leaves.	Prunus Lauro-cerasus.
17	Cabbage Rose.	Fresh Petals.	Rosa centifolia.
18	Quince.	Seeds.	Pyrus Cydonia.

DRAWER L.

GENERAL ORDER.	HABITAT.	PREPARATIONS.
Ranunculaceæ.	America.	Non-official, B.P. Official in United States' Pharmacopœia.
Aspermatophyceæ.	Tropical India.	Non-official in B.P. Official in Indian Pharmacopœia.
Rubiacæ.	Cultivated in Britain.	Non-official.
Rubiacæ.	Canada.	Official in United States' Pharmacopœia.
Leguminosæ.	India.	Official in Indian Pharmacopœia.
Ericaceæ.	Singapore and Southern India.	Non-official.
Rubiacæ.	South of France, Spain. Cultivated in Britain.	Syrup Althææ, P.L.
Ericaceæ, Rutaceæ.	Mexico and West Indies.	Prepared Cocoa.
Rubiacæ.	Brazil.	Non-official.
Myrsinaceæ.	St. Domingo and Jamaica.	Decoct. Sarzæ Comp.
Rubiacæ.	Brazil.	Non-official.
Rutaceæ.	South of Europe.	Aq. Flor. Aurant. Ol. Syrup.
Rubiacæ.	Indigenous.	Succus. Syrup.
Rubiacæ.	Germany.	Non-official B.P. Official in German Pharmacopœia.
Compositæ.	Southern Europe.	Tinct. German Pharmacopœia. 3j to ʒj. Sp. Rectif.
Rubiacæ.	Cultivated in Britain.	Aqua.
Rubiacæ.	Cultivated in Britain.	Aq. Rosæ. Mist. Ferri Comp. Troch. Bismuth.
Rubiacæ.	Western Asia. Cultivated in Southern Europe.	Decoct. Cydon. Non-official.

DRAWER M.

NO.	NAME.	PART USED.	WHENCE OBTAINED
1	White Sandal.	Wood.	<i>Santalum album.</i>
2	Galbanum.	Gum-resin.	<i>Ferula Galbaniflua.</i>
3	Spearmint.	Flowering Plant.	<i>Mentha viridis.</i>
4	Elecampane.	Dried Root.	<i>Inula Helenium.</i>
5	Lactucarium.	Inspissated Juice.	<i>Lactuca virosa.</i>
6	Stramonium.	Leaves.	<i>Datura Stramonium.</i>
7	Stramonium.	Seeds.	<i>Datura Stramonium.</i>
8	Pennyroyal.	Flowering plant.	<i>Mentha Pulegium.</i>
9	Long Pepper.	Dried Spadices.	<i>Piper officinarum.</i> <i>P. Longum.</i>
10	Willow.	Bark.	<i>Salix fragilis, var. speciosa.</i>
11	Larch.	Bark.	<i>Larix Europæa.</i>
12	Venice Turpentine.	Oleo-resin.	<i>Larix Europæa.</i>
13	Juniper.	Fruit.	<i>Juniperus communis.</i>
14	Sarsaparilla.	Root.	<i>Smilax officinalis.</i>
15	Vanilla.	Pods.	<i>Vanilla planifolia.</i>
16	Tous les Mois.	Starch.	<i>Canna edulis.</i>
17	Sweet Flag.	Rhizome.	<i>Acorus Calamus.</i>
18	Club Moss.	Spores.	<i>Lycopodium clavatum.</i>

DRAWER M.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
Malaceæ.	East Indies.	Non-official.
Umbelliferæ.	India and the Levant.	Emp. Galban.
Rubiataæ.	Indigenous.	Aq.
Compositæ.	Southern Europe. Cultivated in England.	Non-official.
Compositæ.	Indigenous.	Non-official. Ext. Lactuæ B.P.
Papaceæ.	Cultivated in Britain.	No B.P. prep.
Papaceæ.	Cultivated in Britain.	Ext. Tinct. 1 in 8.
Rubiataæ.	Cultivated in Britain.	Non-official.
Geraceæ.	India, Singapore, Batavia.	Non-official.
Salicaceæ.	Indigenous.	Salicin. Acid Salicylic.
Rubiaceæ.	Central Europe. Cultivated in Britain.	Tinct. 1 in 8.
Rubiaceæ.	Central Europe. Cultivated in Britain.	Non-official.
Rubiaceæ.	Northern Europe. Cultivated in Britain.	Ol. Spirit.
Euphorbiaceæ.	Central America. Imported from Jamaica.	Decoct. 1 in 8. " Comp. 1 in 8. Ext. liquid.
Umbellidaceæ.	Eastern Mexico.	Non-official.
Rubiaceæ.	Barbadoes.	Non-official.
Geraceæ.	Indigenous.	Non-official.
Rubiopodiaceæ.	Indigenous, and imported from Central and North- ern Europe.	Non-official.

DRAWER N.

NO.	NAME.	PART USED.	WHENCE OBTAINED.
1	Ceylon Moss.	Entire Alga.	Gracilaria Lichenoides
2	Isinglass.	Swimming Bladder.	Acipenser,
3	Pepsin.	Mucous lining of stomach.	Various.
4	Ox Gall.	Purified Bile.	Bos Taurus.

DRAWER N.

NATURAL ORDER.	HABITAT.	PREPARATIONS.
e.	Ceylon.	Non-official B.P. Official Indian Pharm.
ces. (Class.)	Russia.	Gelatin, 10 grains to 1 fluid ounce.
mmalia.		Pulv.
ninantia.		Fel. Bovin purif.

OPIUM PREPARATIONS.

Confect. Opii, ʒ in 40.

Emplast. Opii, 1 in 10.

Enema ,, ½ drm. Tinct. to
2 oz.

Ext. Opii, 1 from 2.

„ „ Liq. 1 Ext. in 20.

Liniment, 1 Tinct. in 2.

Morphia.

Morphiæ Acetas.

Liquor, 1 in 20.

Inject. Hypoderm. 1 gr. in
12 mins.

Morphiæ Hydrochloras.

„ Liqueur, 1 in 120.

Suppositoria, $\frac{1}{2}$ gr. in each.

id. c. Saponē, $\frac{1}{2}$ gr. in each.

Trochisci, $\frac{1}{8}$ gr. in each.

et Ipecac. $\frac{1}{8}$ gr. in each.

Pil. Ipecac. c. Scillâ, 1 in 23½.

Pil. Plumbi. c. Opio, 1 in 8.

., Saponis Comp. 1 in 54.

Pulv. Cretæ Arom. c. Opio, 1 in 40.

„ Ipecac. Comp. 1 in 10.

„ Kino Comp. 1 in 20.

„ Opii Comp. 1 in 10.

Suppos. Plumbi. Comp. 1 grain in
each.

Tinct. Camph. Comp. 2 grs. to 1 oz.

„ Opīi, 1 in 13 $\frac{1}{3}$.

„ „ Amm. 5 grs. to 1 oz.

Troch. Opii, $\frac{1}{10}$ gr. Ext. in each.

Ung. Gallæ c. Opio, 32 grs. to 1 oz.

Vin. Opii, ʒ Ext. in 20.

Apomorphia, Codeia.

Meconia, Meconic Acid.

Narceia, Narcotia.

Papaverine, Thebaia.

Non-
official.

These are the chief preparations, but there are various others.



INDEX.

A	PAGE.	PAGE.	
Abies balsamea.....	375	Acid carminic439	
excelsa	376	catechuic	147
Abietic acid	373	catechu-tannic	147
Abuta amara.....	27	cathartic	134
rufescens	27	cathartogenic	134
Acacia catechu	146, 198	cetraric	422
gum	143	cholic	466
"sorts"	144	chrysamic	398
suma	146	chrysophanic	134, 312
verek	144	cinnamic	127, 128, 328
Acaciæ mucilago	145	cincho-tannic	206, 207
Acetate of morphia	46	citric	96
Acetum cantharidis	449	copaivic	141
ipecacuanhæ	221	crotonic	345
scillæ	404	equisetic.....	4
Achæmium, <i>vide</i> Rosaceæ	147	ergotic	427
Acid abietic	373	erythric	424
aconitic	4	eugenic	170
Arabic	145	evernic	424
artanthic	320	ferulaic	188
benzoic	248, 250	filicic	419
calumbic	23	fumaric	422
camphoretic	324	gallic	359
camphoric	324	gentianic	259
		glycocholic	465, 466

	PAGE.		PAGE.
Acid guaiaconic	82	Acid sumbulolic	186
guaiaretic	82	tannic	356, 357
hydrocyanic tests....	151, 152	taurocholic	465, 466
hyoscinic	285	tiglinic	345
ipecacuanhic	220	tropic	280
ipomœic.....	266	valerianic	186, 224
kinic	206	Acidi gallici glycerinum	360
kino-tannic	125	tannici glycerinum	358
linoleic	80	,, suppositoria.....	76, 358, 447
meconic.....	40, 51	,, c. saponem	358
myristic	325	,, trochisci	358
oleic	80	Acidum benzoicum	250
ophelic	261	sulphuricum aromaticum.....	329
opianic	50	Acipenser	454
orsellic	424	Aconella	7
oxylinoic.....	80	Aconite Indian	9
palmitic	459	Aconiti extractum	7
pelargonic	85	folia	2
phthalic	250	linimentum	8
pinic	373	radix	2, 3
piperic	317	,, Indica	9
polygalic	65	tinctura	8
pyro-gallic.....	357, 360	Aconitia	4, 10
querci-tannic.....	353	aconitate of	6
quinic	206	crystalline	7
quinine sulphate	210	manufacture of.....	5
quinovic.....	206	Aconitiæ unguentum	8, 461
rhatania-tannic	66, 67	Aconitic acid.....	4
rheo-tannic	312	Aconitina	4
ricinoleic	348	Aconitum Cammarum.....	8
rutic	85	ferox	8, 9
salicylic	169, 363	heterophyllum	8
salicylous	363	Napellus	2, 8, 9
santalac	126	neomontanum	8
silvic	373	paniculatum	8
sulphindigotic	121	Acoraceæ	414
sumbulamic	186	Acorin	414
sumbulic	186	Acorus calamus	414

	PAGE.
Acotyledones	418—430
Acrinyl sulphocyanate	61, 62
Acrogenæ	418—420
Actæa racemosa	16
spicata	14, 17
Actææ racemosæ radix	16
Aculeata	441
Aculeus	441
Adeps benzoatus	249, 461
præparatus	460
Ægle marmelos	97
Æsculus hippocastanum	76
Æthusa cynapium	178
African turmeric	392
Albumen ovi	456
solution of	457
Aldehyd benzoic	128, 150
euodic—lauric	85
Alder bark	103
Aleppo galls	355
Alexandrian mastich	112
senna	133
Algæ	429
Alkaloids, cinchona, analysis,	
216—218	
tests	202, 203
Alkanet root	297
Allspice	171
Allyl	188
sulphocyanate	60, 62
Almond oil	152
Almonds, sweet	148
bitter	149
bitter, essential oil	128,
150, 151, 328	
Aloe Barbadosis	396, 397
Capensis	402
Socotrina	399

	PAGE.
Aloe spicata	402
vulgaris	396
Aloes Barbadosis extractum ..	398
pilula	157, 179, 399
Barbadoes	396, 397
Cape	402
decoctum compositum	108,
120, 394, 400	
enema	398
et asafoetidæ pilula ..	189, 400
et ferri pilula	399
et myrrhæ pilula	108, 157,
394, 401	
hepatic	400
Natal	402
pills	157
Socotrine	398, 399
,, extractum	400
pilula	157, 401
tinctura	120, 401
vinum	389, 391, 401
Aloin	397
Alpinia officinarum	391
Althæa officinalis	72
Althææ radix	72, 78
syrupus	73
Altingiaceæ	360—362
American ginseng	66
turpentine	369
Amidin iodide	417
Ammoniaci c. hydrargyro em-	
plastrum	193, 293
mistura	193
Ammoniæ linimentum	297
spiritus aromaticus	96
fœtidus	187, 188
Ammoniacum	191
African	195

	PAGE.		PAGE.
Ammoniacum distinctive test	192	Anethi aqua	183
lump	191	fructus	182
Persian	195	oleum	183
tear	191	Anethol	180, 181
Ammonium benzoate	250	Anethum graveolens	182
Amomum melegueta	386	Angelic acid	186
Ampelidaceæ	98	Angiospermia	1—364
Amygdalæ amaræ	149	Angosturæ cortex	86
„ oleum	150, 151	Aniseed	21, 181
dulces	148	Anise camphor	180, 181
„ mistura	149	oil of	181
„ oleum	152	Anisi essentia	182
pulvis compositus	146, 149	fructus	181
Amygdalæ	148—155	oleum	181
Amygdalin	150, 154, 155	Anisi stellati fructus	20
Amygdalus communis v. amara	149	Anisyl hydride	181
var. dulcis	148	Annelida	435
Amyli glycerinum	417	Annulated ipecacuanha	219
mucilago	417	Annuloida	434
Amylum	416	Annulosa	435
Amырidaceæ	104—110	Anodyne liniment	52
Amysin	105	Anthelmintic (derivation)	16
Anacanthini	453	Anthemidis extractum	229
derivation of	453	flores	227
Anacardiaceæ	110, 356	infusum	229
Anacardium occidentale	111	oleum	228, 229
Anacyclus	225	Anthemis	225
Anacyclus officinarum	227	nobilis	227
pyrethrum	226	Antimonii tartarati unguentum	447
Analysis of cinchona alka-		Antimonium tartaratum	220
loids	216—218	Apetalæ	310
Analysis of milk	463, 464	Apiaceæ	174—195
Anamirta cocculus	27	Apis mellifica	441, 445
Anarcotine	40, 50	Apocodeia, apocodeine	41
Anarthropoda	435	Apomorphia, apomorphine	41, 48
derivation of	435	Aporetin	312
Anchusa tinctoria	297	Aqua anethi	183
Anethene	183	aurantii floris	94

	PAGE.		PAGE.
Aqua camphoræ	325	Artanthic acid	320
carui	179	Artemisia	225, 232
cinnamomi	328	cina	232
fœniculi	180	maritima	232
laurocerasi	155	Arthropoda	437
menthæ piperitæ	301	Articulata	437
,, viridis	303	derivation of	437
pimentæ	172	Artificial camphor	370
rosæ	159	Artiodactyla	459
sambuci	197	derivation of	459
Arabic acid	145	Artocarpeæ	341
Arabin	104, 107, 117, 145	Asafœtida (assafoetida)	186
Arbutin	244	distinctive test	192
Arctostaphylos uva ursi	243	oil of	187
Areca	412	ordinary (test)	194
Areca catechu	412	pills	157
Areca nut	412	Asafœtidæ enema	188
Aristolochia longa	350	oleum	187
rotunda	350	pilula composita 109, 188, 191	
serpentaria	351	tinctura	189
Aristolochiaceæ	350—352	Asagroeæ officinalis	405
Aristolochiæ radix	350	Ascaris vermicularis	397
Aristolochin	350	Asclepiadaceæ	136, 263
Armoraciæ radix	57	Asparagin	73, 119
spiritus comp.	58, 336	Aspidium filix mas	419
Arnica	225	Astragalus verus	115, 116
montana	230	Atis	8
root	230	Atropa belladonna ..	276, 277, 281
Arniciæ flores	231	Atropaceæ	276—288
radix	230	Atropia	278, 279
tinctura	231	sulphate	123
Arnicin	231	Atropiæ liquor	280
Aroideæ	414	unguentum	280, 461
Arrowroot, Bermuda	392	sulphas	280
Natal	392	sulphatis liquor	280
St. Vincent	392	Attar of roses	159
Artanthe adunca	320	Aurantiaceæ	91—98
elongata	320	Aurantii cortex	91

	PAGE.
Aurantii flores	93
floris aqua	94
,, syrupus.....	94
florum oleum.....	94
infusum	92, 170
,, comp.....	92, 95
syrupus	92
tinctura	93
,, recentis	93
vinum	93
Aves	455

B

Bael	97
Balausta	172, 173
Balaustini flores	173
Balm, common.....	305
Balsam of Canada	375
coast	129
of copaiba	71
Friar's	249
gurjun.....	71
Maracaibo	142
Maranham	141, 142
Para	141, 142
of Peru	127, 129
of tolu.....	126
special distinctive tests ..	128
Balsamodendron mukul	109
myrrha	106
pubescens	109
Balsamum dipterocarpi	71
Peruvianum	129
Tolutanum	126
Banbury.....	315
Barbadoes aloes	396, 397
Barbaloin	397, 398

	PAGE.
Barberry, common	30
Bark, alder	103
Angostura	86
bebeeru	330
canella	63
Carthagenae	215
cassia	329
cinchona (pale).....	213
,, (red)	214
cinnamon	326
Columbian.....	215
cusparia	86
elm	336
flat yellow	200
greenheart	330
larch	364, 365
oak	353
Pitayo	216
quill yellow	201
simaruba root	99
willow	362
Winter's	19, 20, 64
yellow cinchona	200
Barley, pearl.....	415
Barosma betulina.....	87
camphor.....	88
crenulata	87
serratifolia	87, 88, 89
Barracco juice	120
Basidia	425
Bassorin	73, 107, 117
Batavian rhubarb.....	315
Baume de vie	400
Bay berries	28, 330
Bdellium	109
special distinctive tests ..	109
Bean ordeal	122
Beans, calabar	122

	PAGE.
Beans, St. Ignatius	256
Bearberry leaves	243
Beaver, the	468
Beberia	23, 331
Beberia sulphas	331, 332
Bebeeru bark	23, 330
Belæ extractum liquidum	98
fructus	97
Belladonna root	276
Belladonnæ emplastrum	281, 375
extractum	281
folia	281
linimentum	278
radix	276
succus	281
tinctura	282
unguentum	282, 461
Bengal kino	125
Benjamin gum	247
Benzoate ammonium	250
Benzoated lard	461
Benzoic acid	248, 250
aldehyd 150, 151, 154, 155, 328	
Benzoin	247, 461
Siamese	248, 249
Sumatra	248, 249
Benzoini tinctura composita	
128, 249, 362, 402	
Benzoinum	247
Benzoyle cyanide	121
Benzyl cinnamate	130
Berberia	23
Berberidaceæ	11, 29, 30
Berberine	12, 13
Berberis Asiatica	30
vulgaris	30, 173
Bergamot, oil of	97
Berries, buckthorn	101

	PAGE.
Betel nut	412
Bigaradia Citrus	91, 93
Bikh	9
Bile	465
Birthwort	350
Bish	9
Bismuthi trochisci	159
Bissa bö1	106
Bitter almonds	149
essential oil	150, 151, 328
Bitter orange peel	91
Bitter sweet	273
Black catechu	146
cohosh	16
hellebore root	13
mustard seed	59, 60
pepper	316
snake root	16
Blistering paper	450
Blood root	56
Blue cohosh	30
litmus paper	424
Blumea balsamifera	326
Bombay mastich	112
senna	136
Bomboloes	324
Boracis mel	443
Boraginaceæ	297—298
Borneo camphor	325, 326
Borneol	184, 325
Bos taurus	465
Boswellia	104
Botany Bay kino	126, 169
Brassicaceæ	57—62
Brayera anthelmintica	155
Brazil wood	132
Brazilin	132
Bread guarana	76

	PAGE.
Brodie, Sir B.	26
Broom tops	113, 114
Brucia	86, 252, 254
Buchu folia	87
infusum	88
tinctura	88
Buckbean	262
Buckthorn berries.....	101
juice	102
Bugbane	16
Burgundy pitch	376, 377
Burseraceæ	104—110
Butea frondosa	125
gum.....	125
Butter, cocoa	75, 76
kokum	71
Buxus sempervirens	173, 245
Byttneriaceæ	74—76

C

Cabbage rose leaves	158
Cacumina sabinæ	367
scoparii	113
Cacao	74
Cadmii iodidi unguentum	447
Caducous calyx	31
Cæsalpinieæ	113—130
Caffeine	75, 77
Cajuput, oil of	168
Cajuputene, hydrate of	168
Cajuputi oleum.....	168
spiritus	163
Cake gamboge	70
saffron	394
Calabar beans	122, 277
Calami aromatici rhizoma	414
Calamus draco	413

	PAGE.
Calcarate calyx	15
Calcis linimentum	297
Calcium santionate	234
Calefaciens emplastrum.....	375, 447
Calicut	389
Callitris quadrivalvis	373
Calumba	22, 23
Calumbæ extractum	23
infusum	23
radix	21
tinctura	24
Calumbic acid	23
Calumbin	22, 23
Calycifloræ	101—195
Cambogia	68
Cambogiæ pilula composita 70, 399	
Camphor	323
anise	180, 181
artificial	370
Barosma	88
Borneo	325, 326
Dryobalanops	224
Ngai	326
sassafras	333
Camphoræ officinarum	323
Camphoræ aqua	325
,, linimentum 293, 300, 325	
compositum	325
tinctura composita 54, 182,	
250, 325	
Camphoretic acid.....	324
Camphoric acid	324
Campylospermeæ.....	174
Canada balsam.....	375
special distinctive test....	376
Canarium commune	105
Cane sugar	442, 444
Canella alba	19, 63

	PAGE.
Canella bark	63
Canellaceæ	63, 64
Canellæ albæ cortex	63
Cannabene	338
hydride	338
Canna edulis	392
speciosa	392
Cannabinaceæ	337—340
Cannabis indica	337
indicæ extractum	339
sativa	337
Cannon-ball opium	37
Cantharidin	449
Cantharidis acetum	449
emplastrum	375, 447, 450, 468
tinctura	451
unguentum	293, 447, 451
Cantharis vesicatoria	448
Caoutchin	246
Caoutchouc	40
Cape aloes	402
Caprifoliaceæ	196—197
Capsaicin	275
Capsicin	275
Capsici fructus	274
tinctura	276
Capsicum	273, 274
annuum	276
fastigiatum	274
tincture of	275
Capsulæ papaveris	32
Cáyenue pepper	274
Caramania gum	117
Cardamomi, tinctura com-	
posita	98, 179, 329, 388, 439
Cardamomum	387
Aleppy shorts	389
Madras short-longs	389

	PAGE.
Cardamomum Malabar shorts ..	389
Carded cotton	74
Cariceæ	341
Carinatae	456, 457
Carmine	439
Carminic acid	439
Carolina pink root	256
Carragheen	429
Carraway fruits	178
oil of	179
Carthagenæ bark	215
Carthamus	225
tinctorius	394
Carui aqua	179
fructus	178
oleum	179
Carum carui	178
Carvene	179
Carvol	179, 183
Caryophylli	169
infusum	170
Caryophyllin	169
Caryophyllus aromaticus	169
Cascarillæ cortex	342
infusum	343
tinctura	344
Cascarillin	343
Casein	456
Cashew nut	111
Cassia acutifolia	133
angustifolia	136, 137
Brasiliana	138
elongata	136
horse	138
lanceolata	133
obovata	133
purging	137
Cassia cortex	329

	PAGE.
Cassiæ fistulæ fructus.....	137
pulpa	138
Cassiæ grandis fructus	138
Castor fiber	468
oil	347
detection in balsam copaibæ	142, 143
Italian	348
Castorei tinctura	469
Castoreum.....	468
Castorin	469
Cataplasma conii.....	177
fermenti	78
lini	78, 292
sinapis	62
Catechu, black	146
pale.....	197
Catechin	147
Catechu, acacia	146, 198
infusum	199, 329
nigrum	146
pallidum.....	197
pulvis compositus 67, 125,	199, 329, 336
tinctura	199, 329
trochisci.....	199
Catechuic acid	147
Catechu-tannic.....	147
Cathartic acid	134
Cathartogenic	134
Catharto-mannite	134
Caulophyllum thalictroides ..	30
Cellulose	74
Centaury, small	262
Cephaëlis ipecacuanha	218
Cera alba	445
flava	445
Cerati saponis, emplastrum	293, 447

	PAGE.
Cerolein	446
Ceryl cerotate	446
Cetacea	458
Cetacei unguentum..	152, 447, 459
Cetaceum	458
Cetin	458, 459
Cetraria islandica.....	421
Cetrariæ decoctum	422
Cetraric acid.....	422
Cetyl palmitate.....	459
hydrate	459
Cevadilla	405
Ceylon moss	430
Chamomile flowers	227
oil of	228
Charta epispastica 292, 375,	376, 447, 450, 459
sinapis	62, 246
Cherry laurel	90
leaves.....	154
Chian turpentine	111
Chillies	274
Chinese galls	356
musk	467
Chirata	260
Chiratæ infusum	262
tinctura	262
Chiratin	261
Chiretta	260
Chærophyllum anthriscus	178
Cholesterin	427, 469
Cholic acid	466
Chondodendron tomentosum..	24
stems	27
Chondrus crispus.....	429
Christison, Sir R.....	69
Christmas rose	13
Chrysamic acid	398

	PAGE.		PAGE.
Chrysanthemum parthenium..	229	Cinnamomi aqua	328
Chrysobalanæ.....	147	cortex	326
Chrysophanic acid	134, 312	oleum	327
Cichorium	225	pulvis compositus ..	328, 391
Cimifugæ radix.....	16	tinctura	329
Cimicifugin	17	Cinnamomum camphora.....	323
Cinchona alkaloids analysis		cassia	329
216—218		zeylanicum	326
tests	202, 203	Cinnamon bark.....	326
bark (red)	214	oil of	327
„ (yellow).....	200	wild	63
Calisaya.....	200	Cinnamyl hydride	328
condaminea (var.)	213	Cinnyl hydrate	361
lancifolia	215	Cissampelos pareira	24, 26, 27
macrocalyx	213	Citric acid	96
officinalis	213	Citroptene	96
„ test	214	Citrullus colocynthis	165
Pitayensis	216	Citrus aurantium.....	91, 93
succirubra	214	bergamia	97
Cinchonaceæ	197—222	limetta	97
Cinchona-red	207, 333	limonum	94
Cinchonæ flavæ cortex	200	vulgaris, var. Bigaradia	91, 93
decoctum	207, 208	Claviceps purpurea	425
extractum liquidum.....	208	Cloves	169
infusum	208	Club-moss	420
tinctura	208	Coarse gamboge	70
pallidæ cortex	213	Cocci tinctura	439
tinctura composita	214,	Cocculus indicus	27, 330
352, 389, 394, 439		palmatum	22
Cinchonia	205	Coccus	438—440
Cinchonicine.....	206	cacti	438
Cinchonidia	205	Cochineal	438—440
Cincho-tannic acid	206, 207	Honduras	439
Cinnamein.....	130	Mexican	439
Cinnamic acid.....	127, 128, 328	Vera Cruz	439
aldehyd	328, 361	Teneriffe	439
Cinnamodendron corticosum ..	20	Cochinellin	439
Cinnamol	328	Cochlearia armoracia	57

	PAGE.
Cochlearia distinctive character	58
Cocoa butter.....	75, 76
nibs.....	74
Codeia, Codeine	40, 48, 49
syrup of	49
Cod liver oil	453
Cælenterata	434
Cælospermeæ	174
Cohosh, blue.....	30
Colchicia	408
Colchici cormus	407
extractum	409
,, acet	409
semina	409
tinctura	410
vinum	409
Colchicine	411
Colchicum autumnale.....	408
seeds	409
Coleoptera.....	448
derivation of	448
Collodion	74
Collodium flexile	349, 376
Colocynth Levant	165
Mogadore	166
pulp.....	165
Colocynthein.....	167
Colocynthis extractum com-	
positum	167, 389, 402
et hyoscyami pilula	167,
170, 271, 399	
pilula composita	167, 170,
271, 399	
pulpa	165
Colocynthin	166
Columbian bark	215
Colza oil.....	294
Comfrey	298

	PAGE.
Commercial varieties opium ..	36
Compositæ	225—240
Confectio opii 51, 117, 179, 317, 391	
piperis	179, 317, 443
rosæ caninæ	160
gallicæ	157
scammonii 170, 179, 271,	
391, 443	
sennæ 120, 134, 138, 140,	
153, 184, 341	
terebinthinæ.....	120, 370, 443
Conia	175
Coniæ vapor	177
Coniferæ	364
Coniferin	385
Conii cataplasma	177
extractum	177
folia.....	176
fructus	174
pilula composita	177, 221
succus.....	177
tinctura	176
Conium	174
maculatum	174
Constantinople opium.....	36
Convolvulaceæ	264—272
Convolvulin	266, 268
Convolvulus scammonia..	268, 270
Copaiba (copaiva)	71, 140
special distinctive tests	
141. 142	
Copaibæ oleum	143
Copaifera multijuga.....	140
Copaivic acid	141
Copper, Fehling's solution	442, 443
Coriander fruit	183
oil of	184
Coriandri fructus	183

	PAGE.
Coriandri oleum	184
Coriandrum	174
sativum	183
Coriaria myrtifolia	136
Corigliano juice	120
Corollifloræ	195—309
Cortex angosturæ	86
aurantii	91
canellæ albæ	63
cascarillæ	342
cassiæ	329
cinchonæ flavæ	200
,, lancifoliæ	215
,, pallidæ	213
,, rubræ	214
cinnamomi	326
cuspariæ	86
frangulæ	103
granati	30
,, radicis	172
laricis	364, 365
limonis	94
mezerii	321
nectandræ	330
pruni virginianæ	153
quercus	353
salicis	362
simarubæ radicis	99
ulmi	336
winteranus	19
winteri	19
Corylaceæ	353
Cotarnia	50
Cotton, carded	74
wool	74
Cowhage	122
Cow-itch	122
Creasol	372

	PAGE.
Creasoti mistura	372
unguentum	372, 447
vapor	372
Creasotum	371, 372
derivation of	371
Cretæ mistura	146
c. opio pulvis	449
pulvis aromaticus	329, 336,
389, 394	
c. opio pulvis	53
Croci tinctura	394
Crocin	394
Crocus	393
sativus	393
Croton eluteria	342
lucidus	344
oil	345
tiglium	344
Crotonic acid	345
Crotonis linimentum	168, 345
oleum	345
Cruciferae, vel. Brassicaceæ	
	57—62
Cryptogamia	418—430
Cryptopia	41
Cubeb-camphor	319
Cubeba officinalis	318
Cubebæ oleum	319
tinctura	319
Cubebene-hydrate	319
Cubebin	319
Cubebs	318
oil of	319
Cucurbitaceæ	162—167
Cucurbiteæ	162
Cumin fruit	184
Cumini fructus	184
Cuminic aldehyd	185

	PAGE.
Cuminum cyminum.....	184
Cupuliferæ	353—360
Curcuma angustifolia	392
longa	386
Curcumæ rhizoma	386
Curcumin	387
Curd soap	292
Curvembryæ	273
Cuspariæ cortex	86
Cuspariæ infusum	87
Cusparin	86
Cusso	155
infusum	156
Cutch.....	146, 147, 198
Cydoniæ decoctum	162
semina	161
Cymol.....	185
Cynanchum	136
Cynarrhodum, <i>vide</i> Rosacæ..	147

D.

Daphne laureola	321, 322
mezereum	321, 322
Daphnetin	322
Daphnin.....	322
Datura stramonium.....	282, 283
tatula	284
Dāturia	283
Daucus	174
Dead Sea apples	355
Decoctum aloes	108, 120
compositum	394, 400
cetrariæ	422
cinchonæ flavæ	207, 208
cydoniæ	162
granati radicis	173
hæmatoxyli	132, 329

	PAGE.
Decoctum hordei	416
papaveris	33
pareiræ	25
quercus	354
sarsæ	382
sarsæ compositum 81, 120,	333, 382
scoparii	115
taraxaci	237
ulmi	337
Delphinia	16
Delphinium staphisagria.....	15
Derivation of anacanthini	453
anarthropoda.....	435
artiodactyla	459
coleoptera	448
hemiptera	438
homoptera	438
hymenoptera	441
ichthyopsida	453
pachydermata	460
protozoa	433
rhizopoda	433
sauropsida	455
teleostei	453
Detection of umbelliferous gum-	
resins	193
Dextrose	442
Dicotyledones	1—378
Dictyogenæ	379—384
Digitalin	307, 308
Digitalinum	307, 308
Digitalis folia	306
infusum	309
purpurea	306, 307
succus.....	309
tinctura	309
Dill fruit.....	182

	PAGE.
Dill, oil of	183
Dinitro-cellulose	74
Diplolepis gallæ tinctoriæ	354
Dipterocarpeæ	71
Dipterocarpi balsamum	71
Dipterocarpus turbinatus	71
Dissepiments spurious.....	137
Distinctive tests (special)	
ammoniacum African	195
Distinctive tests (special)	
ammoniacum Persian	195
asafoetida	194
,, from ferula alliacea	194
balsam tolu	128
bdellium	109
benzoin	249
Canada balsam.....	376
copaiba	141, 142
dragon's blood	413
elemi	105
galbanum Levant.....	194
,, Persian.....	194
guaiaicum	83
mastich	111, 112
myrrh.....	108
olibanum	195
opoponax	195
resin	374
sagapenum Levant	194
,, Persian	193
umbelliferous gum-resins	193
Dorema ammoniacum.....	191
Dover's powder.....	53
Draconis sanguis	413
Dragon's blood	413
Drimys winteri.....	19, 20
Dryobalanops aromatica.....	325
camphor	224

	PAGE.
Dulcamara.....	273
Dulcamaræ infusum	274
Dulcamarine	274
Dutch trimmed rhubarb	315

E

East Indian myrrh	106, 108
opium	37
rhubarb	315
senna	136
tamarinds	139
Ecballium elaterium	162
officinæ	162
Ecbolia	427, 428
Egg, white of	456
yolk of	457
Egyptian opium	37
Elaterii pulvis compositus	165
Elaterin.....	164, 165
Elaterium	162, 164
Elder flowers.....	196
ointment of	197
'wine	197
Elecampane root	234, 235
Electuary lenitive.....	134
Elemi	105
special distinctive tests ..	105
unguentum	106, 447
Elettaria cardamomum	387
Elm bark	336
Emetine.....	220
Emmenagogue (derivation) ..	14
Emplastrum ammoniaci c.	
hydrargyro	193, 293
belladonnæ	281, 375
calefaciens	375, 447, 450
cantharidis	375, 447, 450, 468

	PAGE.		PAGE.
Emplastrum cerati saponis	293, 447	Essentia anisi	182
„ ferri	377	menthæ piperitæ	302
galbani	190, 193, 447	Essential oil of bitter almonds	
hydrargyri	293, 375	150, 151, 328	
opii	52, 375	of mustard	60, 61
picis	293, 375, 377, 447	Estimation, ipecacuanha	222
plumbi	294, 296	of scammony	272
iodidi	375	Ethal	459
resinæ	374	Etærio, <i>vide</i> Rosaceæ	147
saponis	295, 375	Eucalyptol	168
Empleurum serrulatum	88	Eucalyptus	126
Emulsin	149, 150, 154, 155	globulus	168
English galls	356	resinifera	169
raubarb	315	Eugenia pimenta	171
Enema aloes	398	Eugenic acid	170
asafoetidæ	188	Eugenin	169
magnesiæ sulphatis	293	Eugenol	170
opii	52	Euodici aldehyd	85
tabaci	288	Euphorbia resinifera	346
terebinthinæ	371	Euphorbiaceæ	342—350
Epigynæ 101, 162, 195,		Euphorbium	346
196—242, 384		Euphorbon	346
Epipetalæ	195, 245	Euryangium sumbul	185
Equisetic acid	4	Evernic acid	424
Ergota	425	Exogonium purga	264
Ergot of rye	425—428	Extractum aconiti	7
Ergotæ extractum liquidum	428	aloes Barbadosensis	398
infusum	428	„ Socotrinæ	400
tinctura	428	antheridis	229
Ergotic acid	427	belæ liquidum	98
Ergotine	427	belladonnæ	281
Ericaceæ	242—245	calumbæ	23
Eriocolin	244	cannabis Indicæ	339
Erythræa centaurium	262	cinchonæ flavæ liquidum	208
Erythric acid	424	colchici	409
Erythroretin	312	„ aceticum	409
Eseré nut	122	colocynthis compositum	
Essence of pimento	172	167, 389, 402	

	PAGE.
Extractum conii	177
ergotæ liquidum	428
filicis liquidum	419
gentianæ	259
glycyrrhizæ	119
„ liquidum	119
hæmatoxyli	132
hyoscyami	286
jalapæ	266
kramerizæ	67
lactucæ	240
lupuli	340
mezerii æthereum	323
nucis vomicæ	255
opii	52
„ liquidum	52
papaveris	33
pareiræ	26
„ liquidum	26
physostigmatis	124
quassiæ	100
rhei	313
sarsæ liquidum	382
stramonii	283
taraxaci	238

F

Fabaceæ	113—147
Fabæ physostigmatis	122
Farina lini	78
tritici	416
Fehling's solution of copper	442, 443
Fel bovinum purificatum	465, 466
Fennel fruits	179
oil of, bitter	180
„ Saxon	180

	PAGE.
Fennel, oil of, sweet	180
Fenugreek	115
Fermenti cataplasma	78
Ferri emplastrum	377
iodidi pilula	120
et quinz citras	212
mistura aromatica 23, 170, 214	
composita	108, 159
Ferrum redactum	125
Ferula alliacea	194
galbaniflua	189
Ferulaic acid	188
Feverfew	229
Ficus carica	341
Figs	341
Filices	418
Filicic acid	419
Filicis extractum liquidum	419
Filix mas.	419
Fleming's tincture	8
Fleurs de sureau	196
Florence oil	292
Flores anthemidis	227
arnicæ	231
aurantii	93
balaustini	173
koso	155
sambuci	196
Flowers, chamomile	227
elder	196
orange	93
Flückiger, Professor	390
Fly, Spanish	448
Fœni-græci semina	115
Fœniculum dulce	179
Fœniculi aqua	180
fructus	179
Folia aconiti	2

	PAGE.
Folia buchū	87
conii	175
digitalis	306
hyoscyami	284
laurocerasi	154
maticæ	320
stramonii	282
tabaci	287
uvæ ursi	243
Fools' parsley	178
Foxglove	306
Frangulæ cortex	103
Frankincense	104
Frankincense (common)	375
Fraxin	290
Fraxinus Ornus	289
rotundifolia	289
Friar's balsam	249
Fruit carraway	178
coriander	183
cumin	184
dill	182
fennel	179
hemlock	174
mango	111
sugar	442
Fructus anethi	182
anisi	181
belæ	97
capsici	274
carui	178
cassiæ fistulæ	137
„ grandis	138
conii	174
coriandri	183
cumini	184
fœniculi	179
juniperi	366

	PAGE.
Fructus lauri	330
mangosteen	97
pruni	153
rhamni	101
rosæ caninæ	160
Fumaric acid	422
Fungi	425—428

G

Gadus morrhua	453
Galangal	391
Galbani emplastrum	193, 447
oleum	190
Galbanum	189
distinctive test	192
Levant	194
Persian	194
Galipea cusparia	86
Gallate of veratria	407
Gallæ	354
tinctura	356
unguentum	356, 461
„ c. opio	56, 356
Gallic acid	359
Gallipoli oil	292
Galls Aleppo	355
Chinese	356
English	356
Mecca	355
Mosul	355
Gallus bankiva var. domestica	456, 457
Gambia kino	125
Gambir (Gambier)	197
Gamboge	68
adulteration	70
cake	70

	PAGE.
Gamboge coarse	70
lump	70
pipe	70
special distinctive tests, 69, 70	
Garcinia indica	71
mangostana	97
morella (<i>v. β. pedicellata</i>)	68
Garciniæ oleum	71
Gelatin	454, 456
,, solution of	455
Genoa oil	292
Gentian	258
Gentiana lutea	258
pannonica	260
punctata	260
purpurea	260
Gentianaceæ	257—262
Gentianæ extractum	259
,, infusum compositum	
93, 95, 259	
mistura	93, 184, 259
radix	258
tinctura composita	93, 260, 389
Gentianic acid	259
Gentiogenin	258
Gentiopicrin	258
Gerrard, Mr.	90
Geum urbanum	231
Ginger	389
African	391
Bengal	391
Cochin	391
Jamaica	391
Ginseng American	66
Glucose	61, 442, 444
Glumaceæ	415—417
Gluten	416
Glycerinum acidi gallici	360

	PAGE.
Glycerinum acidi tannici	358
amyli	417
Glycyrretin	119
Glycyrrhiza glabra	118
Glycyrrhizæ extractum	119
,, liquidum	119
pulvis compositus	120
radix	118
Glycyrrhizin	119
Glyceryl hydrate	294
oleate	294, 461
Glycocholic acid	465, 466
Glycocin	466
Gomme Sirop de	146
Gossypium	74
Gracilaria Lichenoides	430
Grains of Paradise	386
Graminaceæ	415—417
Grana Paradisi	386
Granati cortex	30
radicis cortex	172
,, decoctum	173
Granilla	440
Grape sugar	442
Greenheart bark	330
Green hellebore root	411
Guaiaci lignum	80
mistura	83, 146
resina	81, 84
tinctura ammoniata	84
Guaiaconic acid	82
Guaiacum, lump	81
officinale	80, 81
Peruvian	83
sanctum	80
in tears	81
wood	80
Guaiaretate potassium	82

	PAGE.
Guaiaretic acid.....	82
Guarana	76
bread	76
Guimauve, racine de	72
Guinea pepper	274
Gulancha	29
Gum Arabic	143, 144
Benjamin	247
butea	125
caramania	117
kino.....	124
kordofan.....	144
moussul	117
Senegal	144
suakin.....	144
talca	144
tragacanth.....	115, 116
Turkey (picked)	144
wattle	144
Gummi acaciæ	143
Gum-resins, umbelliferous, De- tection of	193
Gun cotton	74
Gurjun balsam	71
Gutta-percha	245, 246
liquor	246
Guttiferæ	68—71
Gymnospermia	364—378

H

Hagenia Abyssinica	155
Hamamelideæ	360
Hanbury, Daniel 24, 106, 341, 395	
Hard soap.....	292, 295
Hay saffron	394
Helenin	235
Helicoid inflorescence.....	297

	PAGE.
Hellebore (black) root.....	13
(green) root	411
(white) root	410
Helleborein	14
Helleboresin	14
Helleboretin	14
Hellebori radix nigri	13
Helleborin	14
Helleborus niger	13
Hemidesmi radix	263
syrupus	263
Hemidesmus Indicus	263
root.....	263
Hemlock fruit	174
leaves	176
Hemp, Indian	337
seed oil	80, 294
Henbane	284
Hepatic aloes	400
Hesperidin.....	92
Hesperidium.....	91, 94
Hiera Picra	64
Hips	160
Hirschsohn, E. (umbelliferone)	
193—195	
Hirudinea	436
Hirudo	436
Hemiptera	438
derivation of	438
Hæmatein	131
Hæmatoxyli decoctum ..	132, 329
extractum	132
lignum	130
Hæmatoxylin	131, 132
Hæmatoxylon campechianum	
130, 131	
Hæmostatic (derivation of) ..	320
Hog tragacanth	117

	PAGE.
Homoptera	438
derivation of	438
Honduras cochineal	439
Honey	441—445
Honeydew	425
Hop	339
Hordei decoctum	416
Hordeum decorticatum	415
distichon	415
Horehound	305
Horse cassia	138
turpentine	369
Horseradish root	57
Huile de cade	367
de petit grain	94
Humulus lupulus	339
Huxham's tincture of bark	214
Huxley, Professor	452
Hydragogue (derivation of)	102
Hydrargyri ammoniati unguen-	
tum	447
emplastrum	293, 375
iodidi rubri unguentum	447
linimentum	297
nitratis unguentum	293, 461
oxidi rubri unguentum	152, 447
pilula	120
subchloridi pilula com-	
posita	84, 349
unguentum	461, 468
compositum	293, 325, 447
suppositoria	76, 447
Hydrastin	18
Hydrastis Canadensis	18
radix	18
Hydrate of cajuputene	168
Hydrocyanic acid, tests	151, 152
Hydroquinone	206, 244

	PAGE.
Hymenoptera	441
derivation of	441
Hyoscina	285
Hyoscinic acid	285
Hyoscyami extractum	286
folia	284
succus	286
tinctura	287
Hyoscyamia	285
Hyoscyamia, sulphate	285, 286
Hyoscyamus niger	284
Hyphæ	425
Hypogynæ	395
Hypostamineæ	195, 242—245

I

Ibo (river)	21
Iceland moss	421
Icthyocolla	454
Icthyopsida	452—455
derivation of	452
Igasuria	252
Ignatii semina	256
Illicium anisatum	20
Incense	248
Indian aconite	9
bdellium	109
hemp	337
pink root	256
sarsaparilla	263
Indigo	120
sulphate solution	121
Indigofera	120
tinctoria	121
Indigotin	121
Infusum anthemidis	229
aurantii	92

	PAGE.
Infusum aurantii compositum	92, 95, 170
buchu	88
calumbæ	23
caryophylli	170
cascarillæ	343
catechu	199, 329
chirataë	262
cinchonæ flavæ.....	208
cuspariæ.....	87
cusso	156
digitalis	309
dulcamaræ	274
ergotæ	428
gentianæ compositum	93,
	95, 259
krameriæ	67
lini	78, 120
lupuli	340
maticæ	321
quassiæ	100
rhei	313
rosæ acidum	158
senegæ	65
sennæ.....	135, 391
serpentariæ	352
uvæ ursi	244
valerianæ	224
Injectio morphiæ hypodermica	47
Inosite	237
Insecta	437
Inula	225
helenium	234
Inulæ radix	234
Inulin.....	235, 237
Iodide amidin	417
Iodidi unguentum	461
Ipecacuanha	219

	PAGE.
Ipecacuanha annulated	219
estimation	222
New Granada	219
striated	222
undulated	222
wine	221
Ipecacuan	218
Ipecacuanhæ, acetum.....	221
pulvis compositus	53, 221
radix	218
cum scillâ pilula 53, 193,	
	221, 404
trochisci	221
vinum	221
Ipecacuanhic acid	220
Ipomœa orizabensis.....	267
simulans.....	268
Ipomœic acid	266
Iridaceæ.....	393—395
Irish moss.....	429, 430
Iron carbonate pills.....	157
Isinglass	454
Isonandra gutta	245
Italian castor oil	348
Izmid.....	36

J

Jaborandi	89, 90
Jalap	264
tampico	268
Jalapæ extractum	266
pulvis compositus....	267, 391
resina	265
tinctura	267
Jalapin	266, 271
Jamaica pepper	171
Jateorhiza calumba	22

	PAGE.
Jateorhiza Miersii	22
palmata.....	21, 22
Jecoris aselli oleum	453
Jeffersonia diphylla	30
Jervia	411
Jervine	411
Jordan almonds	148
Juice barracco	120
buckthorn	102
corigliano	120
lemon	96
,, artificial.....	96
pignatelli	120
solazzi	120
Spanish	120
Juniper oil	367
tar	367
Juniperi fructus	366
oleum	367
empyreumaticum	367
spiritus	367
Juniperus communis	366
oxycedrus	367
sabina	367, 368

K

Kamala	349
Kino	124
Bengal	125
Botany Bay	126, 169
Gambia	125
pulvis compositus 54, 125, 329	
red	125
tinctura	125
Kinic acid	206
Kino-tannic acid	125
Kokum butter	71

	PAGE.
Kordofan gum	144
Kosin	156
Koussin	156
Koussou	155
flores	155
Krameria tomentosa	67
triandra	66
Krameria extractum	67
infusum	67
radix	66
tinctura	67

L

Labiatae	298—305
Labiatiflorae	225
Lac	463
Lacmus	422
Lactis saccharum.....	465
Lactuca	225
virosa	238
Lactucæ extractum	240
Lactucarium	239
Lactucerin	237, 239
Lactucin	239
Lævulose	442
Lake	439
Larch bark	364, 365
Lard, benzoated	461
prepared.....	460
Laricis cortex	364, 365
tinctura	365
Larix Europæa.....	364, 366
Larixin	365
Laudanum.....	55
Lauraceæ	323—333
Laurel, cherry	90
spurge.....	321

	PAGE.
Lauri fructus.....	330
Lauric aldehyd.....	85
Laurocerasi aqua.....	155
folia	154
Laurus nobilis	330
Lavandula spicata	300
vera	298
Lavandulæ oleum	298, 299
spiritus	300
tinctura composita 300,	
305, 329, 336	
Lavender compound tincture..	126
Lead with opium pills.....	157
Lead plaster	292
Leaves, bearberry	243
cabbage rose.....	158
cherry laurel	154
hemlock	176
matico	320
rose	156
stramonium	282
Leech, the.....	436
Lefort, M.....	220
Leguminosæ	113—147
Lemon juice	96
artificial	96
oil of	95
peel.....	94
Lenitive electuary	134
Leontodon hispidus.....	238
taraxacum	236
Lettuce-opium	239
Levant colocynth.....	165
galbanum	194
nut	27
sagapenum	194
Lewa	37
Lichenes	421—424

	PAGE.
Lichenin	422
Lignum guaiaci	80
hæmatoxyli	130
pterocarpï	126
quassiæ	99
santali	352
vitæ	80
Ligulifloræ.....	225
Liliaceæ	395—404
Lime-juice	97
Limonis cortex.....	94
oleum	95
succus.....	96
saturating power	97
syrupus	95
tinctura	95
Linaceæ	77—80
Lini cataplasma.....	78, 292
farina	78
infusum	78, 120
oleum	79
semina	77
Linimentum aconiti	8
ammoniæ	297
anodyne.....	52
belladonnæ	278
calcis	297
camphoræ	293, 325
,, compositum 300,	325
crotonis	168, 345
hydrargyri	297
opii	52
potassii iodidi c. sapone	
96, 295, 296	
saponis	292, 296
sinapis compositum 62,	305,
322, 349	
terebinthinæ	371

	PAGE.
Linimentum terebinthinæ	
aceticum	371
Linoleic acid	80
Linoxyn	80
Linseed	77
meal	78
oil	79, 294
Linum usitatissimum	77
Liquidambar altingiana	362
orientale	360
Liquor atropiæ	280
,, sulphat.	280
epispasticus	451
gutta-percha	246
morphiæ acetatis	46
,, hydrochlorat	44
strychniæ	251, 255
Liquorice root	118
Litmus	422
neutral	424
paper, blue	424
,, red	424
tincture of	424
Lobelacrin	242
Lobelia	240
inflata	240
Lobeliaceæ	240—242
Lobeliæ tinctura	242
,, ætherea	242
Lobelina	241
Loganiaceæ	251—257
Logwood	130
Long pepper	318
Lucca oil	292
Lump gamboge	70
guaiaicum	81
Lupuli extractum	340
infusum	340

	PAGE.
Lupuli tinctura	340
Lupulin	340
Lupulus	339
Lycopodiaceæ	420
Lycopodium	420

M

Macroton	17
Magellan, Straits of	19
Magnesiæ sulphatis enema	293
Magnoliaceæ	18—21
Male fern	419
Malva sylvestris	73
Malvaceæ	72—74, 77
Mammalia	458—469
Mandrake	10
Mango fruit	111
Mangofera Indica	111
Mangosteen fruit	97
oil	71
Manna	289—291
Mannite	290
Maracaibo balsam	142
Maranham balsam	141, 142
Maranta arundinacea	392
Marantaceæ	392
Marantæ amylum	392
Marjoram	305
Marrubium vulgare	305
Marshmallow root	72, 78
Marsh trefoil	262
Mastiche (Mastich)	110
Alexandrian	112
Bombay	112
solutions	111
special distinctive tests	

	PAGE.
Masticin.....	111
Matico leaves	320
Maticœ folia	320
infusum	321
Matricaria.....	225
chamomilla	229
Matto grosso (Province).....	218
Maw seed	34
May apple.....	10
Meadow sweet, oil of	181
Meal linseed.....	78
Mecca galls	355
senna	136
Meconia, Meconine.....	40, 49
Meconic acid	40, 51
Mel	441—445
boracis	443
depuratum	443
Melaleuca minor	168
Melampodii radix.....	13
Melanthaceæ	405—412
Melissa officinalis.....	305
Melissyl palmitate	446
Menispermaceæ	21—29
Menispermin.....	29
Mentha piperita	300, 301
pulegium	303
viridis	302
Menthene	301
Menthæ piperitæ aqua	301
„ essentia	302
„ oleum ..	300, 301
„ spiritus	302
pulegii oleum	303
viridis aqua	303
„ oleum	302
Menthol.....	301, 303
Menyanthes trifoliata.....	262

	PAGE.
Mercury pills	157
vegetable	11
Meta-cinnamein	361
Mexican cochineal	439
Mezerii cortex	321
extractum æthereum	323
Mezereon	321
Micropyle	123
Milk	463
analysis of.....	463
sugar of	465
Milkwort	64
Mimoseæ	113, 143
Mistura ammoniaci.....	193
amygdalæ	149
creasoti	372
cretæ	146
ferri aromatica ..	23, 170, 214
„ composita.....	108, 159
gentianæ	93, 184, 259
guaiaci.....	83, 146
scammonii	269
sennæ composita	120, 135
spiritus vini gallici	457
Mogadore colocynth	166
Mollusca	451
Monochlamydeæ	310—378
Monocotyledones	379—417
Monkshood	2
Moraceæ	341
Mori succus	341
syrupus	342
Morphia, morphine	40, 41—44
chemical behaviour	43
estimation	39
preparation, B.P.....	41
explanation of process....	42
tests	43, 44

	PAGE.
Morphiæ acetas	46
acetatis liquor	46
hydrochloras	36, 44
„ liquor	44
injectio hypodermica	47
suppositoria	45, 76, 447
„ c. sapone	45
trochisci	45
„ et ipecacuanhæ	45, 222
Morrhua oleum	453
Morus nigra	341
Moschus	466
moschiferus	466
Moss Ceylon	430
club	420
Iceland	421
Irish	429, 430
Mosul galls	355
Mother cloves	170
Moussul gum	117
Mucilage, acacia	144
Mucilago acaciæ	145
amyli	417
tragacanthæ	117
Mucro	133
Mucuna pruriens	122
Mucunæ setæ	122
mulberry	341
Mullein	309
Musk	466
Chinese	467
deer	466
Russian	467
Siberian	467
Mustard black seed	59, 60
essential oil of	60, 61
white seed	59
Mycose	427

	PAGE.
Myristica	334
fatua	336
officinalis	334
Myristicaceæ	334—336
Myristicæ oleum	325
expressum	325
spiritus	336
Myristic acid	325
Myristin	79, 325
Myronate potassium	60, 61
Myrosin	58, 59, 60, 61
Myroxylon pereiræ	129
toluifera	126, 127
Myrrh, East Indian	106, 108
Turkey	106
Myrrha	106, 107
special distinctive tests ..	108
Myrrhæ tinctura	108
Myrtaceæ	168—173

N.

Napelline	7
Napthalene	378
Napthalin	250
Narceia, narceine	40, 49, 50
Narcotia, narcotine	40, 50
Narthex asa-fœtida	186
Natal aloes	402
Nataloin	398, 402
Nectandra rodiaei	330
Nectandræ cortex	330
Nectaries	13
Nepal	9
Neroli, oil of	94
Neutral litmus	424
New Granada ipecacuanha ..	219
rhatany	67

	PAGE.
Ngai camphor	326
Nhandirobeæ	162
Nibs cocoa	74
Nicotia	287, 288
Nicotianæ tabacum.....	287
Nitro-cellulose	74
Nuciş vomicæ extractum	255
,, tinctura	255
Nut areca	412
betel	412
cashew	111
eseré	122
pistacia	111
Nuculanium	98
Nut galls	354
Nutmeg	334
oil of (expressed)	325
Nux vomica	251

O

Oak bark	353
Œnanthe	174
Oil, almond	152
anise	181
asafoetida	187
bergamot	97
bitter almonds	128
,, (essential)	
150, 151, 328	
cajuput	168
caraway	179
castor	347
chamomile	228
cinnamon	327
cod liver.....	453
colza	294
coriander	184

	PAGE.
Oil, croton.....	345
cubebæ	319
dill	183
fennel, bitter.....	180
,, Saxon	180
,, sweet	180
Florence	292
galbanum	190
Gallipoli.....	292
Genoa.....	292
hemp seed	80, 294
juniper	367
lemon.....	95
linseed.....	79, 294
Lucca.....	292
mangosteen	71
meadow sweet	181
mustard, essential	60, 61
neroli.....	94
nutmeg, exprcssed	325
,, volatile	325
olive	291
pennyroyal	303
Provence	292
rosemary	304
rue	85
salad	291
savin	368
sesame	294
spearmint	302
spike	300
turpentine	370
virgin	291
Ointment of aconitia	4
elder flowers	197
Olea Europea	291
Oleaceæ.....	289—297
Oleate glyceryl.....	294

	PAGE.
Oleate potassium	295
sodium	295
Oleic acid	80
Olein.....	76, 79, 152, 292. 454
Oleum amygdalæ.....	152
,, amaræ.....	150, 151
anethi	183
anisi	181
anthemidis	228, 229
asafœtidæ	187
aurantii florum	94
cajuputi	168
carui	179
cinnamoni	327
copaibæ	143
coriandri	184
crotonis	345
cubebæ	319
galbani	190
garciniæ	71
jecoris aselli	453
juniperi	367
,, empyreumaticum	367
lavandulæ	298, 299
limonis	95
lini	79
menthæ piperitæ	300, 301
,, pulegii	303
,, viridis	302
morrhuæ	453
myristicæ	325
,, expressum	325
olivæ	152, 291
pimentæ	172
ricini	347
rosæ	159
rosmarini	304
rutæ	85

	PAGE.
Oleum sabinæ	368
sinapis	62
terebinthinæ	369
theobromæ.....	75
Olibanum	104, 195
Olive oil.....	291
tests for	293
Olivæ oleum.....	152, 291
Ophelia chirata.....	260
Ophelic acid	261
Opianic acid	50
Opii confectio 51, 117, 179, 317, 391	
emplastrum.....	52, 375
enema.....	52
extractum	52
,, liquidum.....	52
linimentum	52
pulvis compositus 54, 117,	
179, 317, 391	
tinctura	55
,, ammoniata 55,	
182, 250, 394	
trochisci	55, 120
vinum.....	170, 329
opisthotonos	254
Opium	34—40
antidotes	35
assay	39
cannon-ball	37
chemistry	40
collection	34
commercial varieties	36
Constantinople	36
dose.....	35
East Indian	37
Egyptian	37
Patna garden	37
Persian	37

	PAGE.
Opium pharmacopœial test for	
strength	38
pills	53
Smyrna	36
uses	35
Opoponax	195
Otto of roses	159
Ovi albumen	456
vitellus	457
Ovis aries	467
Ox bile purified	465, 466
Oxylnoleic acid	80
Oxymel	443
scillæ	404, 443
Orange flowers	93
peel, bitter	91
tree	91
Orcein	423
Orchidaceæ	384, 385
Orchis mascula	385
Orcin	423
Ordeal bean	122
Origanum vulgare	305
Orsellic acid	424
Orthospermeæ	174

P

Pachydermata	459, 460
derivation of	460
Pale catechu	197
cinchona bark	213
Palmaceæ	412—414
Palmitic acid	459
Palmitin	76, 292, 454
Panax quinquefolium	66
Papaver Rhœas	30, 31
somniaferum, var. album. 32, 34	

Papaver somniferum var.	
nigrum	32
,, var. setigerum	32
Papaveraceæ	30—57
Papaverine	40, 50, 51
Papaveris capsulæ	32
decoctum	33
extractum	33
semina	34
syrupus	34
Paper blistering	450
blue litmus	424
red litmus	424
turmeric	387
Papilionaceæ	113—130
Para balsam	141, 142
Paramenispermin	29
Paregoric	54
,, Scotch	55
Pareira brava false	26
white	27
yellow	27
Pareiræ decoctum	25
extractum	26
liquidum	26
radix	24
Parillin	380
Special Distinctive Tests	381
Patna garden opium	37
Paullinia sorbilis	76
Pearl barley	415
Peel bitter orange	91
lemon	94
Pelargonic acid	85
Pellitory root (of Spain)	226
Pelosin	25
Pennyroyal, oil of	303
Pepper, black	316

	PAGE.
Pepper	274
cayenne	274
Guinea	274
Jamaica	171
long	318
white	317
Pepsin	462
Peptone	463
Periderm.	200
Perigynæ	101—162
Persian ammoniac	195
galbanum	194
guaiacum	83
opium	37
sagapenum	193
Peruvian balsam	127, 129
rhatany	68
Pepo	162, 163
Petaloidæ	384
Petit grain, huile de	94
Phanerogamia	1—417
Phenyl hydrate.....	363
Phæoretin	312
Phthalic acid	250
Physeter macrocephalus.....	458
Physostigmia	123
Physostigma venenosum.....	122
Physostigmatis extractum	124
fabæ	122
Picis emplastrum	293, 375, 377, 447
liquidæ unguentum	378, 447
Picræna excelsa	99
Picrotoxin	28
Pignatelli juice.....	120
Pills aloes	157
Barbadoes	157
myrrh	157
Socotrine	157

	PAGE.
Pills asafœtida	157
iron carbonate	157
lead with opium	157
mercury	157
opium	53
Pilocarpia	90
Pilocarpus pinnatifolius.....	89
Pilula aloes et asafœtidæ 189, 400	
,, Barbadosensis 179, 399	
,, et ferri.....	399
,, et myrrhæ ..	108,
394, 401	
,, Socotrinæ	401
asafœtidæ composita 109,	
188, 191	
cambogiæ composita 70, 399	
colocynthidis composita	
167, 170, 271, 399	
,, et hyoscyami	
167, 170, 271, 399	
conii composita	177, 221
ferri iodidi	120
hydrargyri	120
subchloridi composita 84, 349	
ipecacuanhæ cum scillâ	
53, 193, 221, 404	
plumbi c. opio	53
quinæ.....	160, 211
rhei composita 109, 302,	
313, 402	
saponis composita....	53, 296
scammonii composita 266, 269	
scillæ composita 193, 391, 404	
Pimenta, Pimento	171
Pimentæ aqua	172
oleum	172
Pimpinella anisum	181
Pinic acid	373

	PAGE.		PAGE.
Pink Carolina root	256	Plumbi suppositoria composita	54, 76, 447, 461
Indian root	256	unguentum	447
Pinus palustris.....	369, 375	Poayeros	218
pinaster	369	Podophylli radix	10
sylvestris	378	resina	12, 17
toëda	369, 375	preparation of	12
Pipe gamboge	70	Podophyllum peltatum	10
Piper album	317	Poison nut.....	251
angustifolium ..	320	Polarimeter	445
caudatum	318	Polychroit.....	394
longum	318	Polygala Senega	64
nigrum	316	Polygalaceæ.....	64—68
officinarum.....	318	Polygalic acid	65
Piperaceæ.....	316—321	Polygonaceæ	310—315
Piperia	316	Pome	161
Piperic acid	317	Pomeæ	161
Piperidia	317	Pomegranate root bark	172
Piperine	317	Poppy heads	32
Piperis confectio	179, 317, 443	Porifera	433
Pisces	452	Portland, Duke of, powder....	350
Pistacia lentiscus.....	110	Potassæ sulphuratæ unguentum	461
nut	111	Potassii iodid unguentum	461
terebinthus	111	c. sapone linimentum	96, 295, 296
vera.....	111	Potassium guaiaretate	82
Pitayo bark	216	myronate	60, 61
Pitch Burgundy	376, 377	oleate	295
Pix Burgundica	376, 377	Pounce	373
liquida	378	Powder, Dover's ..	53
nigra	378	Prepared lard	460
Plaster, lead	292	storax.....	360
soap	292	suet.....	467
Plumbi acetatis unguentum ..	461	Propyl	454
carbonatis unguentum....	447	hydrate	454
emplastrum	294, 296	Propylamine	454
iodidi emplastrum	375	Protozoa	433, 434
c. opio pilula.....	53	derivation of	433
subacetatis unguentum			
compositum	152, 325		

	PAGE.
Provence oil	292
Prunes	153
Pruni Fructus	153
Virginianæ Cortex	153
Prunus domestica	153
laurocerasus	154
serotina	153
Pseud-aconitine	10
Psychotria emetica	222
Pterocarpi lignum	126
Pterocarpus draco	413
erinaceus	125
marsupium	124
santalinus	126
Puccoon, red.....	56
Pulp cassia	138
colocynth	165
Pulpa colocynthidis.....	165
Pulvis amygdalæ compositus,	
146, 149	
catechu compositus 67,	
125, 199, 329, 336	
cinnamomi compositus,	
328, 389, 391	
cretæ aromaticus, 329, 336,	
389, 394	
cretæ aromaticus c. opio..	53
cretæ c. opio.....	449
elaterii compositus	165
glycyrrhizæ compositus ..	120
ipecacuanhæ compositus	
53, 221	
jalapæ compositus ..	267, 391
kino compositus 54, 125, 329	
opii compositus 54, 117,	
179, 317, 391	
rhei compositus 312, 314, 391	
scammonii compositus 267,	
271, 391	

	PAGE.
Pulvis tragacanthæ compositus	
117, 146, 417	
Punica granatum.....	172
Purging cassia	137
Purified ox-bile.....	465, 466
Pyrethri radix	226
tinctura	227
Pyrethrin	226
Pyro-gallic acid	357, 360
Pyroxylin	74
Pyrus cydonia	161
Pyxidium	284
Pyxis	284

Q

Quassia wood.....	99, 100
Quassia extractum	100
infusum	100
lignum	99
tinctura	101
Quassin	99, 100
Quercetin	157
Querci-tannic acid	353
Quercitrin	157
Quercus cortex.....	353
decoctum	354
infectoria	354
pedunculata	353
Quill bark, yellow	201
Quinia	204
tests	204
Quiniæ pilula	160, 211
sulphas	202, 209
,, acid	210
tests	211
tinctura	212
,, ammoniata.....	212

	PAGE.
Quiniae vinum	212
Quince seeds.....	161
Quinic acid	206
Quinicine	206
Quinidia.....	205
Quinone.....	206, 244
Quinovic acid	206

R

Racine de Guimauve	72
Radicis simarubæ cortex	99
Radix aconiti	2, 3
actææ racemosæ	16
althææ	72, 78
aristolochiæ	350
armoraciæ	57
arnicæ	230
belladonnæ	276
calumbæ	21
cimifugæ	16
galangæ minoris	391
gentianæ	258
glycyrrhizæ	118
hellebori nigri	13
hemidesmi.....	263
hydrastis	18
inulæ	234
ipecacuanhæ	218
krameriæ	66
melampodii	13
pareiræ	24
podophylli	10
pyrethri	226
rhei.....	310
sanguinariæ	56
sassafras	332
scammoniæ	268

	PAGE.
Radix senegæ	64
serpentariæ	351
spigeliæ	256
sumbul	185
taraxaci	236
valerianæ	223
veratri albi.....	410
,, viridis	411
Raisins	98
Ranunculaceæ	1—18
Raphé.....	123
Rasores	456
Rectembryeæ	273
Red cinchona bark	214
litmus paper	424
puccoon	56
sandal wood	126
sanders wood	126
Reich, M.	220
Resina	373
common.....	84
special distinctive tests ..	374
guaiaci	81, 84
jalapæ	265
podophylli	12, 17
,, preparation of..	12
scammoniæ	268
Resinæ emplastrum.....	374
unguentum	374, 447
Rhœadin	31, 40
Rhamnaceæ	101—103
Rhamnegine	102
Rhamni fructus.....	101
succus	102
syrupus	103, 391
Rhamnine	102
Rhamnus cathartica	101, 102, 103
frangula.....	103

	PAGE.
Rhatania-tannic acid	66, 67
Rhatany, New Granada	67
Peruvian red	67
root	66
savanilla	67
Rhei extractum.....	313
infusum	313
pilula composita 109, 302,	
313, 402	
pulvis compositus 312, 314, 391	
radix	310
syrupus	184, 314
tinctura....	184, 314, 389, 394
vinum	64, 314
Rheo-tannic acid	312
Rheum officinale	310, 311
palmatum, var. Tanguti-	
cum	311
Rhaponticum	315
Rhizoma calami aromatici....	414
Rhœados petala	30
syrupus	31, 32
Rhizopoda	433
derivation of	433
Rhubarb root.....	310
Batavian	315
Dutch trimmed.....	315
East Indian	315
English	315
Rhus semialata.....	356
Richardsonia scabra	222
Ricini oleum	347
semina	346
Ricinolamide.....	348
Ricinoleic acid	348
Ricinolein	348
Ringed animals.....	435
Roccella.....	422

	PAGE.
Roccella fuciformis	423
tinctoria	423
Rodentia	468, 469
Root, alkanet	297
arnica	230
belladonna.....	276
Carolina pink	256
cimicifuga	16
elecampane	234, 235
green hellebore.....	411
hemidesmus	263
horseradish	57
Indian pink	256
ipecacuanha	218
liquorice.....	118
marshmallow	72, 78
pareira	24
pellitory	226
rhatany	66
rhubarb	310
sassafras	332
scammony.....	268
senega	30
serpentary.....	257, 351
sumbul	185
valerian	223, 224
white hellebore.....	410
Rosa canina	160
centifolia	158
Gallica	156
Rosaceæ	147—162
Rosamala storax	362
Rosæ aqua.....	159
caninæ confectio	160
,, fructus	160
centifoliæ petala.	158
Gallicæ confectio.....	157
,, petala	156

	PAGE.
Rosæ Gallicæ syrupus.....	158
„ infusum acidum	158
„ oleum	159
Rosæ	155—160
Rose attar	159
leaves	156
otto.....	159
Rosemary, oil of	304
Rosin	373
colophony	373
Rosmarini oleum	304
spiritus	305
Rosmarinus officinalis.....	304
Rottlera tinctoria.....	349
Rottlerin	350
Rubiaceæ	197—222
Rue, oil of	85
Rumex	36, 37
Ruminantia	459, 465, 467
Rusot	30
Russian musk	467
Ruta graveolens	85
Rutaceæ	84—90, 91
Rutæ oleum	85
Rutic acid	85
Rye, ergot of	425—428

S

Sabadilla	405
Sabadilline	411
Sabinæ cacumina.....	367
oleum	368
tinctura	368
unguentum	369, 447, 461
Saccharum lactis	465
purificatum	417
Saffron	393

	PAGE.
Saffron adulteration.....	395
cake	394
hay	394
Safrol	333
Sagapenum Levant.....	194
Persian	193
Sage (garden)	305
Sago	414
Sagus	414
St. Ignatius' beans	256
Salad oil	291
Salep	385
Salicaceæ	362—364
Salicin	363
Salicis cortex	362, 363
Salicylic acid	169
Salicylol.....	363
Salicylous acid	363
Saligenin	363
Salix alba	362
caprea	362
fragilis	362
Salvia officinalis	305
Sambuci aqua	197
flores	196
unguentum	197
Sambucus nigra	196
San Salvador	129
Sandal wood, red.....	126
Sandarach	373
Sanders wood, red	126
Sanguinaria	57
Canadensis	56
Sanguinariæ radix	56
Sanguis draconis	413
Sanguisorbæ	161
Sanguisuga medicinalis	436
officinalis	436

	PAGE.
Santalaceæ	352
Santali lignum	352
Santalic acid.....	126
Santalin.....	126
Santalum album	352
Santonica	232, 233
Santonin	233, 234
Sapindaceæ	76, 77
Sapo animalis	468
durus	295
mollis	295
Saponin.....	12, 65
Saponis emplastrum	295, 375
cerati emplastrum	293
pilula composita	53, 296
linimentum	296
„ compositum	305
Sapotaceæ	245—247
Sarothamnus scoparius	113
Sarsaparilla	383
Brazilian ..	383
Guatemala.....	383
Guayaquil	384
Honduras	383
Indian	263
mealy	383
Mexican.....	383
non-mealy	383
official	383
varieties	383
Sarsæ decoctum	382
compositum	81, 120, 333, 382
extractum liquid	382
Sassafras camphor	333
officinale	332, 333
radix	332
Sauropsida	455—457
derivation of	455

	PAGE.
Savanilla rhatany.....	67
Savin	367
oil of	368
Saxon oil of fennel	180
Scammoniæ radix	268
resina	268
Scammonii confectio	170, 179,
271, 391, 443	
mistura	269
pilula composita	266, 269
pulvis compositus	267, 271, 391
Scammonium	269, 270
Scammony, estimation	272
resin	268
root	268
Scilla	402
maritima	402
Scillæ acetum	404
oxymel	404, 443
pilula composita	193, 391, 404
syrupus	404
tinctura	404
Scillitin	403
Sclerotium.....	426
Scoparii cacumina	113
decoctum	115
succus.....	115
Scoparin	114
Scorpioid inflorescence	297
Scotch paregoric	55
Scrophulariaceæ	306—309
Secale cereale	425
Seed, black mustard	59, 60
colchicum	409
quince.....	161
stramonium	283
white mustard.....	59
Semen-contra	232

	PAGE.		PAGE.
Semina colchici	409	Serum sanguinis	456
cydoniæ	161	Sesame oil	294
fœni-græci	115	Setæ mucunæ	122
Ignatii	256	Sevum præparatum	467
lini	77	Sialogogue, derivation of	89
papaveris	34	Siamese benzoin	248, 249
ricini	346	Siberian musk	467
sinapis albæ	59	Siceæ	162
,, nigræ	59	Silvic acid	373
staphisagriæ	15	Simaruba amara	99
stramonii	283	root bark	99
theobromæ	74	Simarubaceæ	99—101
tigllii	344	Simarubæ radicis cortex	99
Senega root	30	Simplex unguentum	152, 461
Senegal gum	144	Sinalbin	61
Senegæ infusum	65	Sinapin, sulphate of	61
radix	64	Sinapis alba	59
tinctura	65	albæ semina	59
Senegin	65	cataplasma	62
Senna Alexandrian	133	charta	62, 246
Alexandrina	133	linimentum compositum	
Bombay	136	62, 322, 349	
confection of	341	nigra	59
East Indian	136	nigræ semina	59
Indica	136	oleum	62
Mecca	136	Sinigrin	58, 60, 61
Tinnivelly	136	Sirop de Gomme	146
Sennæ confectio 120, 134, 138,		Skulëin	403
140, 153, 184		Smilaceæ	379—384
infusum	135, 391	Smilacin	380
mistura composita ..	120, 135	Smilax medica	383
syrupus	135, 184	officinalis	379, 380, 383
tinctura	98, 135, 179, 184	papyracea	383
Serons	200	Smyrna opium	36
Serpentariæ infusum	352	Snake root, Virginian	351
radix	351	Soap curd	292
tinctura	352	hard	292, 295
Serpentary root	257, 351	liniment	292

	PAGE.
Soap plaster	292
soft	292, 295
Soaps	294
Socaloin	398
Socotrine aloes	398, 399
Sodium oleate	295
Solanaceæ	272—276, 297
Solanine	274
Solanum	273
dulcamara	273
Solazzi juice	120
Solenostemma argel	136
Solidungula	459
Solution albumen	457
gelatin	455
sulphate of indigo	121
Sori	418, 429
"Sorts" acacia	144
Spanish fly	448
juice	120
Sparteia	114
Spearmint, oil of	302
Special distinctive tests,	
ammoniacum	195
asafoetida	194
balsam tolu	128
bdellium	109
benzoin	249
Canada balsam	376
copaiba	141, 142
dragons' blood	413
elemi	105
galbanum	194
guaiaicum	83
mastich	111, 112
myrrh	108
olibanum	195
opoponax	195

	PAGE.
Special distinctive tests,	
parillin	381
resin	374
sagapenum	193, 194
spermaceti	458, 459
Spigelia marilandica	256
Spigeliæ radix	256
Spike, oil of	300
Spiritus ammoniæ aromaticus	96
,, foetidus	187, 188
armoraciæ compositus	58, 336
cajuputi	168
camphoræ	325
juniperi	367
lavandulæ	300
menthæ piperitæ	302
myristicæ	336
rosmarini	305
vini gallici mistura	457
Sponge	433
tents	434
Spongia officinalis	433
usta	434
Spongida	433
Sporangia	418
Spurge laurel	321
Spurious dissepiments	137
Squill	402
Staphisagriæ semina	15
Staphisagrine	16
Star anise fruit	20
Starch	416, 445
Starches, classification	473, 479
groups	479—485
mounting	473—476, 477
measurements	477, 478
observing	473—476
sampling	473—476

	PAGE.		PAGE.
Starches, table for detection		Succus mori	341
	481—485	rhamni	102
Stavesacre seeds	15	scoparii	115
fatty oil of	16	taraxaci	238
Stearin	76, 79, 292, 468	Sucrose	442
Stearopten	224	Suet prepared	467
otto	159	Sugar, cane	442, 444
Sterculiaceæ	74—76	fruit	442
Stoddart, W. W. (saffron)	394	grape	442
Storax prepared	360	of milk	465
rosamala	362	Sulphate of atropia	123, 280
Straits of Magellan	19	beberia	331, 332
Stramonii extractum	283	hyoscyamia	285
folia	282	indigo solution	121
semina	283	quinine	202
tinctura	284	sinapin	61
Stramonium leaves	282	Sulphindigotic acid	121
seeds	283	Sulphuris unguentum	461
Striated ipecacuanha	222	iodidi unguentum	461
Strychnia	252—254	Sulphocyanate acrinyl	61, 62
Strychniæ liquor	251, 255	allyl	60, 62
Strychnos Ignatia	256	Sumatra benzoin	248, 249
Nux vomica	251	Sumbul radix	185
,, bark	86	tinctura	186
Student's Museum Cabinet.		Sumbulamic acid	186
<i>Vide</i> Appendix II.		Sumbulic acid	186
Styraceæ	247—250	Sumbulolic acid	186
Styracin	361	Suppositoria acidi tannici	76,
Styrax benzoin	247		358, 447, 461
præparatus	360	,, c. sapone	358
Styrol	328, 361	hydrargyri	76, 447, 461
Suakin gum	144	morphiæ	45, 76, 447, 461
Succus belladonnæ	281	,, c. sapone	45
conii	177	plumbi composita	54, 76,
digitalis	309		447, 461
hyoscyami	286	Sureau, fleurs de	196
limonum	96	Sus scrofa	460
,, saturating power	97	Symphytum officinale	298

	PAGE.
Syrupus althææ	73
aurantii	92
„ floris	94
codeia.....	49
hemidesmi.....	263
limonis	95
mori	342
papaveris	34
rhœados	31, 32
rhamni	103, 391
rhei.....	184, 314
rosæ gallicæ	158
scillæ	404
sennæ.....	135, 184
tolutanus	128
zingiberis	390

T

Tabaci enema	288
folia	287

**Table for Detection of
Starches. *Vide Appen-
dix I.***

Talca gum.....	144
Tamarind	138
Tamarinds, East Indian	139
West Indian	139
Tamarindus	138
Indica.....	138
Tampico jalap	268
Tannic acid	356
Tannici acidi suppositoria	76
Taraxacerin	237
Taraxaci decoctum	237
extractum	238
radix	236
succus	238

	PAGE.
Taraxacin	237
Taraxacum	225
dens leonis	236
officinale	236
Tar, juniper	367
wood	378
Taurin	466
Taurocholic acid	465, 466
Tears, guaiacum	81
Tecomates.....	129
Teleostei	453—455
derivation of	453
Teneriffe cochineal	439
Tephrosia apollinea.....	136
Terebinthina Canadensis	375
Veneta	366
Terebinthinæ confectio 120,	
	370, 443
enema	371
linimentum	371
„ aceticum	371
„ oleum	369
unguentum 371, 375, 447, 461	
Terra japonica	197
Tests, cholic acid	466
cinchona alkaloids ..202, 203	
„ officinalis	214
oleum amygd. amar.....	151
olive oil	293
quinia.....	204
quinia sulphas	211
Tests, special distinctive,	
ammoniacum	192, 195
asafoetida	192, 194
balsam Canada.....	376
„ copaiba	141, 142
„ tolu	128
bdellium	109

	PAGE.
Tests, special distinctive,	
benzoin	249
dragon's blood	413
elemi	105
galbanum	192, 194
guaiacum	83
mastich	111, 112
myrrh	108
olibanum	195
opoponax	195
parillin	381
resin	374
sagapenum	193, 194
umbelliferous gum	
resins	193—195
Thalamifloræ	1—101
Thallogenæ	421—430
Thebaia, thebaine	40, 50
Theine	77
Theobroma cacao	75
Theobromæ oleum	75
semina	74
Theobromine	75
Thorn apple	282
Thus Americanum	375
Thyme (garden)	305
Thymelacææ	321—323
Thymol	305
Thymus vulgaris	305
Tigllii semina	344
Tiglinic acid	345
Tinctura aconiti	8
aloes	120, 401
arnicæ	231
asafœtidæ	189
aurantii	93
,, recentis	93
belladonnæ	282

	PAGE.
Tinctura benzoini composita	
128, 249, 362, 402	
buchu	88
calumbæ	24
camphoræ composita	54,
182, 250, 325	
cannabis Indicæ	339
cantharidis	451
capsici	276
cardamomi composita	98,
179, 329, 388, 439	
cascarillæ	344
castorei	469
catechu	199, 329
chiratzæ	262
cinchonæ composita	214,
352, 394, 439	
,, flavæ	208
cinnamomi	329
cocci	439
colchici semina	410
conii	176
croci	394
cubebæ	319
digitalis	309
ergotæ	428
gallæ	356
gentianæ composita	93,
260, 389	
guaiaci ammoniata	84
hyoscyami	287
jalapæ	267
kino	125
kramerizæ	67
laricis	365
lavandulæ composita	300,
305, 329, 336	
limonis	95

	PAGE.
Tinctura lobeliæ	242
„ ætherea	242
lupuli	340
myrrhæ	108
nucis vomicæ	255
opii	55
„ ammoniata 55, 182,	
250, 394	
pyrethri	227
quassiæ	101
quinia	212
„ ammoniata	212
rhei	184, 314, 389, 394
sabinæ	368
scilla	404
sennæ composita 98, 135,	
179, 184	
serpentaria	352
stramonii	284
sumbul	186
tolutana	128
valerianæ	224
„ ammoniata	224
veratri viridis	412
zingiberis	390
„ fortior	391
Tincture of bark (Huxham) ..	214
of capsicum	275
Fleming.....	8
of lavender (compound) ..	126
of litmus	424
of turmeric	387
Tinnivelly senna	136
Tinospora cordifolia	29
Tobacco.....	287
Tolene	127
Tolu balsam	126
Toluene	378

	PAGE.
Tolutana tinctura	128
Tolutanus syrupus	128
Tous les mois	392
Tragacantha	115
Tragacanthæ mucilago	117
pulvis compositus 117,	
146, 417	
Trigonella fœnum-græcum ..	115
Tritici farina.....	416
Triticum vulgare	416
Trinitro-cellulose	74
Trochisci acidi tannici	358
bismuthi.....	159
catechu	199
ipêcacuanhæ	221
morphiæ.....	45
et ipêcacuanhæ	45, 222
opii	55, 120
Tropic acid	280
Tropinia	280
Tubulifloræ	225
Tulasne (M)	425
Turkey gum	144
myrrh.....	106
Turmeric	386
African	392
paper	387
tincture of	387
Turpentine, American.....	369
Chian	111
horse	369
oil of	370
Venice	366
Tussilago	225

U

Ulmaceæ	336
Ulni cortex	336

	PAGE.
Ulmi decoctum	337
Ulmus campestris	336
Umbelliferone ..108, 109, 190,	
193, 194, 195	
Umbelliferæ.....	174—195
Umbelliferous gum resins,	
detection of	193
Uncaria acida	198
gambier	197
Undulated ipecacuanha	222
Unguentum aconitiæ	8, 461
atropiæ	280, 461
antimonii tartarati	447
belladonnæ	282, 461
cadmii iodidi.....	447
cantharidis	293, 447, 451
cetacei	152, 447, 459
creasoti	372, 447
elemi	106, 447
gallæ	356, 461
„ c. opio	56, 356, 461
hydrargyri.....	461, 468
„ ammoniati	447
„ compositum	
293, 325, 447	
„ iodidi rubri	447
„ nitratis....	293, 461
„ oxidi rubri	152, 447
„ subchloridi	461
iodi	461
pici liquidæ	378, 447
plumbi acetatis.....	461
„ carbonatis	447
„ iodidi.....	447
„ subacetatis comp.	
152, 325, 447	
potassæ sulphuratæ.....	461
potassii iodidi	461

	PAGE.
Unguentum resinæ	374, 447
sabinæ	369, 447, 461
sambuci	197
simplex	152, 461
sulphuris	461
„ iodidi	461
terebinthinæ	371, 375, 447, 461
veratriæ.....	293, 407, 461
zinci	461
Ungulata	459—468
Urginea scilla	402
Uvæ passæ	98
Uvæ ursi folia	243
„ infusum	244

V

Vaccinium vitis idœa	245
Valerene	224
Valerian root	223, 224
Valeriana officinalis	223
Valerianaceæ	222—224
Valerianæ infusum	224
radix	223
tinctura	224
„ ammoniata	224
Valerianic acid.....	186, 224
Valerol	224
Vanilla aromatica	385
planifolia	385
Vanillin	385
Vapor coniæ	177
creasoti	372
Varieties gamboge	70
opium commercial	36
sarsaparilla	383
Vegetable mercury	11
Venice turpentine	366

	PAGE.
Vera Cruz cochineal	439
Veratri albi radix	410
viridis radix	411
,, tinct.	412
Veratria	405—407
gallate of	407
Veratriæ unguentum	293, 407, 461
Veratrum album	410
viride	411
Verbascum thapsus	309
Vertebrata	452
Vinum aloes.....	389, 391, 401
aurantii ..	93
colchici	409
ipecacuanhæ	221
opii	56, 170, 329
quinia	212
rhei	64, 314
Virgin oil	291
scammony	270
Virginian snake root	351
Vitaceæ	98
Vitellus ovi	457
Vitis vinifera.....	98
Volatile oil of nutmeg.....	325

W

Wattle gum	144
Wax, white	445
yellow.....	445
West Indian tamarinds	139
Wheat flour	416
White of egg.....	456
hellebore root	410
mustard seed.....	59
pareira	27
pepper	317


	PAGE.
White wax	445, 446
Wild cinnamon.....	63
Willow bark	362
Wine of elder	197
ipecacuanha	221
Winteri cortex	19
Winter's bark.....	19, 20, 64
Wolfsbane	2
Wood, Brazil.....	132
guaiacum	80
quassia.....	99, 100
red sandal	126
red sanders	126
Wood oil	71
tar	378
Woody nightshade	273
Wool, cotton.....	74
Wormseed.....	232

Y

Yellow cinchona bark.....	200
pareira	27
root.....	18
wax.....	445, 446
Yolk of egg	457

Z

Zambesi (river).....	21
Zinci unguentum.....	461
Zingiber.....	389
officinale	389
Zingiberaceæ	386—391
Zingiberis syrupus	390
tinctura	390
,, fortior.....	391
Zygophyllaceæ	80—84



FOR BREAKFAST.

CHOCOLAT MENIER.

AWARDED NINETEEN PRIZE MEDALS.

THE HEALTHIEST, BEST, & MOST DELICIOUS
ALIMENT FOR BREAKFAST.

Annual Consumption exceeds 17,500,000 lbs.

THE QUALITY DEFIES ALL HONEST COMPETITION.

Sold in Half-Pound and Quarter-Pound Packets.

Medals Awarded to M. MENIER at the Philadelphia Centennial
Exhibition :

ONE MEDAL FOR THE CHOCOLATE MADE AT NOISIEL OF
MARNE.

ONE MEDAL FOR THE CHOCOLATE MADE AT THE LONDON
BRANCH FACTORY.

ONE MEDAL FOR THE SUGAR MADE AT ROYE, FRANCE.

ONE MEDAL FOR THE COCOAS GROWN AT VALLE MENIER
NICARAGUA, CENTRAL AMERICA.

WAREHOUSE—

SOUTHWARK STREET & WORCESTER STREET.

SOLD RETAIL EVERYWHERE.

ADVERTISEMENTS.

KNOWLES & PHILLIPS,

Wholesale Druggists,

47, MINORIES, LONDON,

MANUFACTURERS OF

*Pharmaceutical
Preparations.*

PURE ACIDS AND CHEMICALS
FOR ANALYSIS.

PRICE LISTS ON APPLICATION

HANDSOME CHEMIST'S SHOP FITTED COMPLETE,

INCLUDING GLASS SHOW CASES, GOLD
LABELLED SHOP BOTTLES, JARS, &c.,

For £75.

LISTS ON APPLICATION.

An experienced Gold Label Writer, Specie Jar Painter, Embosser, &c., engaged on the premises, and sent to all parts of the Kingdom. Every description of Ornamental Writing done, &c., cheaper than any other house.

LLOYD RAYNER,

Medical & General Shop Fitter, Shop Front
Builder, Show Case Maker,
Fixture Dealer, and Medical Labeller,

SHOW ROOMS AND OFFICES—

333, KINGSLAND ROAD.

FACTORIES—

2, DOWNHAM ROAD, KINGSLAND, & WHITMORE ROAD,
LONDON, N.

ADVERTISEMENTS.



PRIZE MEDAL,
PARIS EXHIBITION,
1867.



BURGOYNE, BURBIDGES,
CYRIAX & FARRIES,

Wholesale and Export Druggists,
MANUFACTURING & OPERATIVE CHEMISTS,
DISTILLERS OF ESSENTIAL OILS,
16, Coleman St., London, E.C.

MANUFACTURERS OF EVERY DESCRIPTION OF
Chemical, Pharmaceutical, Photographic, and
other preparations.

*Importers of Drugs and Chemicals used in the Arts and
Manufactures.*

Speciality:—Pure Acids, Chemicals, and Re-agents for
Analytical Purposes and Scientific Research.

Sole Agents for C. A. KAHLBAUM's Organic & Alcoholic Compounds.

PRICE LISTS AND SPECIAL QUOTATIONS ON APPLICATION.

ADVERTISEMENTS.



ESTABLISHED 1859.

The CHEMIST AND DRUGGIST is the indispensable companion of every intelligent and energetic pharmacist. It is everywhere recognised as the commercial and literary organ of British chemists and druggists, and it is used by the leading firms of Great Britain, America, France and Germany, as the medium of their announcements to the Anglo-Saxon trade.

SUBSCRIPTION :—

TEN SHILLINGS PER ANNUM

Including a copy of the Chemists' and Druggists' Diary at the end of the year, all Post Free.

The CHEMIST AND DRUGGIST contains a hundred pages or more monthly. It presents concisely all the important pharmaceutical items of the month, and contains a large number of original articles by the best writers. Among its special features may be mentioned its Trade Report, Corner for Students, Exchange Column, Foreign Correspondence, Notes of Novelties, &c. For business announcements it is an unequalled medium.

PUBLISHING OFFICE :—

COLONIAL BUILDINGS, 44a, CANNON STREET,
LONDON.

ADVERTISEMENTS.

MUTER'S "Pharmaceutical Chemistry,"

(THEORETICAL AND PRACTICAL.)

Undoubtedly the Best Student's Book. (See
Press Opinions.) 750 pp. Royal 8vo.

PRICE 15/-.

"The Analyst,"

A Monthly Journal of Analytical and Micro-
scopical Research, including the proceedings
of the Society of Public Analysts.

PRICE 6d.; OR 5/- PER ANNUM POST
FREE,

EDITED BY

G. W. WIGNER & JOHN MUTER.

Published by W. BAXTER, at the South London Central Public Laboratory,
Kennington Cross; and Sold by Messrs. BAILLIERE, TINDAL & COX, King
William Street, Strand; and SIMPKIN & MARSHALL, Stationers' Hall Court.

NEW LIST OF
Microscope Specialities
FORWARDED UPON APPLICATION.



MURRAY & HEATH'S

FIVE GUINEA

Student's Microscope,

With Two Achromatic Object Glasses, 1-in. of 15°
and ½-in. of 75°, of excellent quality.

The same form of Stand finished in a very superior
manner, with Coarse Rack and Fine Screw Adjust-
ments, Universal Movement to Stage, Plain and
Concave Mirrors, and Two Object Glasses, as above,
Price £7 7s.

THE STUDENT'S BINOCULAR MICROSCOPE,

With pair of Eye Pieces and Two Object
Glasses, as above, Price £10 15s.

The Pocket or Sea-side
Microscope, Clinical and
Chemical Thermometers,
Chemical, Physical and
Medical Apparatus, &c.

Murray & Heath,

Scientific Instrument
Makers to Her Majesty
and the Government
Departments,
69, JERMYN ST.,
LONDON, S.W.

ADVERTISEMENTS.

The London Medical Bottle Co.,

MANUFACTURERS OF EVERY DESCRIPTION OF

DISPENSING BOTTLES,

Suitable for Surgeons and Chemists.

WORKS:—24, CRANMER ROAD, NORTH BRIXTON, S.W.

WAREHOUSES:—MANOR PLACE, WALWORTH ROAD, S.E.

SLATTER & CO., Proprietors.

PRICE LISTS ON APPLICATION.

NOTICE.

MR. YOUNG begs to state that he was the
FIRST INVENTOR of the

ARNICATED FELT CORN & BUNION
PLASTERS.

Avoid counterfeit preparations,
and observe Trade Mark.

WHITE FELT
CORN & BUNION
PLASTERS

Have a world-wide reputation. They are
acknowledged to be the best in the Market.

FOR SHIPPING PRICE, APPLY TO

H. YOUNG, 16, Carthusian Street, Aldersgate Street, E.C.

May be ordered through the Wholesale Houses.

N.B.—BE CAREFUL THAT YOUNG'S ARE SUPPLIED.



YOUNG'S ARNICATED

ADVERTISEMENTS.

[TWELFTH EDITION.]

Materia Medica Examinations

BRITISH PHARMACOPŒIA, 1867.

*The Original Collection of Specimens of the Organic
Materia Medica, for the use of*

MEDICAL AND PHARMACEUTICAL STUDENTS.

IN order to assist Students in preparing for the above Examinations, we issue COLLECTIONS of carefully selected and characteristic Specimens of the Roots, Barks, Flowers, Fruits, Leaves, Resins, Gums, &c., of the British Pharmacopœia, 1867. Each Specimen bears a Label containing a description of the source from which the article is derived, its natural order, character, and tests, dose, and names of the Pharmacopœia Preparations into the composition of which it enters. The Student is thus enabled, in the most effective manner, to study the Materia Medica with the actual substance as well as the descriptive text before him.

*Testimonial from DR. MUTER, Director of the South-London School
of Pharmacy.*

"South-London School of Pharmacy, 11th April, 1878.

"Messrs. Southall Bros. and Barclay.

"Dear Sirs,—I have great pleasure in expressing the satisfaction which the specimens you provided for our Museum have given, and I also am always ready to say a word in favour of your excellent and economical Student's Collection, which I invariably recommend for the use of country Students.

"Yours faithfully, JOHN MUTER."

Testimonials have also been received from Dr. P. W. LATHAM, Professor of Medicine, Cambridge University; Dr. GEORGE F. DUFFEY, Fellow and Censor (*Examiner in Materia Medica*), King and Queen's College of Physicians, Ireland, Lecturer on Materia Medica in the Carmichael School of Medicine, Dublin, &c., &c.; PROFESSOR CHAS. B. C. TICHBORNE, Dublin, &c.; also from the EDITORS OF THE PHARMACEUTICAL AND MEDICAL JOURNALS, &c., &c.

Price of the Collection, including a copy of "The Organic Materia Medica," by W. SOUTHALL, F.L.S., in a neat Wooden Box, 30/- net.

(The Copyright of these Collections is Registered.)


CATALOGUES supplied FREE BY POST on Application.

SOUTHALL BROS. & BARCLAY,

Manufacturing Pharmaceutical Chemists. Birmingham.

THE SOUTH LONDON
SCHOOL OF PHARMACY,

Its Objects, Premises, Syllabus of Lectures,
Regulations and Terms.



BY WILLIAM BAXTER, SECRETARY.

THIRD EDITION—Revised to January, 1878.

London :

PUBLISHED AT THE OFFICES OF THE SCHOOL,
CENTRAL PUBLIC LABORATORY,
KENNINGTON CROSS, S.E.

[Entered at Stationers' Hall, All Rights Reserved.]

PREFACE.

IN issuing the present prospectus of the arrangements for the coming session, the management may be excused for referring with pride to the unequalled success attending the efforts of the school in raising the standard of Pharmaceutical education during the last ten years. No better proof of such success can be adduced than the fact that over 1500 persons have duly qualified themselves to practise as pharmacists from the ranks of its students. Much of this success is no doubt owing to the ability of the various lecturers, and especially to the fact that the direction of the Laboratory is entirely under the control of a teacher and analytical chemist, having the experience and high standing in his profession possessed by Dr. Muter. The school will continue to be devoted, as heretofore, to both classes of students, viz. Major and Minor; but no student is received who has not passed his Preliminary Examination. In addition, there has now also been added a department for the instruction of medical officers and others in Microscopy and General and Food Analysis.

I. Objects and Aims of the Institution.

AS now constituted, the South London School of Pharmacy receives two distinct classes of pupils, and provides a separate course of study to meet the requirements of each class, as follows :—

CLASS A.

In this division a long course of study, extending over twelve months, is provided for students who, not having previously attended many classes, desire to commence at the very root of matters, and be fully instructed in Pharmacy and its allied sciences. Provision is made for enabling them to become well acquainted with chemical analysis, both in its commercial bearing and as applied to the detection of adulteration in food and drugs, as well as with practical microscopy. Students commence by attending such of the lectures as they select daily, and the laboratory during three hours a day for the first three months, and thereafter they are allowed to work in the laboratory for the whole day. Leave of absence is given when required for any reasonable purpose. The age of entry in this department is optional, and certificates of attendance are granted. For medical officers and food analysts the lectures are not as a rule required; therefore the instruction is entirely practical, and the attendance optional, as may be arranged in each special case.

CLASS B.

For students who, having already obtained instruction in the country, or studied by self help, desire to have a shorter course of advanced instruction to enable them to systematize and arrange their ideas. In this class no student is received who has not attained an age of over twenty years. The length of each course is three months, but in the case of men already very nearly proficient special exceptions are occasionally made.

Students entering under this division are expected to strictly comply with the rules as to perfect regularity of attendance and good and studious conduct during their stay in London, Saturday being allowed as a half-holiday. They attend the lectures and laboratory daily. They are also subjected to frequent examination, written and oral, to ascertain their state of progress.

II. The School Premises.

Are situated at 325, Kennington Road, and consist of—

1. The necessary lecture rooms, properly fitted with abundance of diagrams, apparatus, etc.
2. Two complete museums of specimens of materia medica, with special cases presented by old students and others.
3. A museum of pharmaceutical preparations and dispensing department, upheld by Messrs. Knowles and Phillips.
4. An extensive herbarium and collection of manuscript prescriptions.
5. The laboratory (illustrated above), fitted in so complete and elegant a manner that each student has water, gas, fume chamber, and all such appliances within his reach, without leaving his bench. In addition to this, all the students face the demonstrators, and the benches are so arranged that the occupants can work sitting if desired. The table tops are of white enamelled slate, so that habits of exactitude and cleanliness can be well inculcated. All bottles and reagents are found by the school, except those mentioned in the laboratory list. Expensive apparatus of all kinds are lent to the students when required.

Besides the above premises specially for pharmaceutical students, there is the South London Central Public Laboratory, at Kennington Cross, where the secretary's office is situated, and in which the more advanced students are taught general and food analysis.

III. The Pharmaceutical Educational Course.

The session commences on the 15th of September, and lasts till the 20th of July; and the following courses of lectures, etc., are delivered several times during the session.

LATIN.

This consists of sixty lessons, and includes—

1. The reading of autograph prescriptions.
2. The parsing and rules of grammar involved in prescriptions.
3. The rendering into good Latin of prescriptions written in English.

Textbook :—Pereira's *Selecta e Prescriptis*.

BOTANY.

This involves a full course of sixty lectures on structural botany, and conveys an intimate knowledge of the root, stem, leaves, inflorescence, flower, fruit, and seed. The elementary structure of plants, such as cells, vessels, and the various tissues produced by their union, is fully discussed and illustrated by practical use of the microscope. A general knowledge of vegetable physiology is also inculcated. The organs of reproduction are practically taught by dissections conducted on fresh flowers once a week, under the direction of the lecturer.

During the summer months a lecture is delivered once a week, at Mitcham, on Mr. Bridger's herb farm.

No textbook required in this subject.

MATERIA MEDICA.

This course of sixty lectures opens with instruction in the classification of plants and the characters of the chief natural orders, illustrated by diagrams. It then proceeds to take the orders in detail, the portions of each plant used in Pharmacy, both official and non-official, being fully detailed. Their geographical sources, composition, modes of isolating their active principles, and adulterations, being fully described, as well as the preparations into which they enter. A general knowledge of the classification of animals, and a minute description

of the drugs, etc., obtained from the animal kingdom, is also given.

Textbook : Dr. Muter's *Organic Materia Medica*. At the School Offices, and of Messrs. Simpkin & Marshall, and Messrs. Balliere, Tindal & Cox).

PHARMACY.

This course lasts three months, and includes—

1. A description of the reasons involved in the processes used in pharmaceutical laboratories, and the best methods of conducting the same.
2. A minute knowledge of the B. P. and the chief non-official preparations usually sold.
3. The strengths, doses, and best modes of administration of the various official preparations.
4. The best methods of dispensing emulsions, pills, plasters, etc.
5. Practical instruction in dispensing and elegant pharmacy.

Textbook :—*The British Pharmacopœia*.

THEORETICAL CHEMISTRY.

This subject is taught on Dr. Muter's successful, because eminently systematic, "acidulous radical" method.

This course of sixty lectures includes—

1. The modern theories of chemistry.
2. The metals and chief basylous radicals.
3. Chlorine, chlorides, hypochlorites, chlorates.
4. Bromine, bromides, and bromates.
5. Iodine, iodides, and iodates.
6. Oxygen, hydrates and oxides of metals.
7. Alcohols and their derivatives, organic acids and their salts.
8. Sulphur, sulphides, sulphites, hyposulphites, sulphates, and sulphur substitution products.
9. Carbon, its oxides and carbonates.
10. Nitrogen, its oxides, acids, and their salts. Nitrogen substitution products.
11. Ammonia and its derivatives, amides, amines, and alkaloids.
12. Phosphorus, its oxides, acids, and their salts.
13. Arsenic, its oxides, acids, and their salts.

14. Hydrocarbons, sugars, glucosides, and other organic products not previously treated of.

Textbook : Dr. Muter's *Pharmaceutical Chemistry*. (At the School Offices, and of Messrs. Simpkin & Marshall, and Messrs. Balliere, Tindal & Cox.)

PRACTICAL CHEMISTRY.

I. JUNIOR COURSE.

(*Three months.*)

This includes—

1. The practical manufacture of the chief chemicals of the B. P., in which the processes are typical.
2. Qualitative analysis, and separation of bases and acids.
3. The practical taking of specific gravities, temperatures, and the measurement of gases.

II. SENIOR COURSE.

(*Six months.*)

This extends to—

1. Detection of poisons in organic mixtures.
2. Quantitative analysis of B. P. substances, gravimetric and volumetric.
3. Analysis of manures, water, minerals, etc.
4. Microscopic and chemical examination of food and drugs, for detecting adulterations.
5. Elementary organic analysis.

LIST OF APPARATUS REQUIRED BY EACH STUDENT.

test-tube stand, 8 holes and pegs.

1 dozen test-tubes.

1 test-tube brush.

1 nest of three beakers, from 10 oz. downwards.

3 flasks: 1 pint, $\frac{1}{2}$ pint, and 5 ounces.

2 Clarke's test-glasses.

3 funnels: $2\frac{1}{2}$ inch, 3 inch, and $1\frac{1}{2}$ inch.

2 porcelain crucibles.

1 5-oz. measuring glass.

4 evaporating basins: 3 inch, 4 inch, 6 inch, and 8 inch.

$\frac{1}{4}$ lb. glass rod.

$\frac{1}{4}$ lb. glass tubing.

1 foot small india tube for connecting tubing.

2 dozen corks, various.

1 spatula.

1 pair scissors.

1 three-cornered file.

1 round file.

1 blowpipe.

1 pair crucible tongs.

$\frac{1}{2}$ quire filtering paper.

platinum wire and foil.

1 piece blowpipe charcoal.

1 towel.

1 sponge.

$\frac{1}{4}$ oz. argentic nitrate.

1 dram platinic chloride.

small pestle and mortar.

The student must also find the alcohol, ether, and chloroform, which he may require.

IV. Medals and Prizes

Are awarded in DECEMBER, APRIL, and JULY, on the report of the VISITING EXAMINERS; the examinations being written and practical.

1. A BRONZE MEDAL in each of the subjects of CHEMISTRY (senior and junior), BOTANY, MATERIA MEDICA, and DISPENSING and PRACTICAL PHARMACY.
 2. To every student gaining more than two-thirds of the total possible marks, a CERTIFICATE of MERIT.
 3. The SILVER MEDAL of the Institution, competed for in July and open to all the Bronze Medallists of the year.
-

V. Fees.

1. PERPETUAL FEE for the whole course of lectures (including admission to the museum) £8 8s.

For example, a student entering the School, by payment of this fee, can continue to attend the lectures until the end of the session, and return *as often as he pleases for any number of following sessions without extra expense*, provided he also attends the laboratory.

2. For the Laboratory—

Junior Course, per month, £2 2s.

Senior Course, first month, £4 4s.; each succeeding month, £3 3s.

By these terms an industrious young man, who has paid attention to his business during his apprenticeship, may thoroughly complete his education as "Chemist and Druggist," so far as the school expenses are concerned, for a very moderate sum in comparison with the usual cost of a professional education. The cost of the junior laboratory instruction is exactly one half that usually charged at kindred institutions.

Old students who have already attended the school under the former regulations will be liberally dealt with.

VI. Conditions of Attendance.

- I. Unless from ill health, or by special permission, no student shall at any time be absent from his appointed classes, nor leave the premises during business hours. In cases of absence an explanatory note is required.
- II. After the ringing of the bell indicating a class, each student must be settled in his place and ready for work before the entry of the lecturer.
- III. Conversation during lecture, or conduct tending to attract the attention of other students from the subjects in hand, is forbidden.
- IV. Each student must provide himself with the necessary books recognised by the school, as well as with the small set of laboratory articles, and find his own sponge, towel, and apron.
- V. Except on Saturdays or holidays, students are expected to refrain from frequenting places of public entertainment, billiard rooms, or other such aids to the ruin of their professional prospects.
- VI. Students are required to register the address of their lodgings with the Secretary, and notify any change to him.
- VII. In event of any damage to institution property, should the perpetrator remain undiscovered for twelve hours, the whole class in the room at the time must subscribe to replace the article broken.
- VIII. All apparatus lent to students must be returned clean and in good order, or replaced at their own cost. On leaving for the day, students must clean and arrange their benches, and put away their apparatus in the cupboards.

These Rules are made with a sincere desire to aid the industrious student, and press hardly on no one. Retiring and smoking rooms, as well as an exercise yard, are provided within the walls, and everything is done to make the student comfortable while at school. Any violation of these conditions terminates the student's contract with the school.

SCIENTIFIC APPARATUS.

M. JACKSON & Co.,

(J. ORME & H. M. CAPNER,)

MANUFACTURERS OF EVERY DESCRIPTION OF

Apparatus for use in Teaching or Experimenting
in the Sciences of Chemistry, Electricity, Mag-
netism, Acoustics, Light, Heat, Hydrostatics,
Mechanics, &c.

Importers of finest Bohemian Glass, Berlin Porcelain,
Pure Chemicals for Analysis, &c.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.

CHEMICAL.



ARSENIC APPARATUS, Marsh's £ s. d.
0 6 0
BASINS, FOR EVAPORATING—

BERLIN PORCELAIN, finest quality, with Spout—							
No.	Inch diam.	Price each.	Price per doz.	No.	Inch diam.	Price each.	Price per doz.
00.....	2½	5d.	4/0	6.....	6	2/0	21/0
0.....	3	6d.	4/9	7.....	7	2/6	27/0
1.....	3½	9d.	8/3	8.....	8½	3 0	33 0
2.....	3½	10d.	9/0	9.....	10	5 0	54 0
3.....	4	1/0	10/6	10.....	12	7/6	82 0
4.....	4½	1/3	13/3	11.....	14	9/0	100 0
5.....	4½	1/8	18 0	12.....	15½	25/0	280 0

MEISSEN (Dresden) Porcelain, with Spout, shallow and very thin—
inch diam. 2½ 3½ 4½ 5 5½ 6½ 7½ 8½ 10 11 12 inch diam.
price 4d. 5d. 8d. 10d. 1/3 1/9 2/4 3/0 4/6 5/0 6/6 each.

M. JACKSON & CO.. 65, BARBICAN, LONDON.

ADVERTISEMENTS.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.



BASINS, FOR EVAPORATING—

GERMAN Porcelain, with Spout, glazed inside (*a cheap basin for ordinary use*).

No.	Inch diam.	Price each.	Price per doz.	No.	Inch diam.	Price each.	Price per doz.
1.....	2½	3d	2/6	6.....	5	7d	6 0
2.....	3	4d	3/3	7.....	5½	9d	8 0
3.....	3½	4d	3/6	8.....	6	1/0	10 0
4.....	4	5d	4/3	9.....	6½	1/3	12 0
5.....	4½	6d	5/0				

BEAKERS, for Hot Solutions, finest Bohemian Hard Glass, and of the best make—

No.	Ozs. cap.	Inch deep.	Price each.	Price doz.	No.	Ozs. cap.	Inch deep.	Price each.	Price doz.
00.....	1	1¼	2d	1/9	6..	23	5½	9d	8 0
0.....	1½	2	3d	2/9	7..	32	6	10d	9 0
1.....	3½	2½	4d	3/6	8..	48	7½	1/0	10 0
2.....	4½	3	5d	4/6	9..	67	8½	1/3	12 6
3.....	7	3½	6d	5/6	10..	90	9	1/6	15 6
4..	10	4	7d	6/3	11..	120	10	1/8	17 0
5..	16	4½	8d	7/3	12..	154	11	1/10	19 0



BEAKERS, similar to the above, nested—

Sets of	Nos.	Price per set.	Sets of	Nos.	Price per set.
3.....	00 to 1	7d	5.....	2 to 6	2/6
3.....	1, 3	1/0	6.....	0, 5	2 3
5.....	0, 4	1/8	8.....	1, 8	4 6
5.....	1, 5	2/0	12.....	1, 12	6 6

BLOWPIPES—

Berzelius', with Platinum Jet.....	each	£	s.	d.
Black's Japanned Tin, 9d. Brass do.	"	0	1	6
Hera path's for Bunsen's Burner	2/3, 2/6,	"	0	3 0
Do. on stand, complete	6/	"	0	7 6

BLOWPIPE TABLE, with Bellows, double action J & T (Hera path), with universal joint, for, above.....

BOTTLES, Bohemian white glass, stoppered—

ozs. cap.	½	1	2	4	6	8	12	16	20	40	ozs. cap
price N.M.	1/6	1/8	1/10	2/3	2/8	3/0	3 6	4/0	4 6	5 6	per doz.
" W.M.	1/8	2/3	2/8	3/0	3 8	4/0	4 6	5 3	5 9	7 0	"

BURETTES—


MOHR'S, with India-rubber Tube and Glass Jet:—

CC.	Divs.	Price each.	Grains.	Divs.	Price each.
20.....in.....	200.....	1/10	500.....	100.....	2 8
25.....,,.....	250.....	2 0	1,000.....	100.....	3 6
50.....,,.....	250.....	2/10	1,000.....	200.....	4 0
50.....,,.....	500.....	3 2	Clips for the above, 4d. ea. extra		

MOHR'S, with Glass Stopcock:—

50 cc. in 250 divs.	5/0	50 cc. in 500 divs.	0 5 6
1,000 grains in 100 divs.	5/6	1,000 grains in 200 divs.	0 6 6


ERDMAN'S FLOATS



M. JACKSON & CO., 65, BARBICAN, LONDON.

ADVERTISEMENTS.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.



BURETTE STANDS—

Teak wood for one, 2/0; for two.....	0	2	6
Black wood, with clamp for one Burette, 3/0; for two	0	4	6
Mahogany do. do. 4/6; for two.....	0	6	6
Do. with Brass Rod and Screw Clip (Dr. Percy's)	0	7	6

CONDENSERS, Liebig's Glass.....	2/6	and	0	3	6
Do. Copper Body, Japanned.....	12/0,	15/0,	and	1	1

CRUCIBLES AND COVERS—

MEISSEN (Dresden) Porcelain—							
Nos.	Oz. cap.	Price each.	Price per doz.	Nos.	Oz. cap.	Price each.	Price per doz.
9.....	1	4d.	3 9	4.....	2 1/2	10d.	9 0
8.....	1	5d.	4 6	3.....	3 1/2	1/0	10 6
7.....	1	6d.	5 6	2.....	5	1/2	12 6
6.....	1	7d.	6 6	1.....	8	1 6	16 0
5.....	1 1/4	8d.	7 0				

DRYING APPARATUS—

HOT AIR OVENS, square copper—				
inches square	6	7	8	9 10 inches square.
price	18/0	21/0	24/0	30/0 33/0 each.

HOT WATER OVENS, square, copper—				
inches square	6	7	8	9 inches square.
price	18/6	24/0	32/0	38/0 each.

DESICCATOR (as used at the Science Schools, Kensington.....)		£	s.	d.
		0	8	6

FILTER PAPER—

DUTCH, very thick and coarse.....	per quire	0	2	6
ENGLISH, best white.....	"	0	1	0
RHENISH, nearly equal in quality to Swedish	"	0	2	0

FILTERS, CUT (Circular), in packets of 100—

in. diam.	2 1/4	2 3/4	3 1/2	4 1/2	5 1/2	7 1/2	in. diam.
ENGLISH	2d.	4d.	5d.	6d.	9d.	1/0	per packet.
RHENISH	3d.	4d.	8d.	9d.	1/0	1/6	"

FLASKS, finest BOHEMIAN Hard Glass, for Boiling—

Ozs.	Price cap.	Price each.	Price per doz.	Ozs.	Price cap.	Price each.	Price per doz.
1.....	3d.	2/9	20.....	9d.	8/0
2.....	3d.	3/0	24.....	10d.	9/0
4.....	4d.	3 9	32.....	11d.	10/0
6.....	5d.	4 6	48.....	1/1.	11/7
8.....	6d.	5 6	64.....	1/2.	12/6
12.....	7d.	6 6	100.....	1/6.	16/0
16.....	8d.	7 3	160.....	2/0.	21/6

GERMAN—ozs.		2	4	8	12	16	24	32 ozs. cap.
price	2/0	2/3	3/0	4/0	5/0	6/0	7/0	per doz.

GAUGED and marked in neck, Grains—

to contain	500	1,000	5,000	10,000 grains.
price plain	9d.	1/0	1/6	1/9 each.
price stoppered	—	1/3	1/9	2/0 "

Do., Cubic Centimetres—

to contain	5	10	25	50	100	250	500	1,000 c.c.
price plain	3d.	4d.	6d.	7d.	9d.	1/2	1/4	1/9 each.
price stoppered	—	—	8d.	9d.	1/2	1/9	2/0	2/6 "

M. JACKSON & CO., 65, BARBICAN, LONDON.

ADVERTISEMENTS.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.

FUNNELS, BOHEMIAN GLASS, plain or ribbed, ground edges—



Inch diam.	Price each.	Price per doz.	Inch diam.	Price each.	Price per doz.
1	3d.	2/6	5	11d.	10/0
1½	3d.	2/9	6	1½	13/0
2	4d.	3/9	7	1/6	16/6
2½	5d.	4/6	8	2/0	22/0
3	6d.	5/6	9	2/6	27/0
3½	7d.	6/3	10	3/0	34/0
4	8d.	7/0			

ENGLISH GLASS, plain or ribbed—

in. diam.	1	1½	2	2½	3	4	5	6	in. diam.
price	1/6	1/9	2/0	2/6	3/0	4/0	6/0	9/0	per doz.

	per lb.	£	s.	d.
GLASS TUBING, SOFT, for bending, all sizes	„	0	1	0
Do. do. HARD, for combustions, finest	„	0	1	0
Bohemian	„	0	1	4
Do. do. Barometer and Thermometer	„	0	3	0

GRADUATED MEASURES, Glass—

1	2	drms.	1	2	4	6	8	10	16	20	40	ozs.
price	9d.	1/0	8d.	9d.	1/0	1/3	1/6	1/9	2/0	2/3	2/6	4/0 each.

HYDROMETERS, &c.—

	£	s.	d.
Argentometer, with Immersion Tube	each	1/6	and
Beaume's, for light or heavy Fluids	each	1/6	and
Lactometer, complete, 1/6 and 2/3	Ivory Scale	0	3 0
Saccharometer, 2/6	Ivory Scale	0	3 0
Salinometer for Aquarium, 1/6; for Sea Water	„	0	2 6
Sikes', with Weights, Thermometer, and Tables complete	„	3	3 0
Specific Gravity, for Acids, &c., 1,000—1,850	„	0	2 6
Do. for Spirits, 700—1,000	„	0	2 6
Do. for Spirits, from 40° under to 60° over proof, 2/6	Ivory Scale	0	3 0

TWADDEL'S, Nos. 1 to 6, each 1/6.	per doz.	0	15 0
Urinometer, Ivory Scale, in Leather Case	„	0	3 0
Do. with graduated Trial Jar in ditto	„	0	4 6



HYDROMETER IMMERSION TUBES—

in. high	6	8	10	12	14	in. high.
price	8d.	10d.	1/0	1/3	1/6	each.
per doz.	7/6	9/0	11/0	13/6	16/0	per doz.

LAMPS, BERZELIUS', Brass, on Stand, with Rings, &c.

GAS, Bunsen's	1/3, 2/3, and	0	3 6
Rose for ditto	1/0, 1/3, 1/9, and	0	2 0
Star support for ditto, with Chimney	6d. 8d. and	0	10 0

SPIRIT, GLASS, with Stoneware or Brass Tube wick Holder—

ozs. cap.	1	2	4	8	ozs. cap.
price	9d.	10d.	1/0	1/3	each.
with Screw Cap	1/3	1/6	1/9	2/0	



M. JACKSON & CO., 65, BARBICAN, LONDON.

ADVERTISEMENTS.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.



MORTARS and PESTLES—

WEDGWOOD Ware, with mark, "Wedgwood"—

Inch diam.	Price each.	Inch diam.	Price each.	Inch diam.	Price each.
3	1/3	5	2/6	8	5/6
3½	1/4	6½	3/6	9	7/6
4	1/9	7	4/6	10	9/0

PIPETTES, Bulb, 2d. 3d. and 4d.; per doz. 1/6, 2/0 and 0 2 6
Cylindrical, 3d. 4d. and 6d. per doz. 2/6, 3/0, and 0 4 0

GAUGED to deliver—Cubic Centimetres—

c.c.	1	5	10	20	25	50	100 c.c.
price	4d.	5d.	6d.	8d.	9d.	1/0	1/3 each.
per doz.	3/6	4/6	5/3	7/0	8/0	10/0	13/0 per doz.

RETORTS, finest BOHEMIAN hard glass—

Ozs.	Plain.	Tubulated.	Stoppered.
cap.	Each.	Per doz.	Each Per doz.
2	3d.	2/9	4d. 3/9 9d. 8/6
4	4d.	3/9	5d. 4/10 10d. 9/6
8	5d.	4/6	6d. 6/0 1/0 11/0
12	6d.	5/9	7d. 6/6 1/1 12/0
16	7d.	6/6	8d. 7/2 1/2 13/0
24	8d.	7/6	9d. 8/0 1/4 14/0
32	9d.	8/0	10d. 9/0 1/6 16/0
48	1/0	10/6	1/3 13/6 2/3 24/0

Tubulated for MANUFACTURING PURPOSES—

lbs. cap.	8	10	14	18	24	36	48 lbs. cap.
price	2/3	2/9	3/0	3/6	4/6	5/6	8/0 each.

RECEIVERS for above at the same prices as retorts.

RETORT STANDS, IRON, with three Brass Rings—

Length of Rod, inch	13	15	17	20	24 inch.
price	2/0	2/6	3/6	4/6	8/0 each.

SPATULAS—

Best Steel—inch	3	4	5	6	7 inch blade.
price	7d.	8d.	10d.	1/0	1/3 each.

Bone, 1/0; Ivory, 2/6; Platinum, at per oz. 1 12 0

STILLS, Copper or Tin, with pure Tin Worm Condensers—

	1 pt.	1 qt.	½	1	2	4 galls.
COPPER, price	11/6	15/0	21/0	30/0	45/0	80/0 each.
TIN.....price	7/6	10/0	12/6	15/0	24/0	35/0 each.

PHILLIPS' Revenue Standard, accurately made, as supplied to the Laboratories of Her Majesty's Customs 3 10 0


TEST PAPERS, Blue and Red Litmus, Turmeric, Lead, Starch, &c., book, 2d; per doz. 1/3, 1/4, 1/6 and 2/0

TEST TUBE CLEANERS, or Brushes, 2d.; per doz. 1/6

TEST TUBE HOLDERS, for the hand, 6d. & 1/0

TEST TUBE STANDS, Teak Wood—

holes	6	8	12	24 holes.
price	7d.	9d.	1/2	2/0 each.
do. with Pegs	1/0	1/3	2/0	3/6 each.



M. JACKSON & CO., 65, BARBICAN, LONDON.

ADVERTISEMENTS.

EXTRACTS FROM M. JACKSON & CO.'S PRICE LISTS.

TEST TUBES, finest German Glass—

Size.	Price	Size.	Price
Inches.	Per doz. Per gross.	Inches.	Per doz. Per gross.
3 × 3. 4l.	3/3	6 × 3. 10 1/4.	9/3
4 × 3. 6d.	5/3	6 × 3. 11d.	10/0
5 × 3. 8d.	7/3	6 × 1. 1 1/2.	12/6
5 × 3. 8d.	7/3	7 × 1 1/4. 1/6.	16/0
5 × 3. 9d.	8/3	8 × 1 1/2. 2/0.	21/0

THERMOMETERS—

	£	s.	d.
Box Wood Scale, 1/0 and 1/6; jointed, to 300° F.	0	3	6
For Baths, Japanned Tin Scale.	0	1	9
Chemical, to pass through corks—			

	400° F.	600° F.	300° C.	
Paper Scale.	2/6	3/0	3/0	price each.
Milk Scale.	3/6	4/0	4/0	"
Grad. on stem with enamelled back .. }	5/0	6/0	5/6	"

TUBES, HOFFMAN'S—

	£	s.	d.
V form, for the electrolysis of H ₂ O, HCl or NH ₃	0	5	0
Stand for do., 5/0; complete.	0	10	0
U form, to show the volume of H in HCl, with one Stopcock, 5/6; with stand.	0	13	0
Do., to show that 3 vols H combine with 1 vol. N to form NH ₃ , each, 6/6 and 7/6; complete 14/0 and ..	0	16	0
Do. for the Decomposition of Water, 7/0; complete ..	0	14	0
Do. for the Decomposition of HCl, 10/6; complete.	0	18	0
Do. to show that when a mixture of 2 vols. H and 1 vol. O are exploded, the 3 vols. of Gas become 2 vols., with Steam Jacket, 8/6; stand, 6/6; complete	0	15	0
Do. Eudiometer, graduated, 12/6; stand, 6/6; complete	0	19	0

TUBING, SOFT GLASS, for bending, all sizes, per lb. 0 1 0

Do. HARD GLASS, for combustions, finest Bohemian, per lb.	0	1	4
---	---	---	---

INDIA-RUBBER, Grey

inside diam. 1/8 3-16 1/4 3/8 1/2 5/8 3/4 1 in. diam.	
price 3d. 4d. 5d. 7d. 9d. 1/0 1/4 2/2 per foot.	
Black, 1d. per foot extra.	

WATCH GLASSES—

inch 1 1/2 1 3/4 2 2 1/2 3 inch diam.	
price 9d. 1/0 1/3 3/0 5/0 per doz.	

WOULFFE'S BOTTLES—

Ozs.	Price each	Ozs.	Price each
cap. 2 necks 3 necks		cap. 2 necks 3 necks	
8 1/0 1/3		64 2 6 3 0	
16 1/3 1/6		100 4 0 4 6	
32 1/9 2/0		130 5/6 6/0	
48 2/3 2/6		160 6/3 7/3	

The price per dozen applies strictly to orders for not less than one dozen.

M. JACKSON & CO., 65, BARBICAN, LONDON.

ADVERTISEMENTS.



ESTABLISHED TWENTY YEARS.

JOHN DAVIS,
Medical and Commercial Printer,
DESIGNER,
Lithographer, Engraver, etc., etc.,
STEAM WORKS:
OLD KENT ROAD, LONDON.

IN soliciting the attention of Members of the Medical Profession and others, J. D. begs to inform them that his Establishment is replete with every requisite for the rapid and economical execution of High-class Printing. It contains Machinery from the first makers of the world, and large founts of Type suitable for every class of work, from extensive volumes on Chemistry, Materia Medica and Scientific Subjects, to the smallest requirement of the retail Chemist.

Illustrated Catalogues; Prices Current; Lists of Chemicals, Scientific Apparatus, &c., &c.; executed in first-style at moderate charges.

ESTIMATES ON APPLICATION.

ADVERTISEMENTS.

LAXORA.

Professional Chemists, and Studying Men generally, who frequently suffer all the miseries of Constipation and its attendant discomforts, will be glad to learn that by using the above preparation, they can secure all the effects of the well-known *Electuaire lenitif* of the French Codex in an exceedingly palatable form. One trial will ensure continuance in its use.

N.B.—None genuine without signature P. GUYOT on each box.

SOLD BY ALL CHEMISTS AND DRUGGISTS.

In boxes at 1s. 1½d. each. Forwarded on receipt of 1s. 3d. in Stamps.

WHOLESALE—82, SOUTHWARK STREET, LONDON, S.E.

RIGOLLOT'S MUSTARD LEAVES.

Sold in Tins, 1/6 and 1/-; 2/6 Packets; also in Tins containing 100 leaves for dispensing.

DR. LELIEVRE'S ICELAND MOSS POULTICE (PATENTED)

Is exempt from all the inconveniences inseparable from poultices made with linseed or bread. These spoil the linen, bedding, &c., dry up on the edges, rapidly ferment, giving a most unpleasant odour, and no dependence can be placed on the quality and freshness of the linseed meal.

This new Poultice is instantaneous, for it is ready in a few seconds.

SOLD RETAIL BY ALL CHEMISTS.

Wholesale—82, SOUTHWARK STREET, LONDON.

